

**LEADER MILK OF MAGNESIA- magnesium hydroxide suspension**  
**Cardinal Health 110, LLC. dba Leader**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Cardinal Health Milk of Magnesia Drug Facts**

**Active ingredient (in each 15 mL)**

Magnesium hydroxide 1200 mg

**Purpose**

Saline laxative

**Uses**

- relieves occasional constipation (irregularity)
- generally produces bowel movement in  $\frac{1}{2}$  to 6 hours

**Warnings**

**Ask a doctor before use if you have**

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

**Ask a doctor or pharmacist before use if you are**

taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

**Stop use and ask a doctor if**

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

## Directions

- do not exceed the maximum recommended daily dose in a 24 hour period
- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- mL = milliliter

adults and children 12 years and older	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

## Other information

- **each 15 mL contains:** calcium 20 mg and magnesium 595 mg
- does not meet USP requirements for preservative effectiveness
- store at 20°-25°C (68°-77°F)
- close cap tightly after use
- avoid freezing

## Inactive ingredient

purified water

## Questions or comments?

**1-800-719-9260**

## Principal Display Panel

Genuine

Milk of Magnesia

Magnesium Hydroxide, 1200 mg per 15 mL

Saline Laxative

Original Flavor

Stimulant-Free

Comfortable, Cramp-Free Relief

Sugar-Free

COMPARE TO PHILLIP'S® MILK OF MAGNESIA active ingredient

100% Money Back Guarantee

12 FL OZ (355 mL)



<b>Drug Facts</b> <small>DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING.</small>	
<b>Active ingredient (in each 15 mL)</b> Magnesium hydroxide 1200 mg.....	<b>Purpose</b> Saline laxative
<b>Uses</b> ■ relieves occasional constipation (irregularity) ■ generally produces bowel movement in ½ to 6 hours	
<b>Warnings</b> Ask a doctor before use if you have ■ kidney disease ■ a magnesium-restricted diet ■ stomach pain, nausea, or vomiting ■ a sudden change in bowel habits that lasts more than 2 weeks Ask a doctor or pharmacist before use if you are taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work. Stop use and ask a doctor if ■ you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition. ■ you need to use a laxative for more than 1 week If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).	
<b>Directions</b> ■ do not exceed the maximum recommended daily dose in a 24 hour period ■ shake well before use ■ dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor ■ drink a full glass (8 oz) of liquid with each dose ■ for accurate dosing, use dose cup provided ■ mL = milliliter	
adults and children 12 years and older	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor
<b>Other information</b> ■ each 15 mL contains: calcium 20 mg and magnesium 595 mg ■ does not meet USP requirements for preservative effectiveness ■ store at 20°-25°C (68°-77°F) ■ close cap tightly after use ■ avoid freezing	
<b>Inactive ingredient</b> purified water	
<b>Questions or comments?</b> 1-800-719-9260	

\*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Phillips'® Milk of Magnesia.  
DISTRIBUTED BY CARDINAL HEALTH, DUBLIN, OHIO 43017

www.myleader.com 1-800-200-6313

© 2019 Cardinal Health. All Rights Reserved. CARDINAL HEALTH, the Cardinal Health LOGO, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners.

**100% Money Back Guarantee**  
Return to place of purchase if not satisfied.

CIN 5521554 REV. 11/19



## LEADER MILK OF MAGNESIA

magnesium hydroxide suspension

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:70000-0027
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>MAGNESIUM HYDROXIDE</b> (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	1200 mg in 15 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	

**Product Characteristics**

<b>Color</b>	WHITE (opaque)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0027-1	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/29/2020	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	01/29/2020	

**Labeler** - Cardinal Health 110, LLC. dba Leader (063997360)

Revised: 6/2021

Cardinal Health 110, LLC. dba Leader