

**CHLORAPREP ONE-STEP- chlorhexidine gluconate and isopropyl alcohol solution**  
**CareFusion 213, LLC**

-----  
**BD Chloraprep™ Scrub Teal™ 26 mL Applicator with Sterile Solution**

**WARNING**  
**FLAMMABLE**

**Keep away from fire or flame.**

**To reduce the risk of fire, PREP CAREFULLY:**

- do not use 26-mL applicator for head and neck surgery or on an area smaller than **8.4 in. x 8.4 in.** Use a smaller applicator instead.

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to

dry. **Wet hair is flammable.**

- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (**minimum of 3 minutes on hairless skin; up to 1**

**hour in hair)**

- do not allow solution to pool
- remove wet materials from prep area

***Active ingredients***

Chlorhexidine gluconate 2% w/v

Isopropyl alcohol 70% v/v

***Purposes***

Antiseptic

Antiseptic

**Use**

for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

***Warnings***

**For external use only. Flammable, keep away from fire or flame.**

**To reduce risk of fire, PREP CAREFULLY:**

- do not use 26-mL applicator for head and neck surgery
- do not use on an area smaller than **8.4 in. x 8.4 in.** Use a smaller applicator instead.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry.

**Wet hair is flammable.**

- do not drape or use ignition source (e.g., cautery, laser) until solution is

completely dry (**minimum of 3 minutes on hairless skin; up to 1 hour in hair**)

- do not allow solution to pool
- remove wet materials from prep area

**Allergy alert:**

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing • shock • facial swelling • hives • rash

If an allergic reaction occurs, stop use and seek medical help right away.

**Do Not Use**

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

**When using this product**

keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

**Stop use and ask a doctor if**

irritation, sensitization, or allergic reaction occurs.

These may be signs of a serious condition.

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately **13.2 in. x 13.2 in. (1126 cm<sup>2</sup>)**. Do not use 26-mL applicator for area smaller than **8.4 in. x 8.4 in.** Use a smaller applicator instead.
- do not use 26-mL applicator for head and neck surgery
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wing **only once** to activate the ampules and release the antiseptic
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- **allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)**. Do not blot or wipe away
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

### **Other Information**

- store between 15-30 °C (59-86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)
- the tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired.

### ***Inactive Ingredients***

FD&C green #3 dye, USP purified water

### ***Questions?***

[www.chloraprep.com](http://www.chloraprep.com) • **call 1-800-523-0502** (M-F 8 a.m.-5 p.m. CST)

**BD ChloroPrep™ Scrub Teal™ 26 mL Applicator Package Label**

REF 930825  
NDC 54365-400-39



# BD ChloroPrep™ Scrub Teal™

## 26 mL Applicator

2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v Isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation

Sterile Solution

Applicator is sterile if package is intact

Scrub Teal™

### **WARNING** **FLAMMABLE**



Keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- do not use 26-mL applicator for head and neck surgery or on an area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (**minimum of 3 minutes on hairless skin; up to 1 hour in hair**)
- do not allow solution to pool
- remove wet materials from prep area

#### Drug Facts

##### Active Ingredients

Chlorhexidine gluconate 2% w/v..... Antiseptic  
Isopropyl alcohol 70% v/v..... Antiseptic

##### Purposes

**Use** for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

##### Warnings

For external use only. Flammable, keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery
- do not use on an area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (**minimum of 3 minutes on hairless skin; up to 1 hour in hair**)
- do not allow solution to pool
- remove wet materials from prep area

##### Allergy alert:

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing - chest - facial swelling - hives - rash

• wheezing/difficulty breathing • shock • facial swelling • hives • rash

If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use** • on patients allergic to chlorhexidine gluconate or any other ingredient in this product • for lumbar puncture or in contact with the meninges • on open skin wounds or as a general skin cleanser

**When using this product** keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

**Stop use and ask a doctor** if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

### Directions

• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. • use in a well ventilated area • maximal treatment area for one applicator is approximately 13.2 in. x 13.2 in. (1126 cm<sup>2</sup>). Do not use 26-ml applicator for area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead. • do not use 26-ml applicator for head and neck surgery • remove applicator from package; do not touch sponge • hold the applicator with the sponge down. Pinch wing **only once** to activate the ampules and release the antiseptic. • wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin • completely wet the treatment area with antiseptic • **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds • **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes • do not allow solution to pool; tuck prep towels to absorb solution, and then remove • **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away. • discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

**Other information** • store between 15–30 °C (59–86 °F) • avoid freezing and excessive heat above 40 °C (104 °F) • the tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired.

**Inactive ingredients** FD&C green #3 dye, USP purified water

### Questions?

• [www.chloraprep.com](http://www.chloraprep.com) • call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion 213, LLC, El Paso, TX 79912, subsidiary of Becton, Dickinson and Co.

Artwork No. 6-930842 • A050619 • Made in the USA of US and imported parts • [www.bd.com](http://www.bd.com)



(01)00354365400396

External Use Only  
Professional Use Only



STERILE EO



Package Label

BD Chloraprep™ Scrub Teal™

26 mL Applicator

2% w/v chlorhexidine gluconate (CHG)

and 70% v/v Isopropyl alcohol (IPA)

Patient Preoperative Skin Preparation

Sterile Solution

Applicator is sterile if package is intact

REF 930825

NDC 54365-400-39

## CHLORAPREP ONE-STEP

chlorhexidine gluconate and isopropyl alcohol solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:54365-400
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CHLORHEXIDINE GLUCONATE</b> (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)	CHLORHEXIDINE GLUCONATE	20 mg in 1 mL
<b>ISOPROPYL ALCOHOL</b> (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	0.7 mL in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>FD&amp;C GREEN NO. 3</b> (UNII: 3P3ONR6O1S)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54365-400-39	1 in 1 POUCH	05/24/2019	
1		26 mL in 1 APPLICATOR; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020832	05/24/2019	

**Labeler** - CareFusion 213, LLC (826496312)

**Registrant** - Becton, Dickinson and Company (832696038)

# Establishment

Name	Address	ID/FEI	Business Operations
CareFusion 213, LLC		826496312	analysis(54365-400) , label(54365-400) , manufacture(54365-400) , pack(54365-400) , sterilize(54365-400)

Revised: 10/2025

CareFusion 213, LLC