

4240 FIRST AID KIT - 4240 first aid kit
Honeywell Safety Products USA, INC

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).

4240 First Aid Kit (1st aid Sp, EW, ASA, aypanal, miralac, amm. Inh, triple, burn spray WS, alcohol wipe, BZK wipes - 68400LNODEA)

Eyewash

Active ingredient

Sterile Water 99%

Eyewash

Purpose

Eyewash

Eyewash

Uses

- for flushing the eye to remove loose foreign material, air pollutants or chlorinated water

Eyewash

Warnings

For external use only Obtain immediate medical treatment for all open wounds in or near eyes. To avoid contamination, do not touch tip of container to any surface. Do not reuse. Once opened, discard.

Do not use

- if solution changes color or becomes cloudy
- if you have open wounds in or near the eyes, get medical help right away.

Stop use and ask a doctor if

- you experience eye pain
- changes in vision
- continued redness or irritation of the eye
- condition worsens or persists

Keep out of reach of children

- If swallowed, get medical help or contact a Poison Control Center right away.

Eyewash

Directions

- remove contacts before using
- twist top to remove
- flush the affected area as needed
- control rate of flow by pressure on the bottle
- if necessary, continue flushing with emergency eyewash or shower

Eyewash

Inactive ingredients

sodium chloride, sodium phosphate dibasic, sodium phosphate monobasic

Eyewash

Questions

1-800-430-5490

Aspirin

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)* *nonsteroidal anti-inflammatory drug

Aspirin

Purpose

Pain reliever/fever reducer

Aspirin

Uses

temporarily reduces fever and relieves minor aches and pains associated with:

- a cold
- headache
- toothache
- muscular aches
- backache
- minor pain of arthritis
- premenstrual and menstrual periods

Aspirin

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:are:

- age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- ringing in the ears or loss of hearing occurs
- any new symptoms appear

If pregnant or breast-feeding,

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last three months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

- In case of overdose, get medical help or contact Poison Control Center right away.

Aspirin**Directions**

- drink a full glass of water with each dose
- adults and children 12 years of age and older: take 1 or 2 tablets every 4 hours while symptoms last, not more than 12 tablets in 24 hours
- children under 12 years of age: consult a doctor

Aspirin**Other information**

- store at room temperature 15° - 30°C (59° - 86°F)
- TAMPER EVIDENT PACKETS
- DO NOT USE IF OPEN OR TORN

Aspirin***Inactive ingredients***

corn starch, croscarmellose sodium*, hypromellose*, microcrystalline cellulose*, mineral oil*, polyethylene glycol*, povidone, propylene glycol, silicon dioxide, stearic acid*, titanium dioxide*

*may contain these ingredients

Aspirin***Questions or Comments***

1-800-430-5490

Ammonia***Active ingredient***

Ammonia 15%

Ammonia***Purpose***

Respiratory stimulant

Ammonia***Uses***

- to prevent or treat fainting

Ammonia***Warnings*****For external use only****Do not use**

- if you have breathing problems such as asthma or emphysema

Stop use and ask a doctor if

- condition persists

Keep out of reach of children

- If swallowed get medical help or contact a Poison Control Center right away.

Ammonia***Directions***

- hold inhalant away from face and crush ampoule between thumb and forefinger at position indicated on sleeve.
- hold near nostrils for inhalation of volatile vapor

Ammonia***Other information***

- store at room temperature away from light

Ammonia***Inactive ingredient***

alcohol USP, FD&C red #40, lavender oil, lemon oil fcc, nutmeg oil, purified water

Ammonia***Questions or Comments?***

1-800-430-5490

Water Soluble 1st Aid Spray***Active ingredient***

Benzethonium chloride 0.2% w/w - Benzocaine 10% w/w

Water Soluble 1st Aid Spray***Purpose***

Topical antiseptic

Topical anesthetic

Water Soluble 1st Aid Spray***Uses***

for temporary relief of pain and itching and helps protect against infection in

- minor cuts and scrapes
- insect bites
- minor skin irritations

Water Soluble 1st Aid Spray***Warnings*****For external use only****Flammable**

- keep away from fire or flame
- contents under pressure
- do not puncture or incinerate container
- do not expose to temperature above 120 0 F

Do not use

- in the eyes or other mucous membranes
- in cases of serious burns
- in case of deep orpuncture wounds
- for a prolonged period of time
- on large portion of the body

Stop use and ask a doctor if

- condition worsens or symptoms persist for more than 7 days
- condition clears up and occurs again within a few days
- redness, swelling, or irritation occurs

Keep out of reach of children

- If swallowed, get medical help or contact a Poison Control Center right away

Water Soluble 1st Aid Spray***Directions***

- clean the affected area
- shake can well before using
- hold 4 - 6 inches from surface and spray area until wet
- may be covered with a sterile bandage. If bandaged, let dry first
- for adult institutional use only
- not intended for use on children

Water Soluble 1st Aid pray***Other information***

- avoid inhaling
- use only as directed
- intentional misuse by deliberately concentrating and inhaling the contents may be harmful or fatal

Water Soluble 1st Aid pray***Inactive ingredients***

dipropylene glycol, isobutane, N-butane, propane

Burn Relief Water Soluble***Active ingredients***

Benzethonium chloride 0.2% w/w

Benzocaine 10% w/w

Menthol 0.33% w/w

Burn Relief Water Soluble***Purpose***

Topical antiseptic

Topical anesthetic

Topical anesthetic

Burn Relief Water Soluble***Uses***

for the temporary relief of pain and itching and helps protect against infection in:

- minor cuts and scrapes
- burns
- sunburn
- insect bites
- minor skin irritations

Burn Relief Water Soluble

Warnings

For external use only

Flammable keep away from fire or flame

- contents under pressure
- do not puncture or incinerate container
- do not expose to temperatures above 120 0 F

Do not use

- in or near the eyes or other mucous membranes
- in case of serious burns
- in case of deep or puncture wounds
- for prolonged period of time
- on large portion of the body

Stop use and ask a doctor if

- condition worsens or symptoms persist for more than 7 days
- condition clears up and recurs within a few days
- redness, swelling, or irritation occurs

Keep out of the reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Burn Relief Water Soluble

Directions

- clean the affected area
- shake can well before using
- hold 4 - 6 inches from surface and spray area until wet
- may be covered with a sterile bandage, if bandaged let dry first
- for adult institutional use only
- not intended for use on children

Burn Relief Water Soluble

Other information

- avoid inhaling
- use only as directed
- intentional misuse by deliberately concentrating or inhaling the contents may be harmful or fatal

Burn Relief Water Soluble

Inactive ingredients

dipropylene glycol, isobutane, n-butane, propane

Triple

Active ingredients

Bacitracin zinc 400 units

Neomycin sulfate (5 mg equivalent to 3.5 mg Neomycin base)

Polymyxin B sulfate 5000 units

Triple

Purpose

First aid antibiotic

First aid antibiotic

First aid antibiotic

Triple

Uses

- first aid to help prevent infection in
- minor cuts
- scrapes
- burns

Triple

Warnings

For external use only

Allergy alert do not use if you are allergic to any of the ingredients

Do not use

- in the eyes
- over large areas of the body

Ask a doctor before use if you have

- a deep or puncture wounds
- animal bites
- serious burns

Stop use and ask a doctor if

- the condition persists or gets worse
- a rash or other allergic reaction develops
- you need to use longer than 1 week

Keep out of the reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Triple

Directions

- clean the affected area
- apply a small amount of the product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage

Triple

Other information

- store at 15⁰ to 25⁰ C (59⁰ to 77⁰ F)
- tamper evident sealed packets
- do not use if packet is torn or opened

Triple

Inactive ingredient

petrolatum

Alcohol

Active ingredient

Isopropyl alcohol 70%

Alcohol

Purpose

First aid antiseptic

Alcohol

Uses

- first aid to help prevent infection in minor cuts, scrapes, and burns

Alcohol

Warnings

For external use only

Flammable, keep away from fire or flame.

Do not use

- in the eyes
- over large areas of the body

Ask a doctor before use if you have

deep or puncture wounds

animal bites

serious burns

When using this product

- do not use longer than one week unless directed by a doctor

Stop use and consult a doctor

- if condition persists or gets worse

Keep out of reach of children

- If swallowed, get medical help or contact a Poison Control Center right away.

Alcohol

Directions

- clean the affected area
- apply wipe to affected area 1 to 3 times daily
- may be covered with a sterile bandage
- discard wipe after single use

Alcohol

Other information

store at room temperature 15⁰ to 25⁰ C (59⁰ to 77⁰ F)

Alcohol

Inactive ingredient

water

Miralac

Active ingredient (in each chewable tablet)

Miralac

Active ingredient (in each chewable tablet)

Calcium carbonate 420 mg

Miralac

Purpose

Antacid

Miralac

Uses

for the relief of

- acid indigestion
- heartburn
- sour stomach
- upset stomach associated with these symptoms

Warnings

Ask a doctor before use if you have

- kidney stones
- calcium-restricted diet

Ask a doctor before use if you are

- presently taking a prescription drug. Antacids may interfere with certain prescription drugs

When using this product

- do not take more than 12 tablets in a 24- hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a doctor.

Keep out of the reach of children.

In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

Miralac

Directions

- chew 1 to 2 tablets every 4 hours as symptoms occur, or as directed by a doctor.

Miralac

Other information

- each tablet contains: calcium 170 mg
- sucrose free
- lactose free
- store at room temperature
- TAMPER EVIDENT PACKETS- DO NOT USE IF OPEN OR TORN

Miralac

Inactive ingredients

magnesium stearate, mint flavor, silicon dioxide, sorbitol, starch

Miralac

Questions or comments?

1-800-430-5490

BZK

Active ingredient

Benzalkonium chloride 0.13% w/v

BZK

Purpose

First aid antiseptic

BZK

Uses

Antiseptic cleansing of face, hands, and body without soap and water

BZK

Warnings

For external use only

Do not use

- in the eyes or over large areas of the body
- on mucous membranes
- on irritated skin
- in case of deep puncture wounds, animal bites or serious burns, consult a doctor

- longer than 1 week unless directed by a doctor

Stop use and ask a doctor if

- if irritation, redness or other symptoms develop
- the condition persists or gets worse

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away

BZK

Directions

tear open packet and use as a washcloth

BZK

Other information

- store at room temperature 15⁰ to 30⁰ C (5⁰ - 86⁰ F)
- do not reuse towelette

BZK

Inactive ingredient

water

BzK

Questions

1-800-430-5490

Aypanal

Active ingredient

Acetaminophen 500 mg

Aypanal

Purpose

Pain reliever/fever reducer

Aypanal

Uses

- temporarily relieves minor aches and pains due to the common cold and headache
- temporarily reduces fever

Ask a doctor before use if you have

- liver disease

Aypanal

Warnings

Liver Warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount.
- with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If skin reaction occurs, stop use and seek medical help right away

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription).
- If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

If pregnant or breastfeeding

- ask a health professional before use.

Keep out of reach of children.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Aypanal

Directions

- **do not take more than directed (see overdose warning)**
- adults and children 12 years of age and over: Take 2 tablets with water every 6 hours while symptoms last.
- do not take any more than 8 tablets in 24 hours.
- children under 12: consult a doctor

Aypanal

Other information

- store at room temperature 15⁰ -30⁰ C (59⁰ -86⁰ F)
- TAMPER EVIDENT- DO NOT USE IF OPEN OR TORN

Aypanal

Inactive ingredients

microcrystalline cellulose, povidone, sodium starch glycolate, starch, stearic acid

Aypanal

Questions or Comments

1-800-430-5490

4240

68400LNODEA KIT CONTENTS

1 3/4 X 3 PLAS 100/BOX
1 1X3 PLASTIC 100/BOX
1 WOVEN 2" X 3" 25/BOX
1 FINGERTIP "T" WOVEN 40/BOX
1 1X3 WOVEN SING 50/BOX
1 SWIFT KNUCKLE 40/BX
1 AMMONIA INHALANTS 10 PER
1 ELASTIC TAPE 1" X 5YD
1 FORCEPS POINTED METAL
1 O/H TAPE ADHESIVE TRI-CUT
1 FIRST AID GUIDE ASHI
6 GAUZE CLEAN-WRAP BDGE N/S 2"
2 BLOODSTOPPER
1 NON ADHERENT PADS 2"X3" 50'S
2 GZE PADS STERILE 2"X 2" 25'S
1 GZE PADS STERILE 4"X 4" 25'S
1 CO-FLEX BANDAGE 2"X 5YDS TAN
1 COTTON TIPS 100 PER VIAL
1 ANTISEPTIC WIPES BZK CHL 20'S
1 FIRST AID SPRAY AEROSOL 3 OZ
1 ALCOHOL WIPES 50'S
1 ASPIRIN IND PK 5 GR 2/ENV 250
1 AYPANAL NON-ASP IND 2/ENV 100
1 MIRALAC TABS IND PK 2/ENV 250
1 BURN SPRAY 3 OZ
1 TRIPLE BIOTIC .5 GRAM PKT 20
1 COLD PACK 5"X9" BOXED
1 4OZ BFS EYEWASH TRILINGUAL BOTTLE
1 SCISSOR BDGE 4" RED PLS HDL
1 400 EMPTY KIT BLANK

1 POCKET INSERT RED #400 KIT 5R
1 TONGUE BLADES SR WRAPPED 6'S
1 LBL STOCK 6-3/8"X4"
1 LBL STOCK 4"X2-7/8"
1 LBL STOCK 3"x1-7/8"
2 PR LRG NITRILE GLVES ZIP BAG
2 TRI BNDG NON WOVEN 40"X40"X56"
1 RED BIO BAGS 2/BX

Eyewash

Principal Display Panel

Honeywell

TAMPER-EVIDENT CAP.
TAPA CON SELLO DE SEGURIDAD.
BOUCHON INDICATEUR D'INFRACTION.

eyesaline®

LAVAOJOS
EYESALINE

Solución
Isotónico Estéril

EYESALINE
EYEWASH

Sterile
Isotonic Solution

LAVAGE
OCULAIRE
EYESALINE

La Solution
Isotonique Stérile

16 fl. oz. (473 mL)

NPN: 80057528
64809 1 45033 117

Drug Facts (for USA only)

Active ingredient Sterile water 99%
Purpose Eyewash
Uses for flushing the eye to remove loose foreign material, air pollutants, or chlorinated water.
Warnings For external use only - Obtain immediate medical treatment for all open wounds in or near the eyes. To avoid contamination, do not touch tip of container to any surface. Do not reuse. Once opened, discard.
Do not use if solution changes color or becomes cloudy
if you have open wounds in or near the eyes, get medical help right away
Stop use and consult a doctor if:
• you experience eye pain • changes in vision
• continued redness or irritation of the eye
• condition worsens or persists
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.
Directions
• remove contacts before using • twist top to remove
• flush the affected area as needed
• control rate of flow by pressure on the bottle
• if necessary, continue flushing with emergency eyewash or shower
Inactive ingredients sodium chloride, sodium phosphate dibasic, sodium phosphate monobasic
Questions? Call 1-800-430-5490
Honeywell Safety Products USA, Inc. Smithfield, RI. 02917

LABEL #32-0045/0 Rev. J REORDER / NUEVO PEDIDO / REAPPROVISIONNEMENT #32-000454-0000

space for lot code and supplier part number

PEEL / PELAR / PELEL


Datos de medicamento (Para EE.UU. solamente)

Ingrediente Activo Agua estéril 99%
Propósito Lavaojos
Usos para el lavado de ojo para quitar las partículas sueltas y extrañas, los contaminantes aeros, o agua de cloruro
Advertencias Para el uso externo sólo - Obtenga tratamiento médico inmediato para todas las heridas abiertas en o cerca de los ojos. Para evitar la contaminación, no toque la punta del envase con ninguna superficie. No vuelva a usar. Vez abierto, descarte.
No se use si la solución se enturbia o cambia de color
si tiene heridas abiertas en o cerca del ojo, obtenga ayuda medica de inmediato
Deje de usar y consulte a un médico si:
• experimenta dolor de ojo • cambio de visión
• rojez continuo o irritación del ojo
• la condición empeora o persiste
Manténgase fuera del alcance de los niños. En caso de ingestión accidental, obtenga atención médica o llame a un Centro de control de envenenamiento inmediatamente.
Instrucciones
• quite los lentes de contacto antes de usar la solución
• tuerza la tapa para quitar
• enjuague el área afectada según se necesite
• controle el chorro haciendo presión el la botella
• si es necesario, sigue enjuagado con un lavaojos o ducha de emergencia
Ingredientes inactivos cloruro de sodio, fosfato de sodio dibásico, fosfato de sodio monobásico
¿Preguntas? Llame al 1-800-430-5490
Honeywell Safety Products USA, Inc. Smithfield, RI. 02917

Information

Usages Pour le rinçage des yeux afin d'enlever un corps étranger, des polluants atmosphériques ou de l'eau chlorée.
Advertissements Pour usage externe seulement - Obtenir immédiatement des soins médicaux pour toutes les plaies ouvertes dans ou près des yeux. Pour éviter toute contamination, ne pas toucher la pointe du récipient à n'importe quelle surface. Ne pas réutiliser. Une fois ouvert, jetez-le.
Ne pas utiliser si la solution a changé de couleur ou si elle est brouillée
si vous avez des plaies ouvertes aux yeux ou à proximité, consultez immédiatement un médecin
Cesser d'utiliser la solution et consulter un médecin si vous ressentez une douleur oculaire si votre vision change
rougeur ou irritation persistante des yeux
condition empire ou persiste
Garder hors de la portée des enfants. En cas d'ingestion, communiquer immédiatement avec un médecin ou avec un centre antipoison.
Mode d'emploi enlever les verres de contact avant l'utilisation dévisser le bouchon pour l'enlever rincer la zone touchée selon les besoins ajuster le débit d'écoulement de la solution en augmentant ou en réduisant la pression exercée sur le contenant si nécessaire, continuer de rincer avec une solution de rinçage oculaire d'urgence ou une douche
Ingédients eau stérile, chlorure de sodium, phosphate dibasique de sodium, phosphate monobasique de sodium
Des questions? Faites le 1-800-430-5490
Honeywell Safety Products USA, Inc. Smithfield, RI. 02917

Aspirin
Principal Display Panel



7722-01 Rev C

Drug Facts (continued)
Active ingredient (in each tablet): Aspirin 325mg (NSAID)
Purpose: Pain reliever / fever reducer
Nonsteroidal anti-inflammatory drug
Uses: temporarily reduces fever and relieves minor aches and pains associated with:
 - cold - headache - toothache - menstrual aches - backache - minor pain of arthritis
Warnings:
 - aspirin allergy: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, changes in behavior with nausea and vomiting occur. Consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness.
Warnings (continued):
 - aspirin allergy: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, changes in behavior with nausea and vomiting occur. Consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness.



6 96351 53168 2
 Distributed by
 Honeywell Safety
 Products, USA, Inc.
 Smithfield, RI 02917

PACKAGE NOT CHILD-RESISTANT
 Total quantity 50 Tablets
 25 Packets (2 Tablets per Packet)
Pain Reliever / Fever Reducer
ASPIRIN 325 MG
(NSAID)
Honeywell

ASPIRIN 325 MG
 25 Packets (2 Tablets per Packet)
 Total quantity 50 Tablets
 35222AA

Drug Facts (continued)
Allergy alert: Aspirin may cause a severe allergic reaction which may include:
 • hives • facial swelling • asthma (wheezing) • shock
Stomach bleeding warning: This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if you:
 • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed
Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.
Ask a doctor before use if: - stomach bleeding warning applies to you - you have a history of stomach problems such as heartburn - you have high blood pressure, heart disease, liver cirrhosis, or kidney disease - you are taking a diuretic - you have asthma
Ask a doctor or pharmacist before use if you are: taking a prescription drug for diabetes, gout or arthritis
Stop use and ask a doctor if: - you experience any of the following signs of stomach bleeding:
 • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better
 • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days
 • redness or swelling is present in the painful area • ringing in the ears or loss of hearing occurs
 • any new symptoms appear
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center immediately.



Drug Facts (continued)
Directions - drink a full glass of water with each dose
 • adults and children 12 years of age and older: take 1 - 2 tablets every 4 hours while symptoms last, not more than 12 tablets in 24 hours
 • children under 12 years of age: consult a doctor
Other information - Store at room temperature 15° - 30° C (59° - 86° F)
 • **TRIPLEX EVIDENCE PACKETS - DO NOT USE IF OPEN OR TORN**
Inactive ingredients - corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, polyethylene glycol, povidone, propylene glycol, silicon dioxide, stearic acid, titanium dioxide
 *may contain these ingredients
Questions or comments? 1-800-430-5490

ASPIRIN 325 MG
 25 Packets (2 Tablets per Packet)
 Total quantity 50 Tablets

7722-01 Rev C.pdf 1 9/5/2018 2:18:46 PM

Ammonia
Principal Display Panel

796006 Rev. E Unit Carton Printing Plate for "A" size carton.



796006 Rev. E (page 3 of 3)

Principal Display Panel

SHAKE WELL BEFORE USING

Honeywell

FIRST AID ANTISEPTIC SPRAY

Water soluble
Benzethonium chloride
 Topical antiseptic
Benzocaine
 Topical anesthetic

Helps prevent infection and relieves pain.

CAUTION: FLAMMABLE
 Contents under pressure
 Read warning on back panel.

NET WT. 3 OZ (85gm.)

47.0303

Cat. No. 151019

DRUG FACTS	
Active ingredients	Purpose
Benzethonium chloride 0.2% w/w.....	Topical antiseptic
Benzocaine 10% w/w.....	Topical anesthetic
Uses	
• for the temporary relief of pain and itching and helps to protect against infection in • minor cuts and scrapes • insect bites • minor skin irritations	
Warnings	
For external use only	
• keep away from fire or flame	• contents under pressure
• do not puncture or incinerate container	• do not expose to temperatures above 120°F
Do not use • in or near eyes or other mucus membranes • in case of serious burns • in case of deep or puncture wounds • for a prolonged period of time • on large portion of the body	
Stop use and ask a doctor if:	
• conditions worsens or symptoms persist for more than 7 days • condition clears up and recurs within a few days • redness, swelling or irritation occurs	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions • clean the affected area • shake can well before using • hold 4-6 inches from surface and spray area until wet • may be covered with a sterile bandage. If bandaged, let dry first • for adult institutional use only • not intended for use on children	
Other information • avoid inhaling • use only as directed • intentional misuse by deliberately concentrating and inhaling the contents may be harmful or fatal	
Inactive ingredients dipropylene glycol, isobutane, n-butane, propane	
Questions or comments? 1-800-430-5490	



Distributed by
 Honeywell Safety
 Products USA, Inc.
 Smithfield, RI 02917

Honeywell

Burn Relief Water Soluble Principal Display Panel

SHAKE WELL BEFORE USING

Honeywell

BURN SPRAY

Water soluble
Benzethonium chloride
 Topical antiseptic
Benzocaine
 Topical anesthetic
Menthol
 Topical anesthetic

Provides antiseptic treatment and helps relieve the pain of minor burns and sunburn.

CAUTION: FLAMMABLE
 Contents under pressure
 Read warning on back panel.

NET WT. 3 OZ (85gm.)

47.0306

Cat. No. 201005


DRUG FACTS	
Active ingredients	Purpose
Benzethonium chloride 0.2% w/w.....	Topical antiseptic
Benzocaine 10% w/w.....	Topical anesthetic
Menthol .33%.....	Topical anesthetic
Uses • for the temporary relief of pain and itching and helps to protect against infection in • minor cuts and scrapes • burns • sunburn • insect bites • minor skin irritations	
Warnings	
For external use only	
• keep away from fire or flame	• contents under pressure
• do not puncture or incinerate container	• do not expose to temperatures above 120°F
Do not use • in or near eyes or other mucus membranes • in case of serious burns • in case of deep or puncture wounds • for a prolonged period of time • on large portion of the body	
Stop use and ask a doctor if:	
• conditions worsens or symptoms persist for more than 7 days • condition clears up and recurs within a few days • redness, swelling or irritation occurs	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions • clean the affected area • shake can well before using • hold 4-6 inches from surface and spray area until wet • may be covered with a sterile bandage. If bandaged, let dry first • for adult institutional use only • not intended for use on children	
Other information • avoid inhaling • use only as directed • intentional misuse by deliberately concentrating and inhaling the contents may be harmful or fatal	
Inactive ingredients dipropylene glycol, isobutane, n-butane, propane	
Questions or comments? 1-800-430-5490	




Distributed by
 Honeywell Safety
 Products USA, Inc.
 Smithfield, RI 02917

Honeywell


Triple Principal Display Panel



Alcohol Principal Display Panel



Miralac
Principal Display Panel



BZK
Principal Display Panel

47001083
Rev B

Antiseptic Towelettes

Honeywell

02-16-35MD

Antiseptic Towelettes

Benzalkonium chloride
First aid antiseptic

Six-Saturated Towelettes

Distributed by
Honeywell Safety
Products USA, Inc.
Smithfield, RI 02917

47001083
Rev B

Antiseptic Towelettes

Honeywell

Drug Facts

Active Ingredient	Purpose
Benzalkonium chloride 0.133% w/v	First aid antiseptic

Uses • antiseptic cleaning of face, hands and body without soap and water. • air dries in seconds

Warnings

For external use only

When using this product • do not use in the eyes or apply over large areas of the body

Ask a doctor before use • in case of deep or puncture wounds, animal bites, or serious burns

Stop use and consult a doctor if

• irritation, redness or other symptoms develop • condition persists or gets worse

Do not use • longer than 1 week unless directed by doctor

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions • tear open packet, unfold and use as washcloth

Other information

• store at room temperature 15° -30° C (59° -86° F) • do not reuse towelette

Inactive ingredient water

Questions or comments 1-800-430-5490

Aypanal
Principal Display Panel

4240 Kit Label
68400LNODEA

AYPANAL
25 Packets (2 Tablets Per Packet)
Quantity 50 Tablets

Drug Facts (continued)

Other information • store at room temperature 15-30°C (59-86°F)
• TABLETS: EXPIRES ON 12/31/15 • DO NOT USE IF CAPS OR TORN

Inactive ingredients corn starch, microcrystalline cellulose, polydioxane, sodium starch glycolate, stearic acid

Questions or comments? 1-800-430-5490

Honeywell

AYPANAL
25 Packets (2 Tablets Per Packet)
Quantity 50 Tablets


AYPANAL
Acetaminophen 325 mg
Pain Reliever-Fever Reducer

**25 Packets (2 Tablets Per Packet)
Quantity 50 Tablets**

PACKAGE NOT CHILD-RESISTANT

UNIT NO. 35225AP

Distributed by
Honeywell Safety
Products USA, Inc.
Smithfield, RI 02917



6 69635 55170 5

Drug Facts

Active ingredient (in each tablet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer

Uses • temporarily relieves minor aches and pains due to the common cold and headache
• temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if:
• adult takes more than 4,000 mg acetaminophen in 24 hours, which is the maximum daily amount
• child takes more than 5 doses in 24 hours, which is the maximum daily amount
• taken with other drugs containing acetaminophen
• adult has 3 or more alcoholic drinks every day while using this product

Drug Facts (continued)

Ask a doctor before use if you have liver disease

Ask a doctor before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if • pain gets worse or lasts more than 10 days in adults
• pain gets worse or lasts more than 5 days in children under 12 years

Keep out of reach of children

Overdose warning: In case of accidental overdose, get medical help or contact a poison control center right away. For more information, call a poison control center or call 1-800-430-5490.

Directions • do not take more than directed (see overdose warning)

Children 6 to under 12 years and older
• take 2 tablets every 4 to 6 hours while symptoms last
• do not take more than 12 tablets in 24 hours

Children under 6 years
• take 1 tablet every 4 to 6 hours while symptoms last
• do not take more than 5 tablets in 24 hours
• consult a doctor

Other information • store at room temperature 15-30°C (59-86°F)
• TABLETS: EXPIRES ON 12/31/15 • DO NOT USE IF CAPS OR TORN

Inactive ingredients corn starch, microcrystalline cellulose, polydioxane, sodium starch glycolate, stearic acid

Questions or comments? 1-800-430-5490

77225-01 Rev.B



Distributed by Honeywell Safety Products USA, Inc Smithfield, RI 02917

4240 FIRST AID KIT

4240 first aid kit kit

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:0498-4240

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-4240-01	1 in 1 KIT; Type 0: Not a Combination Product	10/18/2018	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	125 PACKET	250
Part 2	1 BOTTLE	118 mL
Part 3	125 PACKET	250
Part 4	10 AMPULE	3 mL
Part 5	1 CAN	85 g
Part 6	1 CAN	85 g
Part 7	20 PACKET	10 g
Part 8	50 POUCH	20 mL
Part 9	20 PACKET	28 mL
Part 10	50 PACKET	100

Part 1 of 10

MIRALAC

calcium carbonate tablet

Product Information

Item Code (Source)	NDC:0498-0303
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CALCIUM CARBONATE (UNII: H0G9379FGK) (CALCIUM CATION - UNII:2M83C4R6ZB, CARBONATE ION - UNII:7UJQ50PE7D)	CALCIUM CARBONATE	420 mg

Inactive Ingredients

Ingredient Name	Strength
MAGNESIUM STEARATE (UNII: 70097M6I30)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SORBITOL (UNII: 506T60A25R)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	white	Score	2 pieces
-------	-------	-------	----------

Shape	ROUND	Size	11mm
Flavor	MINT	Imprint Code	FR8
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part331	02/22/2012	

Part 2 of 10

EYESALINE EMERGENCY EYEWASH

purified water liquid

Product Information

Item Code (Source)	NDC:0498-0100
Route of Administration	OPHTHALMIC

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
WATER (UNII: 059QF0KO0R) (WATER - UNII:059QF0KO0R)	WATER	98.6 mL in 100 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM PHOSPHATE, MONOBASIC, MONOHYDRATE (UNII: 593YOG76RN)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM PHOSPHATE, DIBASIC (UNII: GR686LBA74)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0100-02	118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

Part 3 of 10

ASPIRIN

aspirin tablet

Product Information

Item Code (Source)	NDC:0498-0114
--------------------	---------------

Route of Administration	ORAL
-------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE 2208 (100 MPA.S) (UNII: B1QE5P712K)	
MINERAL OIL (UNII: T5L8T28FGP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	FR21
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0114-01	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	09/18/2018	

Part 4 of 10

AMMONIA INHALENT

ammonia inhalent inhalant

Product Information

Item Code (Source)	NDC:0498-3334
Route of Administration	RESPIRATORY (INHALATION)

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
AMMONIA (UNII: 5138Q19F1X) (AMMONIA - UNII:5138Q19F1X)	AMMONIA	0.045 g in 0.3 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-3334-00	0.3 mL in 1 AMPULE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	

Part 5 of 10

FIRST AID ANTISEPTIC WATER SOLUBLE

benzethonium chloride, benzocaine spray

Product Information

Item Code (Source)	NDC:0498-0031
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	0.2 g in 100 g
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	10 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
ISOBUTANE (UNII: BXR49TP611)	
BUTANE (UNII: 6LV4FOR43R)	
PROPANE (UNII: T75W9911L6)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0031-40	85 g in 1 CAN; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333E	09/19/2018	

Part 6 of 10**BURN WATER SOLUBLE**

benzocaine, benzethonium chloride, menthol spray

Product Information

Item Code (Source)	NDC:0498-0021
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	0.2 g in 100 g
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	10 g in 100 g
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	0.33 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
ISOBUTANE (UNII: BXR49 TP6 11)	
BUTANE (UNII: 6LV4FOR43R)	
PROPANE (UNII: T75W99 11L6)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0021-40	85 g in 1 CAN; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part348	11/12/2018	

Part 7 of 10

TRIPLE ANTIBIOTIC

bacitracin zinc, polymyxin b sulfate, neomycin sulfate ointment

Product Information

Item Code (Source)	NDC:0498-0750
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96 K)	POLYMYXIN B	5000 [iU] in 1 g
BACITRACIN ZINC (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN	400 [iU] in 1 g
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN	3.5 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
PETROLATUM (UNII: 4T6H12BN9U)	

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0750-36	0.5 g in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333B	09/19/2018	

Part 8 of 10**ALCOHOL WIPE**

isopropyl alcohol swab

Product Information

Item Code (Source)	NDC:0498-0143
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	0.7 mL in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0143-04	0.4 mL in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	09/18/2018	

Part 9 of 10**ANTISEPTIC TOWELETTE**

benzalkonium chloride liquid

Product Information

Item Code (Source) NDC:0498-0501

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	1.3 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0501-00	1.4 mL in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333E	12/22/2017	

Part 10 of 10

AYPANAL NON-ASPIRIN

acetaminophen tablet

Product Information

Item Code (Source) NDC:0498-2001

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	circle;U
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	04/10/2012	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		10/18/2018	

Labeler - Honeywell Safety Products USA, INC (079287321)

Establishment

Name	Address	ID/FEI	Business Operations
James Alexander		040756421	manufacture(0498-3334)

Establishment

Name	Address	ID/FEI	Business Operations
Honeywell Safety Products USA, INC		079287321	pack(0498-4240)

Establishment

Name	Address	ID/FEI	Business Operations
Ultra Seal Corporation		085752004	manufacture(0498-0114, 0498-2001, 0498-0303)

Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

Dixon Investments		115315822	manufacture(0498-0031, 0498-0021)
-------------------	--	-----------	-----------------------------------

Establishment

Name	Address	ID/FEI	Business Operations
Water-Jel Technologies		155522589	manufacture(0498-0750)

Establishment

Name	Address	ID/FEI	Business Operations
Honeywell Safety Products USA, Inc.		167518617	manufacture(0498-0100)

Establishment

Name	Address	ID/FEI	Business Operations
Changzhou Maokang Medical		421317073	manufacture(0498-0143, 0498-0501)

Revised: 4/2019

Honeywell Safety Products USA, INC