

**MUCUS RELIEF COLD FLU AND SORE THROAT MAXIMUM STRENGTH-
acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid
Family Dollar (FAMILY WELLNESS)**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients (in each 20 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Phenylephrine HCl 10 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - sinus congestion and pressure
 - minor aches and pains
 - nasal congestion
 - cough due to minor throat and bronchial irritation
 - sore throat
 - headache
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen

- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product,

do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL every 4 hours while symptoms last
- children under 12 years of age: do not use

Other information

- **each 20 mL contains:** sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, disodium EDTA,, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

Principal Display Panel

COMPARE TO THE ACTIVE INGREDIENTS IN MUCINEX® FAST-MAX® MAXIMUM STRENGTH COLD, FLU, & SORE THROAT*

Maximum Strength

Mucus Relief

Cold, Flu & Sore Throat

Multi-Symptom Relief

Pain Reliever/Fever Reducer - **Acetaminophen**

Cough Suppressant - Dextromethorphan HBr

Expectorant - Guaifenesin

Nasal Decongestant - Phenylephrine HCl

For Relief of:

Nasal Congestion

Sore Throat, Mucus

Headache, Body ache

Cough, Fever

FL OZ (mL)

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Mucinex® Fast-Max® Maximum Strength Cold, Flu, & Sore Throat.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY:

FAMILY DOLLAR SERVICES, INC.

10401 MONROE ROAD

MATTHEWS, NC 28105 USA

FAMILYDOLLAR.COM

PRODUCT OF USA

Package Label

NDC 55319-336-06

COMPARE TO THE ACTIVE
INGREDIENTS IN MUCINEX®
FAST-MAX® MAXIMUM STRENGTH
COLD, FLU & SORE THROAT*

FAMILY
wellness™



Maximum Strength Mucus Relief Cold, Flu & Sore Throat

Multi-Symptom Relief

Pain Reliever/Fever Reducer - **Acetaminophen**

Cough Suppressant - Dextromethorphan HBr

Expectorant - Guaifenesin

Nasal Decongestant - Phenylephrine HCl

For Relief of:
Nasal Congestion
Sore Throat, Mucus
Headache, Body Ache
Cough, Fever

6 FL OZ
(177 mL)

PLD-B282B
LB002607

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY
SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Mucinex® Fast-Max® Maximum Strength Cold, Flu & Sore Throat.

SKU 0903758



DISTRIBUTED BY:
FAMILY DOLLAR SERVICES, INC.,
10401 MONROE ROAD
MATTHEWS, NC 28105 USA
VISIT FAMILYDOLLAR.COM
PRODUCT OF USA

Not 100% satisfied? Return package
and unused product within 30 days
to any Family Dollar store for a
refund (with receipt) or exchange.

PARENTS:
Learn about teen medicine at
www.StopMedicineAbuse.org

PLD-C282C LB002606

Drug Facts

Active ingredients (in each 20 mL)

Acetaminophen 650 mg.....Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....Cough suppressant
Guaifenesin 400 mg.....Expectorant
Phenylephrine HCl 10 mg.....Nasal decongestant

Purposes

PEEL CORNER FOR MORE DRUG FACTS

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

Uses

- temporarily relieves these common cold and flu symptoms ■ sinus congestion and pressure
- minor aches and pains ■ nasal congestion
- cough due to minor throat and bronchial irritation ■ sore throat ■ headache
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash
- If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease ■ heart disease ■ diabetes
- thyroid disease ■ high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

When using this product, do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions ■ do not take more than directed (see Overdose warning)

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product ■ mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL every 4 hours while symptoms last
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments? Call
1-877-753-3935 Monday-Friday 9AM-5PM EST

FAMILY Wellness Mucus Relief Cold, Flu, & Sore Throat Maximum Strength

MUCUS RELIEF COLD FLU AND SORE THROAT MAXIMUM STRENGTH

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-336
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-336-06	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/31/2014	04/26/2024

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	12/31/2014	04/26/2024

Labeler - Family Dollar (FAMILY WELLNESS) (024472631)

