

IBUPROFEN - ibuprofen tablet
MARKSANS PHARMA LIMITED

Ibuprofen Tablets 200mg (Orange-Round and Capsule shaped)

Active ingredient(s)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Use(s)

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - toothache
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease, asthma, or had a stroke.
- you are taking a diuretic

Ask a doctor or pharmacist before use if

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood

- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in painful area
- any new symptoms appear

If pregnant or breast-feeding

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause a problems in the unborn child or complications during delivery.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**

adults and children 12 years and older	<ul style="list-style-type: none"> ▪ take 1 tablet every 4 to 6 hours while symptoms persist ▪ if pain or fever does not respond to 1 tablet, 2 tablets may be used ▪ do not exceed 6 tablets in 24 hours, unless directed by a doctor.
children under 12 years	<ul style="list-style-type: none"> ▪ ask a doctor

Other information

- store at 20-25°C (68-77°F)
- **do not use if carton is opened. Do not use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing**
- see end panel for lot number and expiration date

Inactive ingredient(s)

colloidal silicon dioxide, croscarmellose sodium, FD & C yellow # 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.

Questions or comments?

Call 1-877-290-4008

Manufactured for:

Time-Cap Labs, Inc.

7 Michael Avenue,
Farmingdale,
NY 11735, USA

Manufactured by:

Marksans Pharma Ltd.

Plot No. L-82, L-83,
Verna Indl. Estate,
Verna, Goa-403 722, India.

Principal Display Panel

NDC 25000-136-03, Ibuprofen Tablets, USP 200mg

30 counts Bottle Label

NDC 25000-136-03

Ibuprofen Tablets USP, 200 mg

Pain Reliever / Fever Reducer (NSAID)

30 Coated Tablets



Marksans

Important: Read all product information before using. Keep the box for important information.

Active ingredient (in each tablet) Purpose

Ibuprofen 200 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
■ headache ■ muscular aches ■ minor pain of arthritis →

LOT :

EXP :

R412/21

■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever. **Warnings: Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. →

Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This →

product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you : ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug →

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than →

directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ →

if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel →

faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part →

■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. →

a health professional before use. **Keep out of reach of children. Directions :** ■ do not take more than directed ■ the smallest effective dose should be used. adults and children 12 years and →

or side of body ■ slurred speech → If pregnant or breast-feeding, ask → older : ■ take 1 tablet every 4 to 6 →

hours while symptoms persist ■
if pain or fever does not respond to
1 tablet, 2 tablets may be used ■
do not exceed 6 tablets in 24
hours, unless directed by a doctor.
children under 12 years: ■ ask a
doctor. **Other information** ■ store
→

between 20-25°C (68-77°F)

***Do not use if seal under bottle
cap imprinted with "SEALED
for YOUR PROTECTION" is
broken or missing.***

Mfg. Lic. No. : GO/DRUGS/515 →

Manufactured for:
Time-Cap Labs, Inc.
7 Michael Avenue, Farmingdale,
NY 11735, USA

Manufactured by:
Marksans Pharma Ltd.
Plot No. L-82, L-83, Verna Indl. Estate,
Verna, GOA - 403722, India.

NDC 25000-136-03, Ibuprofen Tablets, USP 200mg

30 counts Carton Label



NDC 25000-135-03, Ibuprofen Tablets, USP 200mg (Caplets)

30 counts Bottle Label

NDC 25000-135-03	
Ibuprofen Tablets USP, 200 mg	
Important: Read all product information before using. Keep the box for important information.	
Active ingredient (in each caplet)	Purpose

Pain Reliever / Fever Reducer
(NSAID)

30 Coated Caplets*
* Capsule-Shaped Tablets



Marksans

Ibuprofen 200 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
■ headache ■ muscular aches ■ minor pain of arthritis →

LOT :

EXP :

R412/21

■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever. **Warnings: Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. →

Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This →

product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you : ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug →

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than →

directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ →

if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel →

faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech →

■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding,** ask →

a health professional before use. **Keep out of reach of children. Directions :** ■ do not take more than directed ■ the smallest effective dose should be used. adults and children 12 years and older : ■ take 1 caplet every 4 to 6 →

hours while symptoms persist ■
if pain or fever does not respond to
1 caplet, 2 caplets may be used ■
do not exceed 6 caplets in 24
hours, unless directed by a doctor.
children under 12 years: ■ ask a
doctor. **Other information** ■ store

between 20-25°C (68-77°F)

***Do not use if seal under bottle
cap imprinted with "SEALED
for YOUR PROTECTION" is
broken or missing.***

Mfg. Lic. No. : GO/DRUGS/515

Manufactured for:
Time-Cap Labs, Inc.
7 Michael Avenue, Farmingdale,
NY 11735, USA

Manufactured by:
Marksans Pharma Ltd.
Plot No. L-82, L-83, Verna Indl. Estate,
Verna, GOA - 403722, India.

NDC 25000-135-03, Ibuprofen Tablets, USP 200mg (Caplets)

30 counts Carton Label



NDC 25000-136-69, Ibuprofen Tablets, USP 200mg

300 counts Bottle Label

NDC 25000-136-69

Ibuprofen Tablets USP, 200 mg

Pain Reliever / Fever Reducer (NSAID)

**THIS PACKAGE FOR HOUSEHOLDS
WITHOUT YOUNG CHILDREN**

300 Coated Tablets



Marksans

Drug Facts

Active ingredient (in each tablet) **Purpose**
Ibuprofen 200 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - toothache
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include :

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

LOT :

EXP :

R112/21



Drug Facts (Continued)

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you :

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAID's except aspirin, increases the risk of heart attack, heart failure, and stroke.

Drug Facts (Continued)

These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke.
- you are taking a diuretic

Drug Facts (Continued)

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problem or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part

Drug Facts (Continued)

- or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Drug Facts (Continued)

Directions

- do not take more than directed
- the smallest effective dose should be used

adults and children
12 years and older

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years

- ask a doctor

Other information

- store between 20-25°C (68-77°F).
- do not use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing

Drug Facts (Continued)

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, FD & C yellow # 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.

Questions or comments?

Call 1-877-290-4008

Manufactured for: **Time-Cap Labs, Inc.**
7 Michael Avenue, Farmingdale, NY 11735, USA

Manufactured by: **Marksans Pharma Ltd.**
Plot No. L-82, L-83, Verna Indl. Estate,
Verna, GOA - 403722, India.

Mfg. Lic. No. : GO/Drugs/515

NDC 25000-136-14, Ibuprofen Tablets, USP 200mg

1000 counts Bottle Label

NDC 25000-136-14

Ibuprofen Tablets USP, 200 mg

Pain Reliever / Fever Reducer
(NSAID)

**THIS PACKAGE FOR HOUSEHOLDS
WITHOUT YOUNG CHILDREN**

1000 Coated Tablets



Marksans

Drug Facts

Active ingredient (in each tablet)	Purpose
Ibuprofen 200 mg (NSAID)*	Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug	

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - toothache
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include :

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you :

- are age 60 or older

LOT :

EXP :

R412/21



Drug Facts (Continued)

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke.
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are →

Drug Facts (Continued)

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. →

Drug Facts (Continued)

Drug Facts (Continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than directed
- the smallest effective dose should be used

adults and children
12 years and older

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor.

children under 12 years

- ask a doctor

Other information

- store between 20-25°C (68-77°F).
- do not use if carton is opened. Do not use if seal under bottle cap imprinted with “SEALED for YOUR PROTECTION” is broken or missing

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, FD&C yellow# 6 aluminum lake, magnesium stearate, microcrystalline cellulose,

polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.

Questions or comments?

Call 1-877-290-4008

Manufactured for:
Time-Cap Labs, Inc.
7 Michael Avenue,
Farmingdale,
NY 11735, USA

Manufactured by:
Marksans Pharma Ltd.
Plot No. L-82, L-83,
Verna Indl. Estate,
Verna, GOA - 403722, India.

Mfg. Lic. No. : GO/DRUGS/515

NDC 25000-136-71, Ibuprofen Tablets, USP 200mg

1000 counts (PET) Bottle Label

NDC 25000-136-71

Ibuprofen Tablets USP, 200 mg

Pain Reliever / Fever Reducer (NSAID)

**THIS PACKAGE FOR HOUSEHOLDS
WITHOUT YOUNG CHILDREN**

1000 Coated Tablets



Marksans

Drug Facts

Active ingredient (in each tablet) *Purpose*
Ibuprofen 200 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - toothache
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen,

Drug Facts (Continued)

or others)

- have 3 or more alcoholic drinks every day while using this product

Take more or for a longer time than directed
Heart attack and stroke warning: NSAID's except aspirin, increases the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke.
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

LOT :

EXP :

R11221



Drug Facts (Continued)

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problem or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Drug Facts (Continued)

Directions

- do not take more than directed
- the smallest effective dose should be used

adults and children
12 years and older

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years

- ask a doctor

Other information

- store between 20-25°C (68-77°F).
- do not use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, FD&C yellow# 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.

Questions or comments?

Call 1-877-290-4008

Drug Facts (Continued)

Manufactured for:
Time-Cap Labs, Inc.
7 Michael Avenue,
Farmingdale,
NY 11735, USA

Manufactured by:
Marksans Pharma Ltd.
Plot No. L-82, L-83,
Verna Indl. Estate,
Verna, GOA - 403722, India.

Mfg. Lic. No. : GO/Drugs/515

NDC 25000-135-14, Ibuprofen Tablets, USP 200mg (Caplets)

1000 counts Bottle Label

NDC 25000-135-14

Ibuprofen Tablets USP, 200 mg

Pain Reliever / Fever Reducer
(NSAID)

Drug Facts

Active ingredient (in each caplet) *Purpose*
Ibuprofen 200 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - toothache
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic

**THIS PACKAGE FOR HOUSEHOLDS
WITHOUT YOUNG CHILDREN**

1000 Coated Caplets*
*Capsule-Shaped Tablets

M Marksans

Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include :

- hives ■ facial swelling ■ asthma (wheezing)
- shock ■ skin reddening ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you :

- are age 60 or older

LOT :

EXP :

R412/21



Drug Facts (Continued)

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke.
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are →

Drug Facts (Continued)

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint ■ vomit blood ■ have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. →

Drug Facts (Continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than directed
- the smallest effective dose should be used

adults and children
12 years and older

- take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor.

children under 12 years

- ask a doctor

Other information

- store between 20-25°C (68-77°F).
- do not use if carton is opened. Do not use if seal under bottle cap

Drug Facts (Continued)

polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.

Questions or comments?

Call 1-877-290-4008

Manufactured for:
Time-Cap Labs, Inc.
7 Michael Avenue,
Farmingdale,
NY 11735, USA

Manufactured by:
Marksans Pharma Ltd.
Plot No. L-82, L-83,
Verna Indl. Estate,
Verna, GOA - 403722, India.

imprinted with "SEALED for YOUR PROTECTION" is broken or missing

Mfg. Lic. No. : GO/DRUGS/515

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, FD&C yellow# 6
aluminum lake, magnesium stearate, microcrystalline cellulose, →

NDC 25000-135-70, Ibuprofen Tablets, USP 200mg (Caplets)

500 counts Bottle Label

NDC 25000-135-70

Ibuprofen Tablets USP, 200 mg

Pain Reliever / Fever Reducer (NSAID)

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

500 Coated Caplets*

*Capsule-Shaped Tablets

Marksans



Drug Facts

Active ingredient (in each caplet)	Purpose
Ibuprofen 200 mg (NSAID)* Pain reliever/fever reducer	
*nonsteroidal anti-inflammatory drug	

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - toothache
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include :

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

LOT :

EXP :

R212/21



Drug Facts (Continued)

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you :

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAID's except aspirin, increases the risk of heart attack, heart failure, and stroke.

Drug Facts (Continued)

These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke.
- you are taking a diuretic

Drug Facts (Continued)

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problem or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part

Drug Facts (Continued)

- or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Drug Facts (Continued)

Directions

- do not take more than directed
- the smallest effective dose should be used

adults and children 12 years and older

- take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor.

Children under 12 years

- ask a doctor

Other information

- store between 20-25°C (68-77°F).
- do not use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing

Drug Facts (Continued)

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, FD & C yellow # 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.

Questions or comments?

Call 1-877-290-4008

Manufactured for: **Time-Cap Labs, Inc.**
7 Michael Avenue, Farmingdale, NY 11735, USA

Manufactured by: **Marksans Pharma Ltd.**
Plot No. L-82, L-83, Verna Indl. Estate,
Verna, GOA - 403722, India.

Mfg. Lic. No. : GO/Drugs/515

NDC 25000-136-02, Ibuprofen Tablets, USP 200mg

24 counts Bottle Label

NDC 25000-136-02

Ibuprofen Tablets USP, 200 mg

Pain Reliever / Fever Reducer
(NSAID)

24 Coated Tablets



Marksans

**Important: Read all product information before using.
Keep the box for important information.**

Active ingredient (in each tablet) Purpose

Ibuprofen 200 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to:
■ headache ■ muscular aches ■ minor pain of arthritis →

LOT :

EXP :

R412/21

■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever. **Warnings: Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. →

Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This →

product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you : ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug →

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than →

directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■ →

if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel →

faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part →

■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear →

a health professional before use. **Keep out of reach of children. Directions :** ■ do not take more than directed ■ the smallest effective dose should be used. adults and children 12 years and →

breathing ■ weakness in one part
or side of body ■ slurred speech →

new symptoms appear.

If pregnant or breast-feeding, ask →

adults and children 12 years and
older : ■ take 1 tablet every 4 to 6 →

hours while symptoms persist ■
if pain or fever does not respond to
1 tablet, 2 tablets may be used ■
do not exceed 6 tablets in 24
hours, unless directed by a doctor.
children under 12 years: ■ ask a
doctor. **Other information** ■ store →

between 20-25°C (68-77°F)

***Do not use if seal under bottle
cap imprinted with "SEALED
for YOUR PROTECTION" is
broken or missing.***

Mfg. Lic. No. : GO/DRUGS/515 →

Manufactured for:

Time-Cap Labs, Inc.

7 Michael Avenue, Farmingdale,
NY 11735, USA


Manufactured by:

Marksans Pharma Ltd.

Plot No. L-82, L-83, Verna Indl. Estate,
Verna, GOA - 403722, India.


NDC 25000-136-02, Ibuprofen Tablets, USP 200mg

24 counts Carton Label



NDC 25000-135-02, Ibuprofen Tablets, USP 200mg (Caplets)

24 counts Bottle Label



■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever. **Warnings: Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This

product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you : ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than

directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** ■

if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. **Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel

faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech

■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. **If pregnant or breast-feeding,** ask

a health professional before use. **Keep out of reach of children. Directions :** ■ do not take more than directed ■ the smallest effective dose should be used. adults and children 12 years and older : ■ take 1 caplet every 4 to 6

hours while symptoms persist ■ if pain or fever does not respond to

between 20-25°C (68-77°F)

Manufactured for:
Time-Cap Labs, Inc.
7 Michael Avenue, Farmingdale

1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. children under 12 years: ■ ask a doctor. **Other information** ■ store

Do not use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

7 Michael Avenue, Farmingdale, NY 11735, USA

Manufactured by:
Marksans Pharma Ltd.
Plot No. L-82, L-83, Verna Indl. Estate, Verna, GOA - 403722, India.

Mfg. Lic. No. : GO/DRUGS/515

NDC 25000-135-02, Ibuprofen Tablets, USP 200mg (Caplets)

24 counts Carton Label



IBUPROFEN
ibuprofen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:25000-135
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	ORANGE	Score	no score
Shape	CAPSULE (Capsule Shaped)	Size	10mm
Flavor		Imprint Code	120
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25000-135-03	30 in 1 BOTTLE; Type 0: Not a Combination Product	11/06/2015	
2	NDC:25000-135-14	1000 in 1 BOTTLE; Type 0: Not a Combination Product	11/06/2015	
3	NDC:25000-135-30	6 in 1 BOX	11/06/2015	
3		6500 in 1 BAG; Type 0: Not a Combination Product		
4	NDC:25000-135-98	1 in 1 BOX	11/06/2015	
4		34065 in 1 BAG; Type 0: Not a Combination Product		
5	NDC:25000-135-93	1 in 1 BOX	11/06/2015	
5		37470 in 1 BAG; Type 0: Not a Combination Product		

5		Product	
6	NDC:25000-135-70	500 in 1 BOTTLE; Type 0: Not a Combination Product	11/06/2015
7	NDC:25000-135-02	1 in 1 CARTON	12/20/2019
7		24 in 1 BOTTLE; Type 0: Not a Combination Product	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091237	11/06/2015	

IBUPROFEN

ibuprofen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:25000-136
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	ORANGE	Score	no score
Shape	ROUND (Round Shaped)	Size	15mm
Flavor		Imprint Code	115
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25000-136-03	30 in 1 BOTTLE; Type 0: Not a Combination Product	11/06/2015	
2	NDC:25000-136-14	1000 in 1 BOTTLE; Type 0: Not a Combination Product	11/06/2015	
3	NDC:25000-136-30	6 in 1 BOX	11/06/2015	
3		6500 in 1 BAG; Type 0: Not a Combination Product		
4	NDC:25000-136-98	1 in 1 BOX	11/06/2015	
4		34065 in 1 BAG; Type 0: Not a Combination Product		
5	NDC:25000-136-93	1 in 1 BOX	11/06/2015	
5		37470 in 1 BAG; Type 0: Not a Combination Product		
6	NDC:25000-136-69	300 in 1 BOTTLE; Type 0: Not a Combination Product	11/06/2015	
7	NDC:25000-136-71	1000 in 1 BOTTLE; Type 0: Not a Combination Product	11/06/2015	
8	NDC:25000-136-02	1 in 1 CARTON	12/20/2019	
8		24 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091237	11/06/2015	

Labeler - MARKSANS PHARMA LIMITED (925822975)

Establishment			
Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	MANUFACTURE(25000-135, 25000-136)

Revised: 12/2021

MARKSANS PHARMA LIMITED