

**SUL-Q-NOX- sulfaquinoxaline solution**  
**Huvepharma, Inc**

-----  
**Sul-Q-Nox®**

**31.92% (Sulfaquinoxaline as sodium and potassium salts)**

ANTIMICROBIAL

**Caution: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.**

**For Use in Animal Drinking Water Only**  
**Sulfonamide**

**KEEP OUT OF REACH OF CHILDREN**  
**KEEP TIGHTLY SEALED WHEN NOT IN USE**

Restricted Drug (California) - Use only as directed. Not for human use.

**Net Contents: 1 Gallon (128 fl. oz.)**

Manufactured for:  
Huvepharma, Inc.  
525 Westpark Drive, Suite 230  
Peachtree City, GA 30269

**HUVEPHARMA®**

HUVEPHARMA and Sul-Q-Nox are registered trademarks of Huvepharma EOOD



Approved by FDA under NADA # 006-891  
P08-4001SL  
Rev. 01-2022

**CAUTION:**

- May cause toxic reactions unless drug is evenly mixed in water at dosages indicated and used according to directions.
- Levels of sulfaquinoxaline higher than 0.025% in feed or 0.012% in water for more than 24 to 36 hours may result in reduced growth rate in chickens as a result of reduced feed or water intake.
- For control of disease outbreaks, medication should be initiated as soon as diagnosis is determined.
- Treated animals must actually consume enough medicated water to provide a necessary dosage of

approximately 10 to 45 mg/lb/day in chickens and 3.5 to 55 mg/lb/day in turkeys and 6 mg/lb/day in cattle and calves, depending on class of animal, ambient temperature, age, and other factors.

- Prolonged administration of sulfaquinoxaline at higher doses may result in depressed feed or water intake, deposition of sulfaquinoxaline crystals in kidney and interference with normal blood clotting.

- Do not mix or administer in galvanized containers.

- The following word is required on this product to comply with Agricultural Code of California:

**HAZARDOUS**

- **CAUSES SKIN AND EYE BURNS.** Avoid contact with eyes, skin, or clothing. In case of contact, flush immediately with water for at least 15 minutes; for eye get medical attention.

**WARNING:**

- Withdraw use of product 10 days before slaughter.
- A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.
- Do not medicate chickens or turkeys producing eggs for human consumption.
- Not for use in lactating dairy cattle.
- **DISPOSE OF ANY WASTE OR UNUSED PORTION PROPERLY.**
- **KEEP ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.**
- Store between 20°- 25°C (68°- 77°F); excursions permitted between 15°- 40°C (59°- 104°F).

**FOR AUTOMATIC DRINKING WATER PROPORTIONERS**

**FOR CONTROL OR TREATMENT OF THE FOLLOWING DISEASES WHEN CAUSED BY ONE OR MORE PATHOGENIC ORGANISMS SUSCEPTIBLE TO SULFAQUINOXALINE  
PREPARE MEDICATED DRINKING WATER FRESH DAILY**

INDICATIONS	DOSAGE PER 128 GALLONS OF WATER	TREATMENT
<p><b>COCCIDIOSIS — CHICKENS:</b> Control of coccidiosis caused by <i>Eimeria tenella</i>, <i>E. necatrix</i>, <i>E. maxima</i>, <i>E. brunetti</i>, <i>E. acervulina</i>, susceptible to sulfaquinoxaline.</p>	<p><b>20½ fl. oz. (0.04%)</b></p> <p><b>13 fl. oz. (0.025%)</b></p>	<p>Give for 2-3 days, skip 3 days. <b>Then</b> Give for 2 days more. If bloody droppings appear, repeat treatment at this level for 2 more days. <b>CAUTION:</b> Do not change litter unless absolutely necessary. Do not give flushing mashes. Give for 2 days - skip 3 days - give for 2 days - skip 3 days and give 2 more days. Repeat if</p>
<p><b>COCCIDIOSIS — TURKEYS:</b> Control of coccidiosis caused</p>		

by *Eimeria meleagridis*, *E. adenoides*, susceptible to sulfaquinoxaline.

**13 fl. oz. (0.025%)**

**ACUTE FOWL CHOLERA — CHICKENS AND TURKEYS:**

Control of acute fowl cholera caused by *Pasteurella multocida* susceptible to sulfaquinoxaline.

**20½ fl. oz. (0.04%)**

**FOWL TYPHOID — CHICKENS AND TURKEYS:**

Control of fowl typhoid caused by *Salmonella gallinarum*, susceptible to sulfaquinoxaline.

**COCCIDIOSIS — CATTLE AND CALVES:**

For the control and treatment of coccidiosis caused by *Eimeria bovis*, *E. zuernii*, susceptible to sulfaquinoxaline.

**7¾ fl. oz. (0.015%)  
6 mg/lb of body wt.**

necessary. **CAUTION:** Do not change litter unless absolutely necessary. Do not give flushing mashes.

Use for 2-3 days. Move birds to clean ground. If disease recurs, repeat treatment. Poultry that have survived fowl typhoid outbreaks should not be kept for laying house replacements or breeders unless tests show they are not carriers.

Give for 3-5 days. As a generalization, cattle and calves will consume approx. 1 gal/100 lbs body weight/day. Cattle or calves not eating or drinking must be treated individually.

**To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Huvepharma, Inc. at 1-877-994-4883 or [www.huvepharma.us](http://www.huvepharma.us). For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.**

<p><b>CAUTION:</b></p> <ul style="list-style-type: none"> <li>May cause toxic reactions unless drug is evenly mixed in water at dosages indicated and used according to directions.</li> <li>Levels of sulfaquinoxaline higher than 0.025% in feed or 0.012% in water for more than 24 to 36 hours may result in reduced growth rate in chickens as a result of reduced feed or water intake.</li> <li>For control of disease outbreaks, medication should be initiated as soon as diagnosis is determined.</li> <li>Treated animals must actually consume enough medicated water to provide a necessary dosage of approximately 10 to 25 mg/lb/day in chickens and 3.5 to 50mg/lb/day in turkeys and 6 mg/lb/day in cattle and calves, depending on class of animal, ambient temperature, age, and other factors.</li> <li>Prolonged administration of sulfaquinoxaline at higher doses may result in depressed feed or water intake, deposition of sulfaquinoxaline crystals in kidney and interference with normal blood clotting.</li> <li>Do not mix or administer in galvanized containers.</li> <li>The following word is required on this product to comply with the Agricultural Code of California: <b>HAZARDOUS</b></li> <li><b>CAUSES SKIN AND EYE BURNS:</b> Avoid contact with eyes, skin, or clothing. In case of contact, flush immediately with water for at least 15 minutes; for eye get medical attention.</li> </ul> <p><b>WARNING:</b></p> <ul style="list-style-type: none"> <li>Withdraw use of product 10 days before slaughter.</li> <li>A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.</li> <li>Do not medicate chickens or turkeys producing eggs for human consumption.</li> <li>Not for use in lactating dairy cattle.</li> <li><b>DISPOSE OF ANY WASTE OR UNUSED PORTION PROPERLY.</b></li> <li>KEEP ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.</li> <li>Store between 20°-25°C (68°-77°F); excursions permitted between 15°-40°C (59°-104°F).</li> </ul>	<p style="text-align: center;"><b>Sul-Q-Nox<sup>®</sup></b></p> <p style="text-align: center;"><b>31.92% (Sulfaquinoxaline as sodium and potassium salts)</b></p> <p style="text-align: center;">ANTIMICROBIAL</p> <p><b>CAUTION:</b> Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.</p> <p><b>For Use in Animal Drinking Water Only</b></p> <p>Sulfonamide</p> <p><b>KEEP OUT OF REACH OF CHILDREN</b></p> <p><b>KEEP TIGHTLY SEALED WHEN NOT IN USE</b></p> <p>Restricted Drug (California) – Use only as directed. Not for human use.</p> <p>Net Contents: 1 Gallon (128 fl. oz.)</p> <p><small>Manufactured for: Huvepharma, Inc. 525 Westpark Drive, Suite 230 Pasadena, CA 91107</small></p> <p><small>HUVEPHARMA AND SUL-Q-NOX are registered trademarks of Huvepharma, Inc.</small></p> <p><small>Approved by FDA under NDND 001 001-20015 001-01-002</small></p>	<p style="text-align: right;"><b>FOR A FOR CONTROL OR OR MORE PAI PREPA</b></p> <p style="text-align: center;"><b>INDICATIONS</b></p> <p><b>COCCIDIOSIS — CHICKENS:</b> Control of coccidiosis caused by <i>meleagridis</i>, <i>E. adenoides</i>, <i>E. zuernii</i>, susceptible to sulfaquinoxaline.</p> <p><b>COCCIDIOSIS — TURKEYS:</b> Control of coccidiosis caused by <i>meleagridis</i>, <i>E. adenoides</i>, <i>E. zuernii</i>, susceptible to sulfaquinoxaline.</p> <p><b>ACUTE FOWL CHOLERA — CHICKENS:</b> Control of acute fowl cholera caused by <i>Pasteurella multocida</i>, susceptible to sulfaquinoxaline.</p> <p><b>FOWL TYPHOID — CHICKENS AND TURKEYS:</b> Control of fowl typhoid caused by <i>Salmonella gallinarum</i>, susceptible to sulfaquinoxaline.</p> <p><b>COCCIDIOSIS — CATTLE AND CALVES:</b> For the control and treatment of coccidiosis caused by <i>Eimeria bovis</i>, <i>E. zuernii</i>, susceptible to sulfaquinoxaline.</p> <p><small>To report suspected adverse drug events, contact FDA at 1-888-FDA-VETS or <a href="http://www.fda.gov/reportanimalae">http://www.fda.gov/reportanimalae</a>.</small></p>
---	--	---

**SUL-Q-NOX**  
sulfaquinoxaline solution

**Product Information**

<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:23243-6771
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>SULFAQUINOXALINE</b> (UNII: WNW8115TM9) (SULFAQUINOXALINE - UNII:WNW8115TM9)	SULFAQUINOXALINE	.3192 g in 1 mL

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:23243-6771-2	4 in 1 BOX		
1		3785 mL in 1 PACKAGE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA006891	07/09/2019	

**Labeler** - Huvepharma, Inc (619153559)

**Registrant** - Huvepharma EOOD (552671651)

### Establishment

Name	Address	ID/FEI	Business Operations
Huvepharma, Inc		883128204	analysis, label, manufacture, pack

Revised: 10/2023

Huvepharma, Inc