

DEHORNING - calcium hydroxide, sodium hydroxide paste
H. W. Naylor Company Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

DEHORNING PASTE

Active Ingredients

Calcium hydroxide 37.8%, Sodium hydroxide 24.9%

Purpose

PREVENTS HORN GROWTH

Directions

Read package insert completely before dehorning young calves, older calves, sheep or goats.

CALVES: As soon as horn button can be felt, when calf is three to seven days old, remove hair over horn button with shears or clippers. Apply Dehorning Paste with wood applicator once only to a spot the size of a quarter (25mm). Keep calf tied away from other animals, out of rain for at least six hours.

Safety

FOR EXTERNAL USE ONLY

FOR ANIMAL USE ONLY

Keep out of the reach of children

Other Ingredients

Iron Oxide Color
Glycerin

Questions?

www.drnaylor.com

Package Insert

Directions for Using

Dr. Naylor® DeHorning Paste

Read this sheet completely before dehorning any animal.

WHEN TO DEHORN

Dr. Naylor Dehorning Paste is best used on calves up to eight (8) weeks of age. It is most easily used on very young calves. The younger the calf is when treated, the better. Shock and pain are negligible and dehorning is quickly and easily done. Dehorning Paste may be used in most areas at any time of the year; there is no cutting, no bleeding, no fly problems!

HOW TO DEHORN

ALL ANIMALS: Be careful to apply a thin, even coating of Dehorning Paste, and only as directed. Remove hair over horn button (bud) with shears or clippers in order to see that application is in the right place and in direct contact with the skin. To contain the paste, apply a ring of Dr. Naylor® Udder Balm or petroleum jelly beyond the Paste application area. If treated calves are confined in pens, tie them away from other animals to prevent rubbing paste on one another. Do not allow treated animals to nurse. Keep animals out of rain after Dehorning Paste has been applied.

YOUNG CALVES: As soon as horn button (bud) can be felt, remove hair over horn button and adjacent area with shears or clippers. On animals under one (1) week old apply a thin even film of Dehorning Paste over horn button and on a small ring of adjacent skin (horn growth epithelium) about the size of a five-cent coin (20 mm). On animals over seven days old apply Paste to a larger spot, the size of a quarter (25mm). Use a wood applicator and apply paste once only.

OLDER CALVES: To dehorn calves six to eight weeks old whose horn buttons have started to grow, remove hair and use a small scraper or file to roughen a ring of skin about 3/8 inches (10mm) wide around the base of the horn button. Do not draw blood. Apply a thin even film of Paste over the horn button and the roughened ring around the base of the horn button. Apply once only.

LAMBS, KID GOATS: Apply Dehorning Paste as soon as possible after birth if animal is in good physical condition. Minimum discomfort is experienced and application is facilitated. Application of the Paste on lambs and kids over two weeks old is generally too late to prevent future horn growth as horns are too large, especially in males. Buck kids (males) usually have palpable horn buttons (buds) at birth. Kid does (females) are a little slower in developing horns. Clip the hair over and around horn buttons. Apply a thin even film of Dehorning Paste over horn buttons and a narrow ring of skin at the base of the horn button. This ring of skin grows the horn. Too thick an application may result in deeper penetration and activity with discomfort. Dehorn dairy bucks at 3 to 4 days, dairy goes at 4 days and always before 10 days. Pygmy bucks should be dehorned at 7 days, pygmy does at 7 to 10 days. Twenty to thirty minutes after Paste application scrape off Paste and neutralize treated area by flooding with water, then vinegar. Wash water and vinegar must not come in contact with animal's eyes. Kids and lambs may be wrapped in a blanket for a short period to prevent rubbing treated area with their hooves.

WARNING

For external veterinary use only. This product is a form of caustic. Apply only with a

wood applicator and as directed. Do not ingest. Avoid eye and skin contact. Use gloves to protect hands. Keep bottle tightly capped and in a safe place.

KEEP FROM REACH OF CHILDREN

www.drnaylor.com
www.dehorning.com

H.W. Naylor Company, Inc.
Morris, NY 13808-019 U.S.A.

Save directions for future use.

Principal Display Panel

Dr. Naylor[®]

DEHORNING PASTE

PREVENTS HORN GROWTH

Net Weight: 4 Oz. (114 grams)

H.W. NAYLOR COMPANY, INC.
Morris, N.Y. 13808-0190 U.S.A.

www.drnaylor.com

POISON

WARNING: This product is caustic. Keep jar closed and store out of the reach of children. Apply with a wood applicator. Do not ingest. Avoid contact with eyes or skin.

FIRST AID:

EXTERNAL: Flood with water for extended period then wash with vinegar.

INTERNAL: Give large amounts of water or milk for dilution, juice of lemon, grapefruit or orange. Follow with olive oil.

EYES: Rinse copiously with water immediately. Wash with 5% boric acid solution.

SEEK MEDICAL ATTENTION

<p>POISON WARNING: This product is caustic. Keep jar closed and store out of the reach of children. Apply with a wood applicator. Do not ingest. Avoid contact with eyes or skin.</p> <p>FIRST AID: EXTERNAL: Flood with water for extended period then wash with vinegar. INTERNAL: Give large amounts of water or milk for dilution, juice of lemon, grapefruit or orange. Follow with olive oil. EYES: Rinse copiously with water immediately. Wash with 5% boric acid solution.</p> <p>SEEK MEDICAL ATTENTION</p>	 <h1>DEHORNING PASTE</h1> <h2>PREVENTS HORN GROWTH</h2> <p>ACTIVE INGREDIENTS Calcium hydroxide 37.8%, Sodium hydroxide 24.9%</p> <p>Net Weight: 4 Oz. (114 grams) H.W. NAYLOR COMPANY, INC. Morris, N.Y. 13808-0190 U.S.A. www.drnaylor.com</p>	<p>DIRECTIONS Read package insert completely before dehorning young calves, older calves, sheep or goats. CALVES: As soon as horn button can be felt, when calf is three to seven days old, remove hair over horn button with shears or clippers. Apply Dehorning Paste with wood applicator once only to a spot the size of a quarter (25mm). Keep calf tied away from other animals, out of rain for at least six hours.</p> <p>FOR EXTERNAL USE ONLY FOR ANIMAL USE ONLY Made in the U.S.A.</p>
--	---	---

DEHORNING

calcium hydroxide, sodium hydroxide paste

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:15878-311
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Calcium Hydroxide (UNII: PF5DZW74VN) (Calcium Hydroxide - UNII:PF5DZW74VN)	Calcium Hydroxide	63 mg in 1 g
Sodium Hydroxide (UNII: 55X04QC32I) (Hydroxide Ion - UNII:9159UV381P)	Sodium Hydroxide	41 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
Ferric Oxide Red (UNII: 1K09F3G675)	
Glycerin (UNII: PDC6A3C0OX)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:15878-311-24	114 g in 1 JAR		

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End

Category	Citation	Date	Date
unapproved drug other		01/01/1964	

Labeler - H. W. Naylor Company Inc. (002080190)

Establishment

Name	Address	ID/FEI	Business Operations
H. W. Naylor Company Inc.		002080190	MANUFACTURE, api manufacture

Revised: 11/2023

H. W. Naylor Company Inc.