

IBUPROFEN ORAL - ibuprofen oral suspension

Strategic Sourcing Services

Ibuprofen Oral Suspension USP, 100 mg/5 mL (OTC)

ACTIVE INGREDIENT(S)

Ibuprofen 100 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

PURPOSE

Pain reliever/ fever reducer

USE(S)

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

WARNINGS

Allergy alert:Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives ■ facial swelling ■ asthma (wheezing)
- shock ■ skin reddening ■ rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning:This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Heart attack and stroke warning:NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning:Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless

directed by doctor.

DO NOT USE

- if the child has ever had an allergic reaction to Ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

ASK A DOCTOR BEFORE USE IF

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

ASK A DOCTOR OR PHARMACIST BEFORE USE IF

- under a doctor's care for any serious condition
- taking any other drug

WHEN USING THIS PRODUCT

- take with food or milk if stomach upset occurs

STOP USE AND ASK DOCTOR IF

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

KEEP OUT OF REACH OF CHILDREN

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

DIRECTIONS

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb) (mL)*	Age (yr)	Dose	Weight (lb) (mL)*	Age (yr)	Dose	Weight (lb) (mL)*	Age (yr)	Dose
under 24	under 2 years	ask a doctor	under 24	under 2 years	ask a doctor	under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL	24-35 lbs	2-3 years	5 mL	24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL	36-47 lbs	4-5 years	7.5 mL	36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL	48-59 lbs	6-8 years	10 mL	48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL	60-71 lbs	9-10 years	12.5 mL	60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL	72-95 lbs	11 years	15 mL	72-95 lbs	11 years	15 mL

**or as directed by a doctor*

OTHER INFORMATION

- each 5 mL contains : sodium 2 mg
- store between 20-25°C (68-77°F)
- do not use if carton is opened or printed bottle neckband is broken or missing
- see bottom panel for lot number and expiration date

INACTIVE INGREDIENT

Acesulfame potassium, anhydrous citric acid, carboxymethylcellulose sodium, D&C Yellow # 10, FD&C Red #40, flavors, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum

QUESTIONS or COMMENTS?

Call 833-358-6431 Monday to Friday 9:00am to 7:00pm EST

©2023 McKesson Corporation

Distributed by:McKesson Corp., via Strategic Sourcing Services LLC.

Memphis, TN 38141

Money Back Guarantee

www.fosterandthrive.com

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Motrin[®].

PRINCIPAL DISPLAY PANEL

Ibuprofen oral suspension berry flavor container carton-118 mL

Foster & Thrive™
Children's
Ibuprofen

Oral Suspension (NSAID)
100 mg per 5 mL
PAIN RELIEVER /
FEVER REDUCER

NDC 70677-1115-1

COMPARE TO THE
 ACTIVE INGREDIENT IN
 CHILDREN'S MOTRIN®**

Foster & Thrive™
Children's
Ibuprofen

Oral Suspension (NSAID)
100 mg per 5 mL
PAIN RELIEVER /
FEVER REDUCER

Lasts Up to 8 Hours

For Ages 2 to 11 Years
ORIGINAL BERRY
FLAVOR
 DOSING CUP INCLUDED

SEE NEW WARNINGS

4 FL OZ (118 mL)

BX-161



LOT:

EXP:

Drug Facts (continued)

- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance

Drug Facts (continued)

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

*or as directed by a doctor

Other information

- each 5 mL contains : sodium 2 mg
- store between 20-25°C (68-77°F)
- **do not use if carton is opened or printed bottle neckband is broken or missing**
- see bottom panel for lot number and expiration date

Inactive ingredients

Acetosulfame potassium, anhydrous citric acid, carboxymethylcellulose sodium, D&C Yellow # 10, FD&C Red #40, flavors, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum

Questions or comments?

Call **833-358-6431** Monday to Friday 9:00am to 7:00pm EST

©2023 McKesson Corporation
 Distributed by: McKesson Corp., via Strategic Sourcing Services LLC.
 Memphis, TN 38141
 Money Back Guarantee
 www.fosterandthrive.com

** This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Motrin®.

Iss. 04/2023

IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USING.
KEEP THIS BOX FOR IMPORTANT INFORMATION.
THIS PRODUCT IS INTENDED FOR USE IN CHILDREN.

Drug Facts

Active ingredient **Purpose**
 (in each 5 mL)

Ibuprofen 100 mg (NSAID)† Pain reliever/
 †nonsteroidal anti-inflammatory drug fever reducer

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed
- **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
- **Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids

Ibuprofen oral suspension original berry flavor with dye container carton-118 mL

Foster & Thrive™
Children's
Ibuprofen

Oral Suspension (NSAID)
 100 mg per 5 mL
**PAIN RELIEVER /
 FEVER REDUCER**

Drug Facts (continued)

- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance ▶

Drug Facts (continued)

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

*or as directed by a doctor

Other information

- each 5 mL contains : sodium 2 mg
- store between 20-25°C (68-77°F)
- **do not use if carton is opened or printed bottle neckband is broken or missing**
- see bottom panel for lot number and expiration date

Inactive ingredients

Acetaminophen potassium, anhydrous citric acid, carboxymethylcellulose sodium, flavors, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum

Questions or comments?

Call **833-358-6431** Monday to Friday 9:00am to 7:00pm EST

©2023 McKesson Corporation
 Distributed by: McKesson Corp., via Strategic Sourcing Services LLC.
 Memphis, TN 38141
 Money Back Guarantee
 www.fosterandthrive.com

** This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Motrin®.

Iss. 04/2023

NDC 70677-1116-1

COMPARE TO THE
 ACTIVE INGREDIENT IN
 CHILDREN'S MOTRIN®**

Foster & Thrive™
Children's
Ibuprofen

Oral Suspension (NSAID)
 100 mg per 5 mL
**PAIN RELIEVER /
 FEVER REDUCER**

Lasts Up to 8 Hours

Dye Free

 **For Ages 2 to 11 Years**
ORIGINAL BERRY
FLAVOR
 DOSING CUP INCLUDED

SEE NEW WARNINGS

4 FL OZ (118 mL)

BX-164



LOT:

EXP:

IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USING.
KEEP THIS BOX FOR IMPORTANT INFORMATION.
THIS PRODUCT IS INTENDED FOR USE IN CHILDREN.

Drug Facts

Active ingredient **Purpose**
 (in each 5 mL)

Ibuprofen 100 mg (NSAID)* Pain reliever/
 *nonsteroidal anti-inflammatory drug fever reducer

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ takes more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids ▶

Ibuprofen oral suspension bubble gum flavor container carton-118 mL

Foster & Thrive™
Children's
Ibuprofen

Oral Suspension (NSAID)
100 mg per 5 mL
PAIN RELIEVER /
FEVER REDUCER

NDC 70677-1117-1

COMPARE TO THE
 ACTIVE INGREDIENT IN
 CHILDREN'S MOTRIN®**

Foster & Thrive™
Children's
Ibuprofen

Oral Suspension (NSAID)
100 mg per 5 mL
PAIN RELIEVER /
FEVER REDUCER

Lasts Up to 8 Hours

 **For Ages 2 to 11 Years**
BUBBLE
GUM FLAVOR
 DOSING CUP INCLUDED

SEE NEW WARNINGS

4 FL OZ (118 mL)

BX-162



LOT:

EXP:

Drug Facts (continued)

- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is
 ■ under a doctor's care for any serious condition
 ■ taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:

- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away: (1-800-222-1222)

Directions

- **this product does not contain directions or complete warnings for adult use**

- do not give more than directed
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance ▶

Drug Facts (continued)

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

*or as directed by a doctor

Other information

- each 5 mL contains : sodium 2 mg
- store between 20-25°C (68-77°F)
- do not use if carton is opened or printed bottle neckband is broken or missing
- see bottom panel for lot number and expiration date

Inactive ingredients

Acetosulfame potassium, anhydrous citric acid, carboxymethylcellulose sodium, D&C Red #93, FD&C Red #40, flavors, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum

Questions or comments?

Call **833-358-6431** Monday to Friday 9:00am to 7:00pm EST

©2023 McKesson Corporation

Distributed by: McKesson Corp., via Strategic Sourcing Services LLC.

Memphis, TN 38141

Money Back Guarantee

www.fosterandthrive.com

** This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Motrin®.

Iss. 04/2023

IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USING.
 KEEP THIS BOX FOR IMPORTANT INFORMATION.
 THIS PRODUCT IS INTENDED FOR USE IN CHILDREN.

Drug Facts

Active ingredient **Purpose**
 (in each 5 mL)

Ibuprofen 100 mg (NSAID) Pain reliever/
 nonsteroidal anti-inflammatory drug fever reducer

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to Ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids ▶

Ibuprofen oral suspension grape flavor container carton-118 mL

Foster & Thrive™
Children's
Ibuprofen
 Oral Suspension (NSAID)
 100 mg per 5 mL
**PAIN RELIEVER /
 FEVER REDUCER**

NDC 70677-1118-1

COMPARE TO THE
 ACTIVE INGREDIENT IN
 CHILDREN'S MOTRIN®**

Foster & Thrive™
Children's
Ibuprofen

Oral Suspension (NSAID)
 100 mg per 5 mL
**PAIN RELIEVER /
 FEVER REDUCER**

Lasts Up to 8 Hours

 **For Ages 2 to 11 Years**
GRAPE FLAVOR
 DOSING CUP INCLUDED

SEE NEW WARNINGS

4 FL OZ (118 mL)

BX-163



LOT:

EXP:

Drug Facts (continued)

- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - has bloody or black stools
 - has stomach pain that does not get better
 - child has symptoms of heart problems or stroke:
 - chest pain
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
 - the child does not get any relief within first day (24 hours) of treatment
 - fever or pain gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance ▶

Drug Facts (continued)

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

* or as directed by a doctor

Other information

- each 5 mL contains: sodium 2 mg
- store between 20-25°C (68-77°F)
- **do not use if carton is opened or printed bottle neckband is broken or missing**
- see bottom panel for lot number and expiration date

Inactive ingredients

Acesulfame potassium, anhydrous citric acid, carboxymethylcellulose sodium, D&C Red #33, FD&C Blue #1, flavors, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum

Questions or comments?

Call **833-358-6431** Monday to Friday 9:00am to 7:00pm EST

©2023 McKesson Corporation
 Distributed by: McKesson Corp., via Strategic Sourcing Services LLC.
 Memphis, TN 38141
 Money Back Guarantee
 www.fosterandthrive.com

** This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Motrin®.

Iss. 04/2023

IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USING. KEEP THIS BOX FOR IMPORTANT INFORMATION. THIS PRODUCT IS INTENDED FOR USE IN CHILDREN.

Drug Facts

Active ingredient Purpose
 (in each 5 mL)

Ibuprofen 100 mg (NSAID) Pain reliever/
 *nonsteroidal anti-inflammatory drug fever reducer

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash

■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
 - takes a blood thinning (anticoagulant) or steroid drug
 - takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 - takes more or for a longer time than directed
- Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
- Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids ▶

Ibuprofen oral suspension berry original flavor container carton-237 mL

Foster & Thrive™

Children's

Ibuprofen

Oral Suspension (NSAID)
100 mg per 5 mL
PAIN RELIEVER /
FEVER REDUCER

NDC 70677-1115-2

COMPARE TO THE
ACTIVE INGREDIENT IN
CHILDREN'S MOTRIN***

Foster & Thrive™

Children's

Ibuprofen

Oral Suspension (NSAID)
100 mg per 5 mL
PAIN RELIEVER /
FEVER REDUCER

Lasts Up to 8 Hours



For Ages 2 to 11 Years
ORIGINAL BERRY
FLAVOR
DOSING CUP INCLUDED

SEE NEW WARNINGS

8 FL OZ (237 mL)

NDC 70677-1115-2

COMPARE TO THE
ACTIVE INGREDIENT IN
CHILDREN'S MOTRIN***

Foster & Thrive™

Children's

Ibuprofen

Oral Suspension (NSAID)
100 mg per 5 mL
PAIN RELIEVER /
FEVER REDUCER

Lasts Up to 8 Hours



For Ages 2 to 11 Years
ORIGINAL BERRY
FLAVOR
DOSING CUP INCLUDED

SEE NEW WARNINGS

8 FL OZ (237 mL)

**IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USING.
KEEP THIS BOX FOR IMPORTANT INFORMATION.
THIS PRODUCT IS INTENDED FOR USE IN CHILDREN.**

Drug Facts

Active ingredient (in each 5 mL) Purpose
Ibuprofen 100 mg (NSAID) Pain reliever/
nonsteroidal anti-inflammatory drug fever reducer

Uses temporarily:
relieves minor aches and pains due to the common cold, flu,
sore throat, headache and toothache
reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction,
especially in people allergic to aspirin. Symptoms may include:
hives facial swelling asthma (wheezing)
shock skin reddening rash ulcers

If an allergic reaction occurs, stop use and seek medical help right
away.

Stomach bleeding warning: This product contains an NSAID,
which may cause severe stomach bleeding. The chance is higher if
your child:
has had stomach ulcers or bleeding problems
takes a blood thinning (anticoagulant) or steroid drug
takes other drugs containing prescription or
nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

takes more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase
the risk of heart attack, heart failure, and stroke. These can be fatal.
The risk is higher if you use more than directed or for longer than
directed.

Sore throat warning: Severe or persistent sore throat or sore throat
accompanied by high fever, headache, nausea, and vomiting may
be serious. Consult doctor promptly. Do not use more than 2 days
or administer to children under 3 years of age unless directed by
doctor.

Do not use

if the child has ever had an allergic reaction to Ibuprofen or any
other pain reliever/fever reducer
right before or after heart surgery

Ask a doctor before use if

stomach bleeding warning applies to your child
child has a history of stomach problems, such as heartburn
child has problems or serious side effects from taking pain
relievers or fever reducers
child has not been drinking fluids
child has lost a lot of fluid due to vomiting or diarrhea
child has high blood pressure, heart disease, liver cirrhosis,
kidney disease, or had a stroke
child has asthma
child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

under a doctor's care for any serious condition
taking any other drug

When using this product

take with food or milk if stomach upset occurs

Stop use and ask a doctor if

child experiences any of the following signs of stomach
bleeding:
feels faint vomits blood
has bloody or black stools
has stomach pain that does not get better
child has symptoms of heart problems or stroke:
chest pain trouble breathing

Drug Facts (continued)

weakness in one part or side of body
slurred speech leg swelling
the child does not get any relief within first day (24 hours) of
treatment
fever or pain gets worse or lasts more than 3 days
redness or swelling is present in the painful area
any new symptoms appear

Keep out of reach of children. In case of overdose, get medical
help or contact a Poison Control Center right away.
(1-800-222-1222)

Directions

**this product does not contain directions or complete warnings
for adult use**
do not give more than directed
shake well before using
mL = milliliter
find right dose on chart. If possible, use weight to dose;
otherwise use age.
use only enclosed dosing cup. Do not use any other dosing
device.
if needed, repeat dose every 6-8 hours
do not use more than 4 times a day
replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

*or as directed by a doctor

Other information

each 5 mL contains : sodium 2 mg
store between 20-25°C (68-77°F)
do not use if carton is opened or printed bottle neckband is
broken or missing
see bottom panel for lot number and expiration date

Inactive ingredients

Acetosulfame potassium, anhydrous citric acid,
carboxymethylcellulose sodium, D&C Yellow #10, FD&C Red #40,
flavors, glycerin, microcrystalline cellulose, polysorbate 80,
propylene glycol, purified water, sodium benzoate, sucrose, and
xanthan gum

Questions or comments?

Call 833-358-6431 Monday to Friday 9:00am to 7:00pm EST

©2023 McKesson Corporation

Distributed by: McKesson Corp., via Strategic Sourcing Services LLC.
Memphis, TN 38141
Money Back Guarantee
www.fosterandthrive.com

*** This product is not manufactured or distributed by McNeil Consumer
Healthcare, distributor of Children's Motrin**.

Iss. 04/2023

BX-171



LOT:

EXP:

IBUPROFEN ORAL

ibuprofen oral suspension

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:70677-1115

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	100 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
GLYCERIN (UNII: PDC6A3C0OX)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	orange	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70677-1115-1	1 in 1 CARTON	08/09/2023	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:70677-1115-2	1 in 1 CARTON	08/09/2023	
2		237 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210602	08/09/2023	

IBUPROFEN ORAL

ibuprofen oral suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70677-1116
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	100 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
GLYCERIN (UNII: PDC6A3C0OX)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	

Product Characteristics

Color	purple	Score	
Shape		Size	
Flavor	BERRY (original berry flavor)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70677-1116-1	1 in 1 CARTON	08/09/2023	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
---------------------------	---	-----------------------------	---------------------------

ANDA	ANDA210602	08/09/2023	
------	------------	------------	--

IBUPROFEN ORAL

ibuprofen oral suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70677-1117
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	100 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
GLYCERIN (UNII: PDC6A3C0OX)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	

Product Characteristics

Color	pink	Score	
Shape		Size	
Flavor	BUBBLE GUM	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70677-1117-1	1 in 1 CARTON	08/09/2023	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210602	08/09/2023	

IBUPROFEN ORAL

ibuprofen oral suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70677-1118
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	100 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
GLYCERIN (UNII: PDC6A3C0OX)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
D&C RED NO. 33 (UNII: 9DBA05BB0L)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:70677-1118-1	1 in 1 CARTON	08/09/2023	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA210602		08/09/2023	

Labeler - Strategic Sourcing Services (116956644)

Establishment			
Name	Address	ID/FEI	Business Operations
AptaPharma Inc.		790523323	manufacture(70677-1115, 70677-1116, 70677-1117, 70677-1118)

Revised: 8/2023

Strategic Sourcing Services