

**SINUS PRESSURE AND PAIN DAYTIME NIGHTTIME MAXIMUM STRENGTH-
acetaminophen, diphenhydramine hcl, guaifenesin, phenylephrine hcl
Walgreen Company**

Walgreens 44-615694-10-SMH

Active ingredients (in each caplet) (Sinus day)

Acetaminophen 325 mg
Guaifenesin 200 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever
Expectorant
Nasal decongestant

Active ingredients (in each caplet) (Sinus night)

Acetaminophen 325 mg
Diphenhydramine HCl 12.5 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever
Antihistamine/cough suppressant
Nasal decongestant

Uses

- temporarily relieves these common cold symptoms:
 - nasal congestion
 - headache
 - minor aches and pains
 - sinus congestion and pressure
 - cough (**Nighttime only**)
 - runny nose and sneezing (**Nighttime only**)
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive (**Daytime only**)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- with any other product containing diphenhydramine, even one used on skin
(Nighttime only)

Ask a doctor before use if you have

- heart disease
- high blood pressure
- liver disease
- thyroid disease
- diabetes
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis **(Nighttime only)**
- glaucoma **(Nighttime only)**

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers **(Nighttime only)**

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children **(Nighttime only)**
- marked drowsiness may occur **(Nighttime only)**
- alcohol, sedatives, and tranquilizers may increase drowsiness **(Nighttime only)**
- use caution when driving a motor vehicle or operating machinery **(Nighttime only)**

- avoid alcoholic beverages (**Nighttime only**)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Do not take DAYTIME and NIGHTTIME products at the same time.

Directions

- **do not take more than directed**
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years and over: take 2 caplets every 4 hours
- children under 12 years: do not use

Other information

- **each caplet contains:** sodium 3 mg (**Daytime only**)
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients (Daytime only)

corn starch, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Inactive ingredients (Nighttime only)

corn starch, croscarmellose sodium, crospovidone, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, iron oxide yellow, magnesium stearate, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol,

povidone, silicon dioxide, sodium bicarbonate, stearic acid, talc, titanium dioxide

Questions or comments?

1-800-426-9391

Principal display panel

DAY & NIGHT PACK NDC 0363-6156-10

• WALGREENS •
PHARMACIST RECOMMENDED†
Walgreens

DAYTIME Sinus Pressure & Pain ACETAMINOPHEN PAIN RELIEVER GUAIFENESIN EXPECTORANT PHENYLEPHRINE HCl NASAL DECONGESTANT Maximum Strength 24 CAPLETS ACTUAL SIZE	NIGHTTIME Sinus Pressure & Pain ACETAMINOPHEN PAIN RELIEVER DIPHENHYDRAMINE HCl ANTIHISTAMINE COUGH SUPPRESSANT PHENYLEPHRINE HCl NASAL DECONGESTANT Maximum Strength 16 CAPLETS ACTUAL SIZE
---	--

TOTAL 40 CAPLETS

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS
TORN,
BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

**Do Not Take Daytime
and Nighttime Products
at the Same Time.**

†Our pharmacists recommend the Walgreens brand.
We invite you to compare to national brands.

50844 REV0723A61569410

DISTRIBUTED BY: **WALGREEN CO.**
DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com
©2023 Walgreen Co.

ADHESIVE AREA

Drug Facts (continued)

- with any other product containing diphenhydramine, even one used on skin (**Nighttime only**)

Ask a doctor before use if you have

- heart disease
- high blood pressure
- liver disease
- thyroid disease
- diabetes
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis (**Nighttime only**)
- glaucoma (**Nighttime only**)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (**Nighttime only**)

When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children (**Nighttime only**)
- marked drowsiness may occur (**Nighttime only**)
- alcohol, sedatives, and tranquilizers may increase drowsiness (**Nighttime only**)

ADHESIVE AREA

Drug Facts (continued)

- use caution when driving a motor vehicle or operating machinery (**Nighttime only**)
- avoid alcoholic beverages (**Nighttime only**)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Do not take DAYTIME and NIGHTTIME products at the same time.

ADHESIVE AREA

Drug Facts (continued)

Directions

- do not take more than directed
- do not take more than 12 capsules in any 24-hour period
- adults and children 12 years and over: take 2 capsules every 4 hours
- children under 12 years: do not use

Other information

- each capsule contains sodium 3 mg (**Daytime only**)

TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients (Daytime only)

corn starch, croscopolone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

ADHESIVE AREA

Drug Facts (continued)

Inactive ingredients (Nighttime only)

corn starch, croscarmellose sodium, croscopolone, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, iron oxide yellow, magnesium stearate, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium bicarbonate, stearic acid, talc, titanium dioxide


Questions or comments? 1-800-426-9391

Walgreens Pharmacist Recommended.
Our pharmacists recommend the Walgreens brand.
We invite you to compare to national brands.

DISTRIBUTED BY: WALGREEN CO.
DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com
©2023 Walgreen Co.

50844
REVU723461568410
R-2201-61569410HN-DF

ITEM 355984
W00000-0000-0



Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-6156
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-6156-10	1 in 1 CARTON; Type 0: Not a Combination Product	06/02/2018	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	2 BLISTER PACK	24
Part 2	2 BLISTER PACK	16

Part 1 of 2

SINUS PRESSURE AND PAIN DAYTIME MAXIMUM STRENGTH

acetaminophen, guaifenesin, phenylephrine hcl tablet, film coated

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
FD&C RED NO. 40 ALUMINUM LAKE (UNII: 6T47AS764T)	
FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	44;615
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/02/2018	

Part 2 of 2

SINUS PRESSURE AND PAIN NIGHTTIME MAXIMUM STRENGTH

acetaminophen, diphenhydramine hcl, phenylephrine hcl tablet, film coated

Product Information

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	12.5 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	

CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
FD&C BLUE NO. 2 ALUMINUM LAKE (UNII: 4AQJ3LG584)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	44;694
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/02/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/02/2018	

Labeler - Walgreen Company (008965063)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(0363-6156)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(0363-6156) , pack(0363-6156)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(0363-6156)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(0363-6156)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(0363-6156)

Revised: 3/2025

Walgreen Company