


CHLORZOXAZONE- chlorzoxazone tablet
PD-Rx Pharmaceuticals, Inc.

Chlorzoxazone Tablets, USP

Rx only

DESCRIPTION

Chlorzoxazone, USP is a centrally-acting skeletal muscle relaxant, available as tablets of 500 mg for oral administration. Its chemical name is 5-Chloro-2-benzoxazolinone, and its structural formula is:



$C_7H_4ClNO_2$

MW 169.57

Chlorzoxazone, USP is a white or practically white, practically odorless, crystalline powder. Chlorzoxazone, USP is slightly soluble in water; sparingly soluble in alcohol, in isopropyl alcohol, and in methanol; soluble in solutions of alkali hydroxides and ammonia.

Chlorzoxazone Tablets, USP contain the inactive ingredients docusate sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose, pregelatinized starch, sodium benzoate, and sodium starch glycolate.

CLINICAL PHARMACOLOGY

Chlorzoxazone is a centrally-acting agent for painful musculoskeletal conditions. Data available from animal experiments as well as human study indicate that chlorzoxazone acts primarily at the level of the spinal cord and subcortical areas of the brain where it inhibits multisynaptic reflex arcs involved in producing and maintaining skeletal muscle spasm of varied etiology. The clinical result is a reduction of the skeletal muscle spasm with relief of pain and increased mobility of the involved muscles. Blood levels of chlorzoxazone can be detected in people during the first 30 minutes and peak levels may be reached, in the majority of the subjects, in about 1 to 2 hours after oral administration of chlorzoxazone. Chlorzoxazone is rapidly metabolized and is excreted in the urine, primarily in a conjugated form as the glucuronide. Less than one percent of a dose of chlorzoxazone is excreted unchanged in the urine in 24 hours.

INDICATIONS

Chlorzoxazone is indicated as an adjunct to rest, physical therapy, and other measures for the relief of discomfort associated with acute, painful musculoskeletal conditions. The mode of action of this drug has not been clearly identified, but may be related to its

sedative properties. Chlorzoxazone does not directly relax tense skeletal muscles in man.

CONTRAINDICATIONS

Chlorzoxazone is contraindicated in patients with known intolerance to the drug.

WARNINGS

Serious (including fatal) hepatocellular toxicity has been reported rarely in patients receiving chlorzoxazone. The mechanism is unknown but appears to be idiosyncratic and unpredictable. Factors predisposing patients to this rare event are not known. Patients should be instructed to report early signs and/or symptoms of hepatotoxicity such as fever, rash, anorexia, nausea, vomiting, fatigue, right upper quadrant pain, dark urine, or jaundice. Chlorzoxazone should be discontinued immediately and a physician consulted if any of these signs or symptoms develop. Chlorzoxazone use should also be discontinued if a patient develops abnormal liver enzymes (e.g., AST, ALT, alkaline phosphatase and bilirubin).

The concomitant use of alcohol or other central nervous system depressants may have an additive effect.

Usage in Pregnancy

The safe use of chlorzoxazone has not been established with respect to the possible adverse effects upon fetal development. Therefore, it should be used in women of childbearing potential only when, in the judgment of the physician, the potential benefits outweigh the possible risks.

PRECAUTIONS

Chlorzoxazone should be used with caution in patients with known allergies or with a history of allergic reactions to drugs. If a sensitivity reaction occurs such as urticaria, redness, or itching of the skin, the drug should be stopped.

If any symptoms suggestive of liver dysfunction are observed, the drug should be discontinued.

ADVERSE REACTIONS

Chlorzoxazone containing products are usually well tolerated. It is possible in rare instances that chlorzoxazone may have been associated with gastrointestinal bleeding. Drowsiness, dizziness, lightheadedness, malaise, or over-stimulation may be noted by an occasional patient. Rarely, allergic-type skin rashes, petechiae, or ecchymoses may develop during treatment. Angioneurotic edema or anaphylactic reactions are extremely rare. There is no evidence that the drug will cause renal damage. Rarely, a patient may note discoloration of the urine resulting from a phenolic metabolite of chlorzoxazone. This finding is of no known clinical significance.

To report SUSPECTED ADVERSE REACTIONS, contact Teva at 1-888-838-2872

or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

OVERDOSAGE

Symptoms

Initially, gastrointestinal disturbances such as nausea, vomiting, or diarrhea together with drowsiness, dizziness, lightheadedness or headache may occur. Early in the course there may be malaise or sluggishness followed by marked loss of muscle tone, making voluntary movement impossible. The deep tendon reflexes may be decreased or absent. The sensorium remains intact, and there is no peripheral loss of sensation. Respiratory depression may occur with rapid, irregular respiration and intercostal and substernal retraction. The blood pressure is lowered, but shock has not been observed.

Treatment

Gastric lavage or induction of emesis should be carried out, followed by administration of activated charcoal. Thereafter, treatment is entirely supportive. If respirations are depressed, oxygen and artificial respiration should be employed and a patent airway assured by use of an oropharyngeal airway or endotracheal tube. Hypotension may be counteracted by use of dextran, plasma, concentrated albumin or a vasopressor agent such as norepinephrine. Cholinergic drugs or analeptic drugs are of no value and should not be used.

DOSAGE AND ADMINISTRATION

Usual Adult Dosage

One tablet three or four times daily. If adequate response is not obtained with this dose, it may be increased to one and one-half tablets (750 mg) three or four times daily. As improvement occurs dosage can usually be reduced.

HOW SUPPLIED

Chlorzoxazone Tablets USP, 500 mg are available as oblong, scored, white tablets debossed with "WPI" on one side and "39|68" on the other side and are packaged in bottles of:

(NDC 55289-633-10) Bottle of 10

(NDC 55289-633-20) Bottle of 20

(NDC 55289-633-24) Bottle of 24

(NDC 55289-633-28) Bottle of 28

(NDC 55289-633-30) Bottle of 30

(NDC 55289-633-40) Bottle of 40

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Dispense contents with a child-resistant closure (as required) and in a tight container as

defined in the USP/NF.

Keep this and all medications out of the reach of children.

PRINCIPAL DISPLAY PANEL

Chlorzoxazone Tablets, USP
500 mg
Rx only

May Cause Drowsiness. Use Caution While Driving
Or Performing Tasks Requiring Mental Alertness.

R only WARNING: KEEP THIS OUT OF THE REACH OF CHILDREN
DOSAGE and STORAGE: SEE PACKAGE INSERT

55289-633-28	55289-633-28	55289-633-28
CHLORZOXAZONE	CHLORZOXAZONE	CHLORZOXAZONE
USP	USP	USP
500 mg	500 mg	500 mg
28 TABLETS	28 TABLETS	28 TABLETS
ReOrder # 100821	ReOrder # 100821	ReOrder # 100821
LOT D23B93	LOT D23B93	LOT D23B93
EXP 10/2024	EXP 10/2024	EXP 10/2024

CALL YOUR DOCTOR FOR MEDICAL ADVICE ABOUT SIDE EFFECTS.
YOU MAY REPORT SIDE EFFECTS TO THE FDA AT 1-800-FDA-1088

TAKE _____ TABLET(S) _____ TIMES A DAY.
TOME _____ TABLETA(S) _____ VECES AL DIA.
Each TABLET Contains: CHLORZOXAZONE USP 500 MG

WHITE

NDC: 55289-633-28



CHLORZOXAZONE
USP

500 mg

28 TABLETS

ORGANOLEPTIC MARKINGS:
WPP: 39 08

OBLONG SCORED



GTIN: 00355289633280
SNO: D23B93000001
EXP: 10/2024
LOT: D23B93

355289633280
00591252001
WATSON PHARMA PRIVATE LTD
VERNA, SALLETTI GOA, INDIA
403 722

CHLORZOXAZONE

chlorzoxazone tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:55289-633(NDC:0591- 2520)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORZOXAZONE (UNII: H0DE420U8G) (CHLORZOXAZONE - UNII:H0DE420U8G)	CHLORZOXAZONE	500 mg

Inactive Ingredients

Ingredient Name	Strength

DOCUSATE SODIUM (UNII: F05Q2T2JA0)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

Product Characteristics

Color	white	Score	2 pieces
Shape	OVAL (oblong)	Size	17mm
Flavor		Imprint Code	WPI;39;68
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55289-633-10	10 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/31/2011	
2	NDC:55289-633-20	20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/31/2011	
3	NDC:55289-633-24	24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/31/2011	
4	NDC:55289-633-28	28 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/31/2011	
5	NDC:55289-633-30	30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/31/2011	
6	NDC:55289-633-40	40 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/31/2011	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA089859	08/22/2011	

Labeler - PD-Rx Pharmaceuticals, Inc. (156893695)

Registrant - PD-Rx Pharmaceuticals, Inc. (156893695)

Establishment

Name	Address	ID/FEI	Business Operations
PD-Rx Pharmaceuticals, Inc.		156893695	repack(55289-633)