

## **RAPIDOX ACTIVATOR- lactic acid solution, concentrate**

### **DeLaval**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click [here](#).*

-----

### **RapidOx™ Activator**

#### **Concentrated Disinfecting Teat Dip**

Aids in reducing the spread of mastitis-causing organisms

#### **ACTIVE INGREDIENTS:**

Lactic Acid ..... 12.5%

#### **KEEP OUT OF REACH OF CHILDREN**

#### **! DANGER**

READ AND UNDERSTAND ALL INSTRUCTIONS BEFORE USING THIS PRODUCT

**RapidOx** is a three-part Chlorine Dioxide teat dip that must be used with the InstaMix G Emollient Concentrate. Recommended for pre/post milking applications and can be used to reduce the spread of mastitis causing organisms.

#### **HAZARD STATEMENTS**

Causes severe skin burns and eye damage. Do not breathe dust/fume/gas/mist/vapors/spray. Wash face, hands and any exposed skin thoroughly after handling. Wear protective gloves/protective clothing/eye protection/face protection.

#### **UN3265**

**Corrosive liquid, acidic, organic, n.o.s.(Lactic Acid)**

#### **FIRST AID**

**Immediately call a POISON CENTER or doctor/physician**

**If in Eyes:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or doctor/physician.

**If on Skin:** Remove/Take off immediately all contaminated clothing. Rinse skin with water/shower. Wash contaminated clothing before reuse.

**If Inhaled:** Remove victim to fresh air and keep at rest in a position comfortable for breathing. Immediately call a POISON CENTER or doctor/physician.

**If Swallowed:** Rinse mouth. Do NOT induce vomiting.

#### **STORAGE AND DISPOSAL**

**THOROUGHLY RINSE ALL MEASURING DEVICES AND STORE OUT OF REACH OF CHILDREN.** Avoid contamination of food or feed. Avoid freezing. Keep out of direct sunlight. Store in suitable labelled containers. Storage temperatures: 32°F to 86°F. Store locked up. Do not mix with bleach or other chlorinated products - will cause

chlorine gas. Dispose of contents/container to an approved waste disposal plant.

## **CHEMTREC EMERGENCY NUMBER 1-800-424-9300**

### **Not for Human Use**

### **Additional Information on Safety Data Sheet**

#### **DIRECTIONS**

IMPORTANT THIS IS A THREE-PART CONCENTRATE - MUST BE MIXED PRIOR TO USE WITH INSTAMIX™ G EMOLLIENT CONCENTRATE.

CREATES CHLORINE DIOXIDE See Dilution Chart for mix ratios.

#### General Use

- Concentrated formula must be diluted into a ready-to-use product.
- Using a manual pump or automatic system, mix into a clean teat cup or container at least 15 minutes prior to use. Prepare only the quantity needed for one milking.
- If cow's teats are sore or chapped, consult a veterinarian before starting to use teat dip.
- If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.
- If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

#### Pre-Dipping Use

- Perform forestripping and inspect teats for clinical mastitis.
- Pre-wash teats thoroughly just prior to milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels.
- Just before each cow is milked, dip, or spray each teat full-length into the teat dip cup containing product.
- Allow product to stand for 30 seconds.
- Wipe teats dry after dipping, using single-service towels to avoid contamination of milk

#### Post-Dipping Use

- Immediately after each milking, use product.
- Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe.
- If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Undiluted product may undergo an increase in viscosity when exposed to cold temperatures. Pumps must be recalibrated in those conditions to maintain accurate dosing. If product freezes, thaw it completely and mix properly before use. Consult your DeLaval representative for questions on specific applications.

#### **DILUTION CHART**

% Emollient (gal/100)	RapidOx Activator (gal/100)	RapidOx Base (gal/100)	Instamix™ G Emollient Conc (gal/100)	Water (gal/100)
--------------------------	--------------------------------	---------------------------	---	--------------------

1%	2	2	2	94
2%	2	2	4	92
3%	2	2	5	91
4%	2	2	7	89
5%	2	2	9	87
6%	2	2	11	85
7%	2	2	12	84
8%	2	2	14	82
9%	2	2	16	80
10%	2	2	18	78
20%	2	2	35	60

Manufactured for: **DeLaval Inc.**  
 11100 N. Congress Avenue  
 Kansas City, MO 6153  
 Phone: 816-891-7700  
 Manufactured by: West Agro, Inc.  
 © 2025, Made in UA



4204-A1125



CIO<sub>2</sub>

Activator

RapidOx™

Activator

Concentrated Disinfecting Teat Dip

Aids in reducing the spread of mastitis-causing organisms

Sellador desinfectante concentrado

Ayuda a reducir la transmisión de microorganismos que causan mastitis

ACTIVE INGREDIENTS:

Lactic Acid ..... 12.5%




KEEP OUT OF REACH OF CHILDREN

 **DANGER**

MANTENER FUERA DEL ALCANCE DE LOS NIÑOS

 **PELIGRO**

4204

NDC NUMBER: 55756-204

READ AND UNDERSTAND ALL INSTRUCTIONS BEFORE USING THIS PRODUCT

SI USTED NO PUEDE LEER ESTA ETIQUETA, POR FAVOR PIDA A ALGUIEN MAS QUE SE LA LEA ANTES DE USAR ESTE PRODUCTO

RapidOx is a three-part Chlorine Dioxide test dip that must be used with the InstaMix G Enrichment Concentrate. Recommended for prepost milking applications and can be used to reduce the spread of mastitis causing organisms.



Causes severe skin burns and eye damage. Do not breathe dust/fume/gas/mist/vapors/spray. Wash face, hands and any exposed skin thoroughly after handling. Wear protective gloves/protective clothing/eye protection/face protection.

Immediately call a POISON CENTER or doctor/physician

If in Eyes:

Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or doctor/physician.

If on Skin:

Remove/Take off immediately all contaminated clothing. Rinse skin with water/shower. Wash contaminated clothing before reuse.

If Inhaled:

Remove victim to fresh air and keep at rest in a position comfortable for breathing. Immediately call a POISON CENTER or doctor/physician.

If Swallowed:

Rinse mouth. Do NOT induce vomiting.

FIRST AID

THOROUGHLY RINSE ALL MEASURING DEVICES AND STORE OUT OF REACH OF CHILDREN. Avoid contamination of food or feed. Avoid freezing. Keep out of direct sunlight. Store in suitable labelled containers. Storage temperatures: 52°F to 86°F. Store locked up. Do not mix with bleach or other chlorinated products – will cause chlorine gas. Dispose of contents/container to an approved waste disposal plant.

CHEMTREC EMERGENCY NUMBER 1-800-424-9300

Not For Human Use

Additional Information on Safety Data Sheet

DILUTION CHART

% Enrichment	RapidOx Activator (gal/100 gal)	RapidOx Base (gal/100 gal)	InstaMix G Enrichment Conc. (gal/100)	Water (gal/100)
1%	2	2	2	94
2%	2	2	4	92
3%	2	2	5	91
4%	2	2	7	89
5%	2	2	9	87
6%	2	2	11	85
7%	2	2	12	84
8%	2	2	14	82
9%	2	2	15	80
10%	2	2	16	78
20%	2	2	35	60

DIRECTIONS

IMPORTANT: THIS IS A THREE-PART CONCENTRATE – MUST BE MIXED PRIOR TO USE WITH INSTAMIX™ G ENRICHMENT CONCENTRATE. CREATES CHLORINE DIOXIDE. See Dilution Chart for mix ratios.

General Use

- Concentrated formula must be diluted into a ready-to-use product.
- Using a manual pump or automatic system, mix into a clean teat cup or container at least 15 minutes prior to use. Prepare only the quantity needed for one milking.
- If cow's teats are sore or chapped, consult a veterinarian before starting to use test dip.
- If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.
- If product in dip cup becomes sticky/dry, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Teat-Dipping Use

- Perform teatdipping and inspect teats for clinical mastitis.
- Pre-wash teats thoroughly just prior to milking with appropriate udder wash solution or pre-milking test dip. Teats should then be dried with single-service towels.
- Just before each cow is milked, dip, or spray each teat full-length into the test dip cup containing product.
- Allow product to stand for 30 seconds.
- Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post-Dipping Use

- Immediately after each milking, use product:
- Dip each teat full-length into the test dip cup containing product. Allow to air dry. Do not wipe.
- If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Undiluted product may undergo an increase in viscosity when exposed to cold temperatures. Pump must be recalibrated in those conditions to maintain accurate dosing. If product freezes, thaw it completely and mix properly before use. Consult your DeLaval representative for questions on specific applications.

Manufactured for: DeLaval Inc.

11100 N. Congress avenue

Kansas City, MO 64153

Phone: 816-891-7700

Manufactured by: West Agro, Inc.

© 2025, Made in UA

NET CONTENTS: 55 US GAL. (208.2 L)

LOT NO.:

EXP. DATE:

UN3265

Corrosive liquid, acidic, organic, non-flammable

**RAPIDOX ACTIVATOR**  
 lactic acid solution, concentrate

Product Information				
Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:55756-204	
Route of Administration	TOPICAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
LACTIC ACID (UNII: 33X04XA5AT) (LACTIC ACID - UNII:33X04XA5AT)		LACTIC ACID	125 g in 1 L	
Inactive Ingredients				
Ingredient Name			Strength	
C9-11 PARETH-8 (UNII: 80E6PSE1XL)				
WATER (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55756-204-01	208 L in 1 DRUM		
2	NDC:55756-204-27	1041 L in 1 TANK		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
unapproved drug other			12/19/2025	

**Labeler -** DeLaval (173704867)

Establishment			
Name	Address	ID/FEI	Business Operations
West Agro, Inc.		147528723	api manufacture, manufacture, pack, label, analysis

Establishment			
Name	Address	ID/FEI	Business Operations
West Agro, Inc.		095906480	api manufacture, manufacture, pack, label, analysis