

**IBUPROFEN PM- ibuprofen, diphenhydramine hcl capsule, liquid filled
Dolgenercorp, Inc. (DOLLAR GENERAL & REXALL)**

Drug Facts

Active ingredient (in each capsule)

Diphenhydramine hydrochloride 25 mg

**Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*
(present as the free acid and potassium salt)**

***non-steroidal anti-inflammatory drug**

Purpose

Nighttime sleep-aid

Pain Reliever

Uses

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery
- unless you have time for a full night's sleep
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have glaucoma
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a breathing problem such as emphysema or chronic bronchitis
- you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers, or any other sleep-aid
- taking any other drug
- taking any other antihistamines
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any continuing medical illness

When using this product

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problem or stroke:
 - chest pain

- trouble breathing
- leg swelling
- slurred speech
- weakness in one part or side of body
- pain gets worse or last more than 10 days
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222)

Directions

- **do not take more than directed**
- adults and children 12 years and over:
 - take 2 capsules at bedtime
 - do not take more than 2 capsules in 24 hours

Other information

- **each capsule contains:** potassium 20 mg
- read all warnings and directions before use. Keep carton.
- store at 20° to 25°C (68° to 77°F)
- avoid high humidity and excessive heat above 40°C (104°F)
- protect from light
- swallow whole; do not crush, chew, or dissolve

Inactive ingredients

anidrisorb, FD&C blue #1, D&C red #33, gelatin, hydrolyzed gelatin, medium chain triglycerides, opacode white ink, polyethylene glycol 600, potassium hydroxide, purified water. Ingredients of opacode white ink, shellac glaze in ethanol, titanium dioxide, n-butyl alcohol, lecithin (soya), simethicone, purified water

Questions or comments?

Call **1-888-309-9030**

Principal Display Panel

Compare to the active ingredients of Advil® PM**

Ibuprofen PM Capsules

Ibuprofen 200 mg / Pain reliever (NSAID)

Diphenhydramine HCl 25 mg / Nighttime Sleep-Aid

SOFTGELS**

(**liquid filled capsules)

**This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® PM.

TAMPER-EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY DOLGENCORP, LLC

100 MISSION RIDGE, GOODLETTSVILLE, TN 37072

Product Packaging

IBUPROFEN PM

ibuprofen, diphenhydramine hcl capsule, liquid filled

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-856	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		IBUPROFEN	200 mg	
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)		DIPHENHYDRAMINE HYDROCHLORIDE	25 mg	
Inactive Ingredients				
Ingredient Name			Strength	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
D&C RED NO. 33 (UNII: 9DBA0SBB0L)				
GELATIN (UNII: 2G86QN327L)				
POLYETHYLENE GLYCOL 600 (UNII: NL4J9F21N9)				
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)				
WATER (UNII: 059QF0KO0R)				
SORBITAN (UNII: 6O92ICV9RU)				
SORBITOL (UNII: 506T60A25R)				
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)				
Product Characteristics				
Color	purple (purple bluish)	Score	no score	
Shape	OVAL	Size	16mm	
Flavor		Imprint Code	1007	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-856-40	1 in 1 BOX	04/30/2018	01/31/2025
1		40 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:55910-856-20	1 in 1 BOX	04/30/2018	01/31/2025
2		20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA200888	04/30/2018	01/31/2025

Labeler - Dolgencorp, Inc. (DOLLAR GENERAL & REXALL) (068331990)

Revised: 10/2022

Dolgencorp, Inc. (DOLLAR GENERAL & REXALL)