

**4267 FIRST AID KIT- 4267 first aid kit  
Honeywell Safety Products USA, INC**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----  
**0498-4267: First Aid Kit (Neomycin, EW, alcohol wipes, Burn Sray, Antiseptic Spray, Hand Sanitizer, aypanal- FAK3SHLF-CLSB)**

**Eyesaline  
Active ingredient**

Sterile Water 99%

**Eyesaline  
Purpose**

Eyewash

**Eyesaline  
Uses**

- for flushing the eye to remove loose foreign material, air pollutants or chlorinated water

**Eyesaline  
Warnings**

**For external use only-**

Obtain immediate medical treatment for all open wounds in or near eyes.

To avoid contamination, do not touch tip of container to any surface.

Do not reuse. Once opened, discard.

**Do not use**

- if solution changes color or becomes cloudy
- if you have open wounds in or near the eyes, get medical help right away.

**Stop use and ask a doctor if**

- you experience eye pain
- changes in vision
- continued redness or irritation of the eye
- condition worsens or persists

**Keep out of reach of children**

If swallowed, get medical help or contact a Poison Control Center right away.

## **Eyesaline**

### ***Directions***

- remove contacts before using
- twist top to remove
- flush the affected area as needed
- control rate of flow by pressure on the bottle
- if necessary, continue flushing with emergency eyewash or shower

## **Eyesaline**

### ***Inactive ingredients***

sodium chloride, sodium phosphate dibasic, sodium phosphate monobasic

## **Eyesaline**

### ***Questions***

1-800-430-5490 Honeywell Safety Products USA, Inc. Smithfield, RI 02917

## **Alcohol Wipe**

### ***Active ingredient***

Isopropyl alcohol 70%

## **Alcohol Wipe**

### ***Purpose***

First aid antiseptic

## **Alcohol Wipe**

### ***Uses***

- first aid to help prevent infection in minor cuts, scrapes, and burns

## **Alcohol Wipe**

### ***Warnings***

**For external use only**

### **Do not use**

- in the eyes
- over large areas of the body

**Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burn

**When using this product**

- do not use longer than one week unless directed by a doctor

**Stop use and consult a doctor**

- if condition persists or gets worse

**Keep out of reach of children**

- If swallowed, get medical help or contact a Poison Control Center right away.

**Alcohol Wipe*****Directions***

- clean the affected area
- apply wipe to affected area 1 to 3 times daily
- may be covered with a sterile bandage
- discard wipe after single use

**Alcohol Wipe*****Other information***

store at room temperature 15 ° to 25 ° C (59 ° to 77 °F)

**Alcohol Wipe*****Inactive ingredient***

water

**Alcohol Wipe*****Questions***

1-800-430-5490

**Aypanal*****Active ingredient***

Acetaminophen 325 mg

**Aypanal*****Purpose***

Pain reliever/ fever reducer

## **Aypanaly**

### **Uses**

- temporarily relieves minor aches and pains due to the common cold and headache - temporarily reduces fever

### **Keep out of reach of children.**

Keep out of reach of children.

## **Aypanal**

### **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product:

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- If a skin rash occurs, stop use and seek medical help right away.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription).
- If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

### **Ask a doctor before use if you have**

- liver disease

### **Ask a doctor or pharmacist before use if**

- you are taking the blood thinning drug warfarin

### **Stop using and ask a doctor if**

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

### **If pregnant or breast-feeding**

If pregnant or breast-feeding, ask a health professional before use.

### **Overdose warning**

- In case of accidental overdose, get medical help or contact a Poison Control Center right away.
- Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Aypanal**

#### ***Directions***

**do not take more than directed (see overdose warning)**

#### **adults and children 12 years of age or older**

- take two tablets every 4-6 hours while symptoms last
- do not take more than 12 tablets in 24 hours

#### **children 6 to under 12 years of age**

- take 1 tablet every 4-6 hours while symptoms last
- do not take more than 5 tablets in 24 hours

**children under 6 years consult a doctor**

### **Aypanal**

#### ***Other information***

- store at room temperature 15<sup>0</sup> to 30<sup>0</sup> C (59<sup>0</sup> - 86<sup>0</sup> F)
- TAMPER EVIDENT PACKETS- DO NOT USE IF OPEN OR TORN

### **Aypanal**

#### ***Inactive ingredients***

corn starch, microcrystalline cellulose, povidone, sodium starch glycolate, stearic acid

### **Aypanal**

#### ***Questions***

1-800-430-5490

### **Neomycin**

#### ***Active ingredient***

Neomycin sulfate (5 mg equivalent to 3.5 mg Neomycin base)

### **Neomycin**

#### ***Purpose***

First aid antibiotic

## **Neomycin**

### ***Uses***

- first aid to help prevent infection in - minor cuts - scrapes - burns

### **Do not use**

- in the eyes
- over large areas of the body

## **Neomycin**

### ***Warnings***

### **For external use only**

### **Do not use**

- in the eyes
- over large areas of the body

### **Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burns

### **Stop use and ask a doctor if**

- a rash or other allergic reaction develops
- you need to use longer than 1 week

### **Stop use and ask a doctor if**

- the condition persists or gets worse
- a rash or other allergic reaction develops
- you need to use longer than 1 week

### **Keep out of reach of children**

- If swallowed, get medical help or contact a Poison Control Center right away

## **Neomycin**

### ***Direction***

- clean the affected area
- apply a small amount of the product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage

## **Neomycin**

## ***Other information***

store at 15 ° to 25 ° C (59 ° to 77 ° F)

## **Neomycin**

### ***Inactive ingredient***

petrolatum

## **Neomycin**

### ***Questions?***

1-800-430-5490

## **Antiseptic Spray**

### ***Active ingredient***

Benzalkonium chloride 0.13%

## **Antiseptic Spray**

### ***Purpose***

First aid antiseptic

## **Antiseptic Spray**

### ***Uses***

- first aid to help prevent infection in minor cuts, scrapes and burns

## **Antiseptic Spray**

### ***Warnings***

#### **For external use only**

#### **Do not use**

- in or near the eyes
- over large areas of the body

#### **Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burns

#### **When using this product**

- do not use longer than one week unless directed by a doctor

#### **Stop use and ask a doctor if**

- the condition persists or gets worse

### **Keep out of reach of children**

- If swallowed, get medical help or contact a Poison Control Center right away

### **Antiseptic Spray**

#### ***Directions***

- clean the affected area
- spray a small amount of this product on the area 1 to 3 times daily
- may be covered with a sterile bandage
- if bandaged, let dry first

### **Antiseptic Spray**

#### ***Other information***

- shake well
- store at room temperature 15<sup>0</sup>-30<sup>0</sup> C (59<sup>0</sup> -86<sup>0</sup> F)

### **Antiseptic Spray**

#### ***Inactive ingredients***

diazolidinyl urea, edetate disodium, glycerin, hypromellose, methylparaben, octoxynol 9, propylene glycol, propylparaben, trolamine, water

### **Antiseptic Spray**

#### ***Questions***

1-800-430-5490

### **Burn Spray**

#### ***Active ingredient***

Benzethonium chloride 0.2% w/w

Benzocaine 10% w/w

Menthol 0.33% w/w

### **Burn Spray**

#### ***Purpose***

Benzethonium chloride 0.2% w/w

Benzocaine 10% w/w

Menthol 0.33% w/w

### **Burn Spray**



## **Uses**

for the temporary relief of pain and itching and helps protect against infection in:

- minor cuts and scrapes
- burns
- sunburn
- insect bites
- minor skin irritations

## **Burn Spray**

### ***Warnings***

#### **For external use only**

#### **Flammable**

- keep away from fire or flame
- contents under pressure
- do not puncture or incinerate container
- do not expose to temperatures above 120 0 F

#### **Do not use**

- in or near the eyes or other mucous membranes
- in case of serious burns
- in case of deep or puncture wounds
- for prolonged period of time
- on large portion of the body

#### **Stop use and ask a doctor if**

- condition worsens or symptoms persist for more than 7 days
- condition clears up and recurs within a few days
- redness, swelling, or irritation occurs

#### **Keep out of the reach of children**

- If swallowed, get medical help or contact a Poison Control Center right away.

## **Burn Spray**

### ***Directions***

- clean the affected area
- shake can well before using
- hold 4 - 6 inches from surface and spray area until wet
- may be covered with a sterile bandage, if bandaged let dry first
- for adult institutional use only
- not intended for use on children

## **Burn Spray**

### ***Other information***

- avoid inhaling

- use only as directed
- intentional misuse by deliberately concentrating or inhaling the contents may be harmful or fatal

## **Burn Spray**

### ***Inactive ingredients***

dipropylene glycol, isobutane, n-butane, propane

## **Hand Sanitizer**

### ***Active ingredient***

Ethyl alcohol 62%

## **Hand Sanitizer**

### ***Purpose***

Antiseptic handwash

## **Hand Sanitizer**

### ***Uses***

- for hand washing to decrease bacteria on skin
- recommended for repeated use

## **Hand Sanitizer**

### ***Warnings***

**For external use only**

**Flammable, keep away from fire or flame**

### **When using this product**

- do not use in the eyes
- discontinue use if irritation and redness develops. If condition persists for more than 72 hours consult a doctor.

### **Keep out of reach of children**

If swallowed, get medical help or contact a Poison Control Center right away.

## **Hand Sanitizer**

### ***Directions***

- wet hands thoroughly with product and allow to dry without wiping

## **Hand Sanitizer**

### ***Other information***

store at 15 ° to 25 ° C (59 ° to 77 ° F)

## **Hand Sanitizer**

### ***Inactive ingredients***

acrylates/C10-30 alkyl acrylate crosspolymer, aloe barbadensis leaf juice, dl-alpha tocopheryl acetate, fragrance, PEG-60 almond glycerides, propylene glycol, purified water, triisopropanolamine

## **Hand Sanitizer**

### ***Questions or Comments?***

1-800-275-3433 info@waterjel.com www.waterjel.com

## **Ammonia**

### ***Active ingredient***

Ammonia 15%

## **Ammonia**

### ***Purpose***

Respiratory stimulant

## **Ammonia**

### ***Uses***

- to prevent or treat fainting

## **Ammonia**

### ***Warnings***

#### **For external use only**

#### **Do not use**

- if you have breathing problems such as asthma or emphysema

#### **Stop use and ask a doctor if**

- condition persists

#### **Keep out of reach of children**

If swallowed get medical help or contact a Poison Control Center right away.

## **Ammonia**

### ***Directions***

- hold inhalant away from face and crush ampoule between thumb and forefinger at

position indicated on sleeve.

- hold near nostrils for inhalation of volatile vapor

### **Ammonia**

#### ***Other information***

- store at room temperature away from light

### **Ammonia**

#### ***Inactive ingredients***

alcohol USP, FD&C red #40, lavender oil, lemon oil fcc, nutmeg oil, purified water

### **Ammonia**

#### ***Questions or Comments?***

1-800-430-5490

### **Eyesaline**

#### **Principal Display Panel**

Honeywell

TAMPER-EVIDENT CAP.  
TAPA CON SELLO DE SEGURIDAD.  
BOUCHON INDICATEUR D'INFRACTION.

eyesaline®

LAVAOJOS  
EYESALINE

EYESALINE  
EYEWASH

LAVAGE  
OCULAIRE  
EYESALINE

Solución  
Isotónico Estéril

Sterile  
Isotonic Solution

La Solution  
Isotonique Stérile

16 fl. oz. (473 mL)

NPN: 80057528  
64809 1 45033 3

Drug Facts (for USA only)

**Active ingredient** Sterile water 99%  
**Purpose** Eyewash  
**Uses** for flushing the eye to remove loose foreign material, air pollutants, or chlorinated water.  
**Warnings**  
**For external use only** - Obtain immediate medical treatment for all open wounds in or near the eyes. To avoid contamination, do not touch tip of container to any surface. Do not reuse. Once opened, discard.  
**Do not use**  
• if solution changes color or becomes cloudy  
• if you have open wounds in or near the eyes, get medical help right away  
**Stop use and consult a doctor if:**  
• you experience eye pain • changes in vision  
• continued redness or irritation of the eye  
• condition worsens or persists  
**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.  
**Directions**  
• remove contacts before using • twist top to remove  
• flush the affected area as needed  
• control rate of flow by pressure on the bottle  
• if necessary, continue flushing with emergency eyewash or shower  
**Inactive ingredients** sodium chloride, sodium phosphate dibasic, sodium phosphate monobasic  
**Questions?** Call 1-800-430-5490  
Honeywell Safety Products USA, Inc. Smithfield, RI. 02917

LABEL #32-0045/0 Rev. J REORDER / NUEVO PEDIDO / REAPPROVISIONNEMENT #32-000454-0000

space for lot code and supplier part number

PEEL / PELAR / PELEL

Datos de medicamento (Para EE.UU. solamente)


**Ingrediente Activo** Agua estéril 99%  
**Propósito** Lavaojos  
**Usos** para el lavado de ojo para quitar las partículas sueltas y extrañas, los contaminantes aeros, o agua de cloruro  
**Advertencias**  
**Para el uso externo sólo** - Obtenga tratamiento médico inmediato para todas las heridas abiertas en o cerca de los ojos. Para evitar la contaminación, no toque la punta del envase con ninguna superficie. No vuelva a usar. Vez abierto, descarte.  
**No se use** • si la solución se enturbia o cambia de color  
• si tiene heridas abiertas en o cerca del ojo, obtenga ayuda medica de inmediato  
**Deje de usar y consulte a un médico si:**  
• experimenta dolor de ojo • cambio de visión  
• rojez continuo o irritación del ojo  
• la condición empeora o persiste  
**Manténgase fuera del alcance de los niños.**  
En caso de ingestión accidental, obtenga atención médica o llame a un Centro de control de envenenamiento inmediatamente.  
**Instrucciones**  
• quite los lentes de contacto antes de usar la solución  
• tuerza la tapa para quitar  
• enjuague el área afectada según se necesite  
• controle el chorro haciendo presión el la botella  
• si es necesario, sigue enjuagado con un lavaojos o ducha de emergencia  
**Ingredientes inactivos** cloruro de sodio, fosfato de sodio dibásico, fosfato de sodio monobásico  
**¿Preguntas?** Llame al 1-800-430-5490  
Honeywell Safety Products USA, Inc. Smithfield, RI. 02917

Information


**Usages** Pour le rinçage des yeux afin d'enlever un corps étranger, des polluants atmosphériques ou de l'eau chlorée.  
**Advertissements**  
**Pour usage externe seulement** - Obtenir immédiatement des soins médicaux pour toutes les plaies ouvertes dans ou près des yeux. Pour éviter toute contamination, ne pas toucher la pointe du récipient à n'importe quelle surface. Ne pas réutiliser. Une fois ouvert, jeter.  
**Ne pas utiliser**  
• si la solution a changé de couleur ou si elle est brouillée  
• si vous avez des plaies ouvertes aux yeux ou à proximité, consultez immédiatement un médecin  
**Cesser d'utiliser la solution et consulter un médecin**  
• vous ressentez une douleur oculaire • si votre vision change  
• rougeur ou irritation persistante des yeux  
• condition empire ou persiste  
**Garder hors de la portée des enfants.**  
En cas d'ingestion, communiquer immédiatement avec un médecin ou avec un centre antipoison.  
**Mode d'emploi**  
• enlever les verres de contact avant l'utilisation • dévisser le bouchon pour l'enlever • rincer la zone touchée selon les besoins • ajuster le débit d'écoulement de la solution en augmentant ou en réduisant la pression exercée sur le contenant  
• si nécessaire, continuer de rincer avec une solution de rinçage oculaire d'urgence ou une douche  
**Ingédients** eau stérile, chlorure de sodium, phosphate dibasique de sodium, phosphate monobasique de sodium  
**Des questions?** Faites le 1-800-430-5490  
Honeywell Safety Products USA, Inc. Smithfield, RI. 02917

Alcohol Wipe  
Principal Display Panel

825362T Rev. A  
 white printed  
 Pms 072 blue, Pms 186 red,  
 Pms 7459 blue and black





**Aypanal  
 Principal Display Panel**



**Neomycin**  
**Principal Display Panel**

796041-25 Rev A Unit Carton Printing Plate for "C" size carton.



# Antiseptic Spray

## Principal Display Panel

	032203		<b>Drug Facts</b>	
	<b>ANTISEPTIC SPRAY</b> <i>Benzalkonium chloride</i> <b>First Aid Antiseptic</b>		<b>Active ingredient</b> Benzalkonium chloride 0.13% ..... First aid antiseptic	<b>Purpose</b> First aid antiseptic
Net contents 2 fl oz (59 mL) ANSI Z308.1-2003			<b>Uses</b> first aid to help prevent infection in minor cuts, scrapes and burns	
<b>Warnings</b> For external use only			<b>Do not use</b> • in or near the eyes • over large areas of the body	
<b>Directions</b> • clean the affected area • spray a small amount of this product on the area 1 to 3 times daily • may be covered with a sterile bandage • if bandaged, let dry first			<b>Ask a doctor before use if you have</b> • deep or puncture wounds • animal bites • serious burns	
<b>Other information</b> • shake well • store at room temperature, 15° to 30°C (59° to 86°F)			<b>When using this product</b> do not use longer than one week unless directed by a doctor	
<b>Inactive ingredients</b> clazolidinyl urea, edetate disodium, glycerin, hypromellose, methylparaben, octoxynol 9, propylene glycol, propylparaben, triethylamine, water			<b>Stop use and ask a doctor if</b> condition persists or gets worse	
<b>Questions or comments?</b> 1-800-430-5490			<b>Keep out of reach of children.</b> If swallowed get medical help or contact a Poison Control Center right away.	
<small>Mfg. for Honeywell Safety Products USA, Inc. Smithfield, RI 02917</small>			<small>002203 Rev. G</small>	

# Burn Spray

## Principal Display Panel



SHAKE WELL BEFORE USING

# Honeywell BURN SPRAY

Water soluble  
**Benzethonium chloride**  
Topical antiseptic  
**Benzocaine**  
Topical anesthetic  
**Menthol**  
Topical anesthetic

Provides antiseptic treatment  
and helps relieve the pain of minor burns  
and sunburn.

**CAUTION: FLAMMABLE**  
Contents under pressure  
Read warning on back panel.

NET WT. 3 OZ (85gm.)

47.0306

Cat. No. 201005

### DRUG FACTS

#### Active ingredients

Benzethonium chloride 0.2% w/w	Topical antiseptic
Benzocaine 1.0% w/w	Topical anesthetic
Menthol 3.3%	Topical anesthetic

#### Purpose

**Uses** • for the temporary relief of pain and itching and helps to protect against infection in  
• minor cuts and scrapes • burns • sunburn • insect bites • minor skin irritations

#### Warnings

##### For external use only

**Flammable** • keep away from fire or flame • contents under pressure  
• do not puncture or incinerate container • do not expose to temperatures above 120°F

**Do not use** • in or near eyes or other mucous membranes • in case of serious burns  
• in case of deep or puncture wounds • for a prolonged period of time  
• on large portion of the body

##### Stop use and ask a doctor if:

• conditions worsens or symptoms persist for more than 7 days  
• condition clears up and recurs within a few days  
• redness, swelling or irritation occurs

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

#### Directions

• clean the affected area • shake can well before using  
• hold 4-6 inches from surface and spray area until wet  
• may be covered with a sterile bandage. If bandaged, let dry first  
• for adult institutional use only • not intended for use on children

#### Other information

• avoid inhaling • use only as directed  
• intentional misuse by deliberately concentrating and inhaling the contents may be harmful or fatal

**Inactive ingredients** dipropylene glycol, isobutane, n-butane, propane

**Questions or comments?** 1-800-430-5490



Distributed by  
Honeywell Safety  
Products USA, Inc.  
Smithfield, RI 02917

**Honeywell**

## Hand Sanitizer Principal Display Panel



**INSTANT**



# Hand Sanitizer

**Antiseptic Gel  
With Vitamin E & Aloe**

***Kills 99.9% of Germs  
Without Water***

**240mL - (8 fl oz)**

796006 Rev. E Unit Carton Printing Plate for "A" size carton.



796006 Rev. E (page 3 of 3)

## FAK3SHLF-CLSB

777028B Rev. A  
white printed four color process  
blue (pms 072C) and red (pms 186C)



### **4267 Kit Contents FAK3SHLF-CLSB**

- 1 1X3 PLASTIC 100/BOX
- 1 FINGERTIP "T" WOVEN 40/BOX
- 1 SWIFT KNUCKLE 40/BX
- 1 1 X 3 WOVEN 100/BOX
- 3 NEOMYCIN ANTIBIOTIC 10 PER
- 1 AMMONIA INHALANTS 10 PER
- 2 EYE DRESS PKT W/4 ADH STRIPS
- 1 TOURNIQUET, 1 PER
- 1 WIRE SPLINT 1 PER
- 1 ADH BAND, EXTRA LARGE, 6 PER
- 1 ALCOHOL PREP PADS 10P
- 1 O/H PUMP ANTISEPTIC 2 OZ ID F
- 1 O/H PUMP BURN RELIEF 2 OZ ID G
- 1 FIRST AID GUIDE ASHI

2 TAPE ADHESIVE 1"X 5 YD PLSTC  
 10 HAND SANITIZER 0.9G WJ BULK  
 4 GAUZE CLEAN-WRAP BDGE N/S 2"  
 4 GAUZE CLEAN-WRAP BDGE N/S 4"  
 4 BLOODSTOPPER  
 1 NON-ADHERENT PADS 2"X3" 10'S  
 1 GZE PADS STERILE 2"X 2" 10'S  
 1 GZE PADS STERILE 3"X 3" 25'S  
 1 ELASTIC BANDAGE 3" X 4.5YD  
 1 CPR FILTERSHIELD 77-100  
 1 COTTON TIPS 100 PER VIAL  
 1 AYPANAL NON-ASP IND 2/ENV 100  
 1 4OZ BFS EYEWASH TRILINGUAL BOTTLE  
 1 SCISSOR BDGE 4" RED PLS HDL  
 1 KIT TWEEZER 3 1/2" SLANTED  
 1 180 EMPTY BLANK NO LOGO  
 1 POCKET INSERT RED #180 KIT 4R  
 1 TONGUE BLADES SR WRAPPED 6'S  
 1 LBL STOCK 6-3/8"X4"  
 1 LBL STOCK 4"X2-7/8"  
 1 LBL STOCK 3"x1-7/8"  
 4 PR LRG NITRILE GLVES ZIP BAG  
 2 WATER-JEL BURN DRESSING 4 X 4  
 1 LBL CONTENTS ANSI 2015 CL B  
 1 LBL CAB CVR ANSI 2015 CL B  
 2 TRI BNDG NON WOVEN 40"X40"X56"  
 2 COLD PACK UNIT 4"X6" BULK  
 1 RED BIO BAGS 2/BX

## 4267 FIRST AID KIT

4267 first aid kit kit

### Product Information

**Product Type**

HUMAN OTC DRUG

**Item Code (Source)**

NDC:0498-4267

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-4267-01	1 in 1 KIT; Type 0: Not a Combination Product	10/18/2018	

## Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	118 mL
Part 2	10 POUCH	4 mL
Part 3	50 PACKET	100
Part 4	1 BOTTLE, SPRAY	59 mL
Part 5	1 BOTTLE, SPRAY	59 mL
Part 6	30 PACKET	27 g
Part 7	10 PACKAGE	9 mL
Part 8	10 AMPULE	3 mL

## Part 1 of 8

### EYESALINE EMERGENCY EYEWASH

purified water liquid

## Product Information

Item Code (Source)	NDC:0498-0100
Route of Administration	OPHTHALMIC

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
WATER (UNII: 059QF0KO0R) (WATER - UNII:059QF0KO0R)	WATER	98.6 mL in 100 mL

## Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM PHOSPHATE, DIBASIC (UNII: GR686LBA74)	
SODIUM PHOSPHATE, MONOBASIC, MONOHYDRATE (UNII: 593YOG76RN)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:0498-0100-02	118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
---	------------------	---	--	--

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M018	12/18/2018	

## Part 2 of 8

### ALCOHOL WIPE

isopropyl alcohol swab

## Product Information

Item Code (Source)	NDC:0498-0143
Route of Administration	TOPICAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	0.7 mL in 1 mL

## Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0143-04	0.4 mL in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	

## Part 3 of 8

### AYPANAL NON-ASPIRIN

acetaminophen tablet

#### Product Information

**Item Code (Source)** NDC:0498-2001

**Route of Administration** ORAL

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

#### Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	

#### Product Characteristics

<b>Color</b>	white	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	circle;U
<b>Contains</b>			

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		2 in 1 PACKET; Type 0: Not a Combination Product		

#### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		04/10/2012	

## Part 4 of 8



# BURN RELIEF

lidocaine hydrochloride spray

## Product Information

**Item Code (Source)** NDC:0498-0221

**Route of Administration** TOPICAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LIDOCAINE HYDROCHLORIDE</b> (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE ANHYDROUS	24.64 mg in 1 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>TROLAMINE</b> (UNII: 9O3K93S3TK)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>OCTOXYNOL-9</b> (UNII: 7JPC6Y25QS)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>TEA TREE OIL</b> (UNII: VIF565UC2G)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>DIAZOLIDINYL UREA</b> (UNII: H5RIZ3MPW4)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0221-59	59 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	

Part 5 of 8

ANTISEPTIC

benzalkonium chloride spray

### Product Information

**Item Code (Source)** NDC:0498-0402

**Route of Administration** TOPICAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BENZALKONIUM CHLORIDE</b> (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII: 7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	0.13 g in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>OCTOXYNOL 9</b> (UNII: 7JPC6Y25QS)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>DIAZOLIDINYL UREA</b> (UNII: H5RIZ3MPW4)	
<b>DIPROPYLENE GLYCOL</b> (UNII: E107L85C40)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>TROLAMINE</b> (UNII: 9O3K93S3TK)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0402-59	59 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	

### Part 6 of 8

### NEOMYCIN

antibiotic ointment

## Product Information

Item Code (Source) NDC:0498-0730

Route of Administration TOPICAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN SULFATE	3.5 mg in 1 g

## Inactive Ingredients

Ingredient Name	Strength
PETROLATUM (UNII: 4T6H12BN9U)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0730-01	0.9 g in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		03/31/2010	

## Part 7 of 8

### INSTANT HAND SANITIZER

alcohol liquid

## Product Information

Item Code (Source) NDC:59898-420

Route of Administration TOPICAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALCOHOL (UNII: 3K9958V90M) (ALCOHOL - UNII:3K9958V90M)	ALCOHOL	62 mL in 100 mL

## Inactive Ingredients

Ingredient Name	Strength
ALOE VERA LEAF (UNII: ZY81Z83H0X)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
.ALPHA.-TOCOPHEROL ACETATE, DL- (UNII: WR1WPI7EW8)	
CARBOMER COPOLYMER TYPE A (UNII: 71DD5V995L)	
WATER (UNII: 059QF0KO0R)	
TRISOPROPANOLAMINE (UNII: W9EN9DLM98)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59898-420-36	0.9 mL in 1 PACKAGE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		04/15/2010	

## Part 8 of 8

### AMMONIA INHALENT

ammonia inhalent inhalant

## Product Information

Item Code (Source)	NDC:0498-3334
Route of Administration	RESPIRATORY (INHALATION)

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
AMMONIA (UNII: 5138Q19F1X) (AMMONIA - UNII:5138Q19F1X)	AMMONIA	0.045 g in 0.3 mL

## Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-3334-00	0.3 mL in 1 AMPULE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	12/01/2022

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		10/18/2018	

**Labeler** - Honeywell Safety Products USA, INC (118768815)

Revised: 1/2024

Honeywell Safety Products USA, INC