SPONGIA TOSTA- spongia officinalis skeleton, roasted pellet Boiron

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

Spongia tosta 1M

Spongia tosta 1M HPUS

Active ingredient**: See product name on front panel (contains 0.443 mg of the active ingredient per pellet).

Uses: See symptoms on front panel.

Relieves dry, barking cough *

Stop use and ask a doctor if symptoms persist for more than 3 days or worsen.

If pregnant or breast-feeding, ask a health professional before use.

Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

lactose, sucrose

BoironUSA.com Info@boiron.com 1-800-BOIRON-1 (1-800-264-7661)

Distributed by Boiron, Inc.

Newtown Square, PA 19073

Do not use if pellet dispenser seal is broken.

Contains approx 80 pellets.

How to dispense pellets? Turn tube upside down. Twist until 5 pellets are dispensed into cap. Carefully remove the cap and use it to pour pellets under the tongue.

*CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED.

*C,K,CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.



Drug Facts

Active ingredient": See product name on front panel (contains 0.443 mg of the active ingredient per pellet).

Uses: See symptoms on front panel.

Warnings: Stop use and ask a doctor if symptoms persist for more than 3 days or worsen. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Directions: • Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.



Spongia tosta

1 m

HPUS NDC 0220-4872-41



Dry, barking cough.*

Made in France

Spongia tosta

MEDICAL

Exp:

69823 4

Contains approx. 80 pellets. US Peel for Drugs Facts and instructions for use.

06960

IOMEOPATHIC PRACTICE, E. NOT FDA EVALUATED.

Drug Facts

Active ingredient**: See product name on front panel (contains 0.443 mg of the active ingredient per pellet).

Uses: See symptoms on front panel.

Warnings: Stop use and ask a doctor if symptoms persist for more than 3 days or worsen. If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children.

Directions: Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.


Other information: ■ Do not use if pellet dispenser seal is broken.

Drug Facts (continued)

Inactive ingredients: lactose, sucrose

Questions or comments?

BoironUSA.com Info@Boiron.com



SPONGIA TOSTA

spongia officinalis skeleton, roasted pellet

Product	Inform	ation
Product	Intorm	ation

Product Type HUMAN OTC DRUG Item Code (Source) NDC:0220-4872

Route of Administration ORAL

Active Ingredient/Active Moiety

Basis of Strength Ingredient Name Strength SPONGIA OFFICINALIS SKELETON, ROASTED (UNII: 1PIP394IID) (SPONGIA

OFFICINALIS SKELETON, ROASTED - UNII:1PIP394IID)

SPONGIA OFFICINALIS SKELETON, ROASTED

1 [hp M] in 1 [hp_M]

Inactive Ingredients

Strength **Ingredient Name** SUCROSE (UNII: C151H8M554)

LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G)

Product Characteristics

Color	white	Score	
Shape	ROUND	Size	4mm
Flavor		Imprint Code	
Contains			

P	ackaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0220-4872- 41	1 [hp_M] in 1 TUBE; Type 0: Not a Combination Product	03/03/1983	

Marketing Information		
Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
	03/03/1983	
	Application Number or Monograph	Application Number or Monograph Marketing Start Citation Date

Labeler - Boiron (282560473)

Registrant - Boiron, Inc. (014892269)

Establishment			
Name	Address	ID/FEI	Business Operations
Boiron		282560473	manufacture(0220-4872)

Revised: 7/2024 Boiron