SPONGIA TOSTA- spongia officinalis skeleton, roasted pellet Boiron

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

Spongia tosta 30C

Spongia tosta 30C HPUS

Active ingredient**: See product name on front panel (contains 0.443 mg of the active ingredient per pellet).

Uses: See symptoms on front panel.

Relieves dry, barking cough *

Stop use and ask a doctor if symptoms persist for more than 3 days or worsen.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

lactose, sucrose

BoironUSA.com Info@boiron.com 1-800-BOIRON-1 (1-800-264-7661)

Distributed by Boiron, Inc.

Newtown Square, PA 19073

Do not use if pellet dispenser seal is broken.

Contains approx 80 pellets.

How to dispense pellets? Turn tube upside down. Twist until 5 pellets are dispensed into cap. Carefully remove the cap and use it to pour pellets under the tongue.

*CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED.

*C,K,CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.



Drug Facts

Active ingredient :: See product name on front panel (contains 0.443 mg of the active ingredient per pellet).

Uses: See symptoms on front panel.

Warnings: Stop use and ask a doctor if symptoms persist for more than 3 days or worsen. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Directions: ■ Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.







Lot:

Exp:

HOMEOPATHIC PRACT

3 06960 69813 5Contains approx. 80 pellets. US

Peel for Drugs Facts and instructions for use.

Drug Facts

Active ingredient**: See product name on front panel (contains 0.443 mg of the active ingredient per pellet).

Uses: See symptoms on front panel.

Warnings: Stop use and ask a doctor if symptoms persist for more than 3 days or worsen. If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children.

Directions: ■ Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.


Other information: ■ Do not use if pellet dispenser seal is broken.

Drug Facts (continued)

Inactive ingredients: lactose, sucrose

Questions or comments?

Dairon ICA com Info@Dairon com



SPONGIA TOSTA

spongia officinalis skeleton, roasted pellet

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:0220-4870

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name

SPONGIA OFFICINALIS SKELETON, ROASTED (UNII: 1PIP394IID) (SPONGIA SPONGIA OFFICINALIS SKELETON, ROASTED - UNII:1PIP394IID)

Basis of Strength

SPONGIA OFFICINALIS SPONGIA OFFICINALIS SKELETON, ROASTED in 30 [hp_C]

Inactive Ingredients		
Ingredient Name	Strength	
SUCROSE (UNII: C151H8M554)		
LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G)		

Product Characteristics			
Color	white	Score	
Shape	ROUND	Size	4mm
Flavor		Imprint Code	
Contains			

Packaging					
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:0220-4870- 41	30 [hp_C] in 1 TUBE; Type 0: Not a Combination Product	03/03/1983	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
unapproved homeopathic		03/03/1983		
homeopathic		03/03/1303		

Labeler - Boiron (282560473)

Registrant - Boiron, Inc. (014892269)

Establishment			
Name	Address	ID/FEI	Business Operations
Boiron		282560473	manufacture(0220-4870)

Revised: 7/2024 Boiron