SPONGIA TOSTA- spongia officinalis skeleton, roasted pellet Boiron

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

Spongia tosta 5C

Spongia tosta 5C HPUS

Active ingredient**: See product name on front panel (contains 0.443 mg of the active ingredient per pellet).

Uses: See symptoms on front panel.

Relieves dry, barking cough *

Stop use and ask a doctor if symptoms persist for more than 3 days or worsen.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

lactose, sucrose

BoironUSA.com Info@boiron.com 1-800-BOIRON-1 (1-800-264-7661)

Distributed by Boiron, Inc. Newtown Square, PA 19073

Do not use if pellet dispenser seal is broken.

Contains approx 80 pellets.

How to dispense pellets? Turn tube upside down. Twist until 5 pellets are dispensed into cap. Carefully remove the cap and use it to pour pellets under the tongue.

*CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED.

*C,K,CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.



Drug Facts

Active ingredient :: See product name on front panel (contains 0.443 mg of the active ingredient per pellet).

Uses: See symptoms on front panel.

Warnings: Stop use and ask a doctor if symptoms persist for more than 3 days or worsen. If pregnant or breast-feeding, ask a health professional before use, Keep out of reach of children.

Directions: • Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.





Lot:

Exp:

VIEOPATHIC PRACTI NOT FDA EVALUATI

3 06960 69807 4

Contains approx. 80 pellets. US

Peel for Drugs Facts and instructions for use.

Drug Facts

Active ingredient***: See product name on front panel (contains 0.443 mg of the active ingredient per pellet).

Uses: See symptoms on front panel.

Warnings: Stop use and ask a doctor if symptoms persist for more than 3 days or worsen. If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children.


Directions: ■ Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

Other information: ■ Do not use if pellet dispenser seal is broken.

Drug Facts (continued) **Inactive ingredients:** lactose, sucrose

Questions or comments?

BoironUSA.com Info@Boiron.com



SPONGIA TOSTA

spongia officinalis skeleton, roasted pellet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0220-4865
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
SPONGIA OFFICINALIS SKELETON, ROASTED (UNII: 1PIP394IID) (SPONGIA OFFICINALIS SKELETON, ROASTED - UNII:1PIP394IID)	SPONGIA OFFICINALIS SKELETON, ROASTED	5 [hp_C] in 5 [hp_C]		

Inactive Ingredients			
Ingredient Name	Strength		
LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G)			
SUCROSE (UNII: C151H8M554)			

Product Characteristics			
Color	white	Score	
Shape	ROUND	Size	4mm
Flavor		Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:0220-4865- 41	5 [hp_C] in 1 TUBE; Type 0: Not a Combination Product	03/03/1983	

Marketing Information			
Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
	03/03/1983		
	Application Number or Monograph	Application Number or Monograph Marketing Start Citation Date	

Labeler - Boiron (282560473)

Registrant - Boiron, Inc. (014892269)

Establishment				
Name	Address	ID/FEI	Business Operations	
Boiron		282560473	manufacture(0220-4865)	

Revised: 7/2024 Boiron