TUSSIN COUGH CHEST CONGESTION- dextromethorphan hydrobromide, guaifenesin solution Rite Aid Corporation

Rite Aid Corporation Tussin Cough Chest Congestion Drug Facts

Active ingredients (in each 20 mL)

Dextromethorphan HBr, USP 20 mg Guaifenesin, USP 400 mg

Purposes

Cough suppressant

Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if

cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- · keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

| age | dose | |
|---------------------------------------|---------------------|--|
| adults and children 12 years and over | 20 mL every 4 hours | |
| children under 12 years | do not use | |

Other information

- each 20 mL contains: sodium 13 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

acetic acid, anhydrous citric acid, carboxymethylcellulose sodium, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, menthol, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to the active ingredients in Robitussin $^{\circledR}$ Maximum Strength Cough + Chest Congestion DM

FREE FROM

GLUTEN FREE

ALCOHOL FREE

MAXIMUM STRENGTH

TUSSIN

COUGH + CHEST CONGESTION

DEXTROMETHORPHAN HBr

COUGH SUPPRESSANT

GUAIFENESIN/EXPECTORANT

ADULT • NON-DROWSY

Relieves chest congestion, controls cough, thins & loosens mucus

For ages 12 & over

DM MAX

RASPBERRY MENTHOL FLAVOR

4 FL OZ (118 mL)



TUSSIN COUGH CHEST CONGESTION

dextromethorphan hydrobromide, quaifenesin solution

| Product Information | | | |
|-------------------------|----------------|--------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:11822-1332 |
| Route of Administration | ORAL | | |
| | | | |

| Active Ingredient/Active Moiety | | | |
|--|----------------------------------|--------------------|--|
| Ingredient Name | Basis of Strength | Strength | |
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 20 mg in 20 mL | |
| GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ) | GUAIFENESIN | 400 mg in 20 mL | |

| Inactive Ingredients | | | |
|---|----------|--|--|
| Ingredient Name | Strength | | |
| ACETIC ACID (UNII: Q40Q9N063P) | | | |
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | | | |
| CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K6790BS311) | | | |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | | | |
| FD&C RED NO. 40 (UNII: WZB9127XOA) | | | |
| GLYCERIN (UNII: PDC6A3C0OX) | | | |
| MENTHOL, UNSPECIFIED FORM (UNII: L7T10EIP3A) | | | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | | | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | | | |
| WATER (UNII: 059QF0KO0R) | | | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | | | |
| SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR) | | | |
| SORBITOL SOLUTION (UNII: 8KW3E207O2) | | | |
| SUCRALOSE (UNII: 96K6UQ3ZD4) | | | |
| XANTHAN GUM (UNII: TTV12P4NEE) | | | |

| Product Characteristics | | | |
|-------------------------|-------|--------------|--|
| Color | RED | Score | |
| Shape | | Size | |
| Flavor | FRUIT | Imprint Code | |
| Contains | | | |

| P | Packaging | | | | | |
|---|----------------------|---|-------------------------|-----------------------|--|--|
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date | | |
| 1 | NDC:11822- 1332-1 | 1 in 1 CARTON | 03/02/2023 | | | |
| 1 | | 118 mL in 1 BOTTLE; Type 0: Not a Combination Product | | | | |
| 2 | NDC:11822- 1332-2 | 1 in 1 CARTON | 03/02/2023 | | | |
| 2 | | 237 mL in 1 BOTTLE; Type 0: Not a Combination Product | | | | |

| Marketing Information | | | | |
|-----------------------|---|-------------------------|-----------------------|--|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | |
| OTC Monograph Drug | M012 | 03/02/2023 | | |

Labeler - Rite Aid Corporation (014578892)

Revised: 10/2024 Rite Aid Corporation