

**VICKS CHILDRENS COUGH CONGESTION NIGHT- phenylephrine hydrochloride and diphenhydramine hydrochloride liquid**  
**The Procter & Gamble Manufacturing Company**

-----  
**VICKS®**  
**children's Cough Congestion Night**

***Drug Facts***

**Active ingredients (in each 15 mL)**

Diphenhydramine HCl 12.5 mg

Phenylephrine HCl 5 mg

**Purpose**

Antihistamine/Cough suppressant  
Nasal decongestant

**Uses**

temporarily relieves common cold symptoms:

- nasal congestion
- runny nose & sneezing
- cough due to minor throat and bronchial irritation

**Warnings**

**Do not use**

- with any other product containing diphenhydramine, even one used on skin
- to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, or emphysema

- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are**

taking sedatives or tranquilizers

**When using this product**

- **do not use more than directed.**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- symptoms do not improve within 7 days or occur with a fever
- cough persists for more than 7 days, comes back or occurs with a fever, rash or persistent headache

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Other information**

- **each 15 mL contains:**sodium 44 mg
- store at no greater than 25°C.

**Directions**

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

---

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	do not use unless directed by a doctor
children under 4 yrs	do not use

---

## **Other information**

- **each 15 mL contains:**sodium 44 mg
- store at no greater than 25°C.

## **Inactive ingredients**

anhydrous citric acid, flavor, glycerin, polysorbate 20, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

## **Questions?**

**1-800-362-1683**

**TAMPER EVIDENT:**Do not use if printed safety seal under cap is missing or damaged.

**DIST. BY: PROCTER & GAMBLE, CINCINNATI, OH 45202**

**Made in Canada**

## **PRINCIPAL DISPLAY PANEL - 177 ml Bottle Label**

**VICKS®**

children's

### **Cough**

Congestion **Night**

Phenylephrine HCl - nasal decongestant

Diphenhydramine HCl - antihistamine/cough suppressant

**FREE**of:

Artificial Dyes & Flavors,

High Fructose Corn Syrup & Alcohol

- Cough
- Runny nose
- Stuffy nose
- Sneezing

Ages 6+

6 FL OZ (177 ml)



Drug Facts	
<b>Active ingredients (in each 15 mL)</b> Diphenhydramine HCl 12.5 mg Phenylephrine HCl 5 mg	<b>Purpose</b> Antihistamine/Cough suppressant Nasal decongestant
<b>Uses</b> temporarily relieves common cold symptoms: • nasal congestion • runny nose & sneezing • cough due to minor throat & bronchial irritation	
<b>Warnings</b> <b>Do not use</b> • with any other product containing diphenhydramine, even one used on skin • to make a child sleepy • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. <b>Ask a doctor before use if you have</b> • heart disease • high blood pressure • thyroid disease • diabetes • glaucoma • trouble urinating due to enlarged prostate gland • cough that occurs with the rash-prone drug (moxal) • a breathing problem or chronic cough that lasts or occurs with smoking, asthma or emphysema • a sodium-restricted diet <b>Ask a doctor or pharmacist before use if you are</b> taking sedatives or tranquilizers	
<b>When using this product</b> • do not use more than directed. • excitability may occur, especially in children • medical conditions may occur • alcohol, sedatives, and tranquilizers may increase drowsiness • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery	
<b>Stop use and ask a doctor if</b> • you get nervous, dizzy or sleepless • symptoms do not improve within 7 days or occur with a fever • cough persists for more than 7 days, comes back or occurs with a fever, rash or persistent headache These could be signs of a serious condition.	
<b>If pregnant or breast-feeding</b> , ask a health professional before use. <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.	
<b>Directions</b> • take only as directed • only use the dose cup provided • do not exceed 4 doses per 24 hrs adults & children 12 yrs & over 30 mL every 4 hrs children 6 to under 12 yrs 15 mL every 4 hrs children 4 to under 6 yrs do not use unless directed by a doctor children under 4 yrs do not use	
<b>Other information</b> • each 15 mL contains: sodium 44 mg • Store at or greater than 20°C.	
<b>Inactive ingredients</b> anhydrous citric acid, flavor, glycerin, polyorbate 20, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose	



## VICKS CHILDRENS COUGH CONGESTION NIGHT

phenylephrine hydrochloride and diphenhydramine hydrochloride liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:37000-712
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	12.5 mg in 15 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>POLYSORBATE 20</b> (UNII: 7T1F30V5YH)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3Z D4)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	

**SACCHARIN SODIUM** (UNII: SB8ZUX40TY)

**SORBITOL** (UNII: 506T60A25R)

### Product Characteristics

<b>Color</b>	white (Clear)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	GRAPE	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37000-712-06	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/22/2019	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/22/2019	

**Labeler** - The Procter & Gamble Manufacturing Company (004238200)

Revised: 10/2025

The Procter & Gamble Manufacturing Company