

**EXTRA STRENGTH NO-PAIN- acetaminophen tablet**  
**Safrel Pharmaceuticals, LLC.**

-----  
**Acetaminophen Extra Strength 500 mg**

**Active Ingredient**  
**(in each tablet)**

Acetaminophen 500mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporary relieves minor aches and pains due to
- the common cold
- headache
- backache
- toothache
- muscular aches
- premenstrual and menstrual cramps and
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- liver disease

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding**

ask a health professional before use

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Do not exceed recommended dosage.**

**Directions**

- do not use more than directed (**see overdose warning**)
- Adults and children 12 years of age and older:
- Take 2 caplets every 6 hours while symptoms last.
- Do not take more than 6 caplets in 24 hours, unless directed by a doctor
- Do not use more than 10 days unless directed by a doctor.
- Children under 12 years of age: Do not use this extra strength product. This will provide more than the recommended dose (overdose) and could cause serious health problems.

**Other Information**

- store at controlled room temperature 20-25°C (68-77°F).
- read all product information before using.
- TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

**Inactive Ingredients**

Hypromellose, Magnesium Stearate, Microcrystalline Cellulose, Pregelatinized Starch, Polyethylene Glycol, Polyvinyl Pyrrolidone, Stearic Acid, Talc, Titanium Dioxide

1-844-384-3723 (Mon-Fri 9AM-5PM EST) or [www.safrelpharma.com](http://www.safrelpharma.com)

\* This product is not manufactured or distributed by McNeil-Consumer Healthcare, owner of the registered trademark Tylenol\*.

## Drug Facts (continued)

**Stop use and ask a doctor if** ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

<ul style="list-style-type: none"> <li>do not take more than directed (see overdose warning)</li> </ul>	<ul style="list-style-type: none"> <li>take 2 caplets every 6 hours while symptoms last</li> <li>do not take more than 6 caplets in 24 hours, unless directed by a doctor</li> <li>do not use for more than 10 days unless directed by a doctor</li> </ul>
adults and children 12 years and over	
children under 12 years	ask a doctor

### Other information

■ store at 20-25°C (68-77°F)

### Inactive ingredients

carnauba wax, hypromellose, mineral oil, polyethylene glycol, povidone, pregelatinized starch, sodium carboxymethyl cellulose, stearic acid, titanium dioxide

### Questions or Comments? 1-844-384-3723

\*Compare to Extra Strength Tylenol® Active Ingredient

NDC 71309-001-05



## Extra Strength NO-PAIN

# ACETAMINOPHEN

- Pain Reliever
- Fever Reducer
- For Adults
- Aspirin Free

500 CAPLETS

actual size

## Drug Facts

Active Ingredient (in each caplet)	Purpose
Acetaminophen USP 500 mg	Pain Reliever/Fever Reducer

**Uses** ■ temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ backache ■ minor pain of arthritis ■ the common cold ■ toothache ■ premenstrual and menstrual cramps

■ temporarily reduces fever

**Warnings** **Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do Not Use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have** liver disease

**Ask a doctor before use if you are taking** the blood thinning drug warfarin

Distributed By: Safrel Pharmaceuticals  
Bridgewater, NJ 08807 USA  
www.safrelpharma.com

LOT.

EXP.

GLUTEN FREE

DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING

## acetaminophen tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71309-001
Route of Administration	ORAL		

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Ingredient Name	Strength
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
TITANIUM DIOXIDE (UNII: 15FIX9V2IP)	

**TALC** (UNII: 7SEV7J4R1U)

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	18mm
<b>Flavor</b>		<b>Imprint Code</b>	BH
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71309-001-05	500 in 1 BOTTLE; Type 0: Not a Combination Product	07/31/2017	
2	NDC:71309-001-40	1 in 1 CARTON	07/31/2017	
2		40 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/31/2017	

**Labeler** - Safrel Pharmaceuticals, LLC. (080566287)

Revised: 4/2025

Safrel Pharmaceuticals, LLC.