

# **IBUPROFEN PAIN RELIEVER/FEVER REDUCER- ibuprofen capsule, liquid filled Amneal Pharmaceuticals LLC**

-----

## **Ibuprofen Capsules (liquid filled, 200 mg)**

### **OTC - ACTIVE INGREDIENT**

(in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*

(present as the free acid and potassium salt)

\* nonsteroidal anti-inflammatory drug

### **OTC - PURPOSE**

Pain reliever/Fever reducer

### **INDICATIONS AND USAGE**

#### ***Uses***

- temporarily relieves minor aches and pains due to:
- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

### **WARNINGS**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause

severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

## **OTC - DO NOT USE**

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

## **OTC - ASK DOCTOR**

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

## **OTC - ASK DOCTOR/PHARMACIST**

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

## **OTC - WHEN USING**

### **When using this product**

- take with food or milk if stomach upset occurs

## **OTC - STOP USE**

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

## **OTC - PREGNANCY OR BREAST FEEDING**

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

## **OTC - KEEP OUT OF REACH OF CHILDREN**

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

## **DOSAGE AND ADMINISTRATION**

### ***Directions***

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

## **OTHER INFORMATION**

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use. Keep carton.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F)

## **INACTIVE INGREDIENT**

FD&C Blue # 1, gelatin, polyethylene glycol, potassium hydroxide, purified water, sorbitol

## OTC - QUESTIONS OR COMMENTS?

Call 1-877-835-5472

Monday through Friday 9AM - 5PM EST.

Read and keep carton for complete warnings and information.

Do not use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Distributed by:

**Amneal Pharmaceuticals LLC**

Bridgewater, NJ 08807

\*Amneal Pharmaceuticals is not affiliated with the owner of the trademark Advil®

Rev. 06-2022-02

## PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

NDC 65162-356-02

Ibuprofen Capsules, 200 mg

20 Capsules

Amneal Pharmaceuticals LLC

\*Compare to the active ingredient in Advil® LIQUI-GELS®  
NDC 65162-356-02  
**SEE NEW WARNINGS INFORMATION**  
**Ibuprofen Capsules**  
**200 mg**  
Pain reliever / Fever reducer (NSAID)  
A356  
Important: Read all product information before using.  
SOFT GELATIN CAPSULES\*\*  
\*\* (Liquid Filled Capsules)  
20 Capsules  
amneal

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**  
**Active ingredient (in each capsule)** Purpose  
Solubilized ibuprofen equal to Pain reliever/  
200 mg ibuprofen (NSAD)\* ..... Fever reducer  
(present as the free acid and potassium salt)  
\*nonsteroidal anti-inflammatory drug

**Uses**  
temporarily relieves minor aches and pains due to:  
■ headache ■ toothache  
■ backache ■ menstrual cramps  
■ the common cold ■ muscular aches  
■ minor pain of arthritis  
■ temporarily reduces fever

Do not use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.  
Distributed by:  
Amneal Pharmaceuticals LLC  
Bridgewater, NJ 08807  
Rev. 06-2022-02

N 3 65162 356 02 1

Lot No:  
Exp. Date:

Non-Varnish Area

(continued)

rofen may cause a  
ion, especially in people  
/mptoms may include:  
acial swelling  
3)  
skin reddening  
dlisters  
1 occurs, stop use and  
ght away.  
**warning:** This product  
which may cause severe  
he chance is higher if  
ar  
1 ulcers or bleeding  
ing (anticoagulant) or  
: containing prescription  
1 NSAIDs (aspirin,  
en, or others)  
coholic drinks every day  
roduct  
1 longer time than  
**stroke warning:**  
rin, increase the risk of  
ailure, and stroke. These  
s is higher if you use  
r for longer than

ad an allergic reaction  
reliever/fever reducer  
er heart surgery

**re use if**  
warning applies to you  
is or serious side effects  
elievers or fever  
y of stomach problems,  
1

## Drug Facts (continued)

### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

### Other information

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use. Keep carton.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F)

### Inactive ingredients

FD&C Blue # 1, gelatin, polyethylene glycol, potassium hydroxide, purified water, sorbitol

### Questions or Comments?

Call 1-877-835-5472  
Monday through Friday 9AM - 5PM EST.

## Drug Facts (continued)

- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

### Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

### When using this product

- take with food or milk if stomach upset occurs

### Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

## Drug Facts (

### Warnings

**Allergy alert** Ibuprofen may cause severe allergic reactions in people who are allergic to aspirin. Symptoms include:

- hives
- asthma (wheezing)
- shock
- rash
- swelling

If an allergic reaction occurs, seek medical help right away.

### Stomach bleeding

Ibuprofen may cause stomach bleeding. This risk is higher if you:

- are age 60 or older
- have had stomach problems
- take a blood thinner or steroid drug
- take other drugs, such as aspirin, or nonprescription pain relievers, such as ibuprofen, naproxen
- have 3 or more alcohol drinks a day while using this product
- take more or for a longer time than directed

### Heart attack and stroke

Ibuprofen, like aspirin, may increase the risk of heart attack, heart failure, and stroke. The risk is higher if you have had heart disease or more than directed or for a longer time than directed.

### Do not use

- if you have ever had a heart attack or stroke
- right before or after surgery

### Ask a doctor before

- stomach bleeding
- you have problems with your stomach or intestines
- you have a history of heartburn

Exp. Date:

Lot No:

N  
3 65162 356 02 1



SEE NEW WARNINGS INFORMATION

NDC 65162-356-02

# Ibuprofen Capsules

## 200 mg

A356

SOFT GELATIN CAPSULES\*\*  
\*\* (Liquid Filled Capsules)

Pain reliever /  
Fever reducer (NSAID)

20 Capsules

**Important:** Read all product information before using.

\*Compare to the active ingredient in Advil® LIQUI-GELS®

DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

### Drug Facts (continued)

#### Questions or Comments?

Call 1-877-835-5472  
Monday through Friday 9AM - 5PM EST.

Distributed by: **Amneal Pharmaceuticals LLC**  
Bridgewater, NJ 08807

\*Amneal Pharmaceuticals is not affiliated with the owner of the trademark Advil®

Rev. 06-2022-02

### Drug Facts (continued)

- leg swelling
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present in the painful area
  - any new symptoms appear
- If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
- Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

### Other information

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use.
- keep carton.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F)

### Inactive ingredients

FD&C Blue # 1, gelatin, polyethylene glycol, potassium hydroxide, purified water, sorbitol

### Drug Facts (continued)

aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

### Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

### When using this product

- take with food or milk, if stomach upset occurs

### Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech

### READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

### Drug Facts

#### Active ingredient (in each capsule)

Sulfated ibuprofen (equal to 200 mg ibuprofen (NSAID)\*, Fever reducer (present as the free acid and potassium salt) \*nonsteroidal anti-inflammatory drug

#### Purpose

Pain reliever/  
Fever reducer

#### Uses

- temporarily relieves minor aches and pains due to:
  - headache
  - toothache
  - backache
  - menstrual cramps
  - muscular aches
- minor pain or arthritis
- temporarily reduces fever

#### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
  - facial swelling
  - skin redness
  - shock
  - asthma (wheezing)
  - rash
  - blisters
- If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
  - have had stomach ulcers or bleeding problems
  - take a blood thinner (anticoagulant) or steroid drug
  - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
  - have 3 or more alcoholic drinks every day while using this product
  - take more or for a longer time than directed
- Heart attack and stroke warning:** NSAIDs, except

## IBUPROFEN PAIN RELIEVER/FEVER REDUCER

ibuprofen capsule, liquid filled

### Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:65162-356

Route of Administration

ORAL

### Active Ingredient/Active Moiety

Ingredient Name

Basis of Strength

Strength

<b>IBUPROFEN</b> (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg
---	-----------	--------

### Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)	
<b>SORBITOL</b> (UNII: 506T60A25R)	

### Product Characteristics

<b>Color</b>	blue	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	18mm
<b>Flavor</b>		<b>Imprint Code</b>	A356
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65162-356-02	1 in 1 CARTON	03/29/2013	
1		20 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:65162-356-69	1 in 1 CARTON	03/29/2013	
2		80 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:65162-356-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	03/29/2013	
4	NDC:65162-356-11	1000 in 1 BOTTLE; Type 0: Not a Combination Product	03/29/2013	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202300	03/29/2013	

**Labeler** - Amneal Pharmaceuticals LLC (123797875)

Revised: 12/2023

Amneal Pharmaceuticals LLC