

# **ALUMIER MD ACNE CLARIFYING CLEANSER- salicylic acid gel**

## **Alumier Labs**

-----

### **Alumier MD Acne Clarifying Cleanser**

#### **Drug Facts**

##### **Active Ingredient**

Salicylic Acid 2.0%

##### **Purpose**

Acne Treatment

##### **Use**

- for the treatment of acne

##### **Warnings**

- **For external use only.**

##### **When using this product**

- avoid contact with eyes. If contact occurs, rinse thoroughly with water.
- skin irritation and dryness is more likely to occur if you use another topical acne product at the same time. If irritation occurs, only use one topical acne product at a time.

**Stop use and ask a doctor** if you develop severe irritation, burning or itching of the skin, hives, swelling of eyes and mouth, blistering or difficulty breathing

- for more severe cases, and/or if improvement is not apparent within 6-8 weeks, consult a health care professional

**Keep out of reach of children.** If swallowed, call a Poison Control Centre or get medical help right away.

##### **Directions**

- use twice daily, in the morning and evening
- moisten face with lukewarm water
- pump a small amount into your hand
- add water and apply to your face and neck, massaging in a circular motion
- rinse thoroughly with lukewarm water and gently pat dry

##### **For New Users**

- apply product to a small area once a day for three days to test if you are sensitive to

this product

- if no discomfort occurs, cover the entire affected area with a thin layer
- start with one application daily, then gradually increase to two to three times daily, if needed
- if dryness or peeling occurs, reduce application to once a day or every other day

### **Other Information**

- Do not use if security seal is broken or missing
- You may report a serious adverse reaction event from using this product to: Alumier Labs Inc., 550 Cochituate Road, Suite 25, Framingham, MA 01701

### **Inactive Ingredients**

Water/Aqua/Eau, Sodium C14-16 Olefin Sulfonate, Propanediol, Dimethyl Lauramide/Myristamide, Glycerin, Cocamidopropyl Dimethylamine, Cocamidopropyl Hydroxysultaine, Glyceryl Caprate, Lauryl Lactyl Lactate, Oligopeptide-10, Aloe Barbadensis Leaf Juice, Boswellia Serrata Gum Extract, Honey Extract (Extrait de Miel), Oryza Sativa (Rice) Bran Extract, Carbomer, Sodium Hydroxide, Disodium EDTA, Caprylyl Glycol, Xylitol, Caprylic Acid, Ethylhexylglycerin.

### **Company Information**

#### **ALUMIER EUROPE LTD.**

5 ST. FREDERICK STREET  
VALLETTA VLT 1470, MALTA

#### **WWW.ALUMIERMD.COM**

EUROPE • UNITED STATES • CANADA  
UNITED KINGDOM

DISTRIBUTED IN CANADA BY  
ALUMIER LABS INC.,

436 LIMESTONE CRESCENT  
TORONTO, ON, M3J 2S4

DISTRIBUTED IN THE USA BY  
ALUMIER LABS INC.,

550 COCHITUATE ROAD, SUITE 25  
FRAMINGHAM, MA 01701

MANUFACTURED IN CANADA UNDER  
LICENSE FROM ALUMIER EUROPE LTD.

# Product Packaging

## Alumier MD

### Acne Clarifying Cleanser

#### ACNE TREATMENT WITH 2%

#### SALICYLIC ACID

NPN: 80064465

177 mL/6 fl oz

#### Alumier®.

ALUMIER EUROPE LTD.  
5 ST. FREDERICK STREET  
VALLETTA VLT 1470, MALTA

[WWW.ALUMIERMD.COM](http://WWW.ALUMIERMD.COM)

EUROPE - UNITED STATES - CANADA  
UNITED KINGDOM

DISTRIBUTED IN CANADA BY/  
DISTRIBUÉ AU CANADA PAR :  
ALUMIER LABS INC.,  
436 LIMESTONE CRESCENT  
TORONTO, ON, M3J 2S4

DISTRIBUTED IN THE USA BY/  
DISTRIBUÉ AUX ÉTATS-UNIS PAR :  
ALUMIER LABS INC.,  
550 COCHITUATE ROAD, SUITE 25  
FRAMINGHAM, MA 01701

MANUFACTURED IN CANADA UNDER  
LICENSE FROM ALUMIER EUROPE LTD.

FABRIQUÉ AU CANADA SOUS LICENCE  
OCTROYÉE PAR ALUMIER EUROPE LTD.

#### Notice

Ingrédient actif	Utilisation
Acide salicylique 2,0 %	Traitement de l'acné

**Usage** ■ pour le traitement de l'acné

#### Avertissements

■ Usage externe uniquement.

#### Quand on utilise ce produit

■ éviter le contact avec les yeux ; en cas de contact, rincer abondamment à l'eau.  
■ une irritation cutanée et une sécheresse sont plus susceptibles de se produire si vous utilisez un autre produit topique contre l'acné simultanément. En cas d'irritation, utilisez un seul produit topique contre l'acné à la fois.

**Arrêtez l'utilisation et demandez à un docteur** si vous développez de l'urticaire, un gonflement des yeux ou de la bouche, des cloques ou si vous avez des difficultés à respirer  
■ pour les cas plus graves, et/ou si une amélioration n'est pas constatée dans les 6 à 8 semaines, consulter un professionnel de santé

**Garder hors de portée des enfants.** En cas d'ingestion, contacter un centre antipoison ou appeler un médecin immédiatement.

**Conseils d'utilisation** ■ utiliser deux fois par jour, le matin et le soir ■ humidifier le visage à l'eau tiède ■ placer une petite quantité de produit dans la main ■ ajouter de l'eau et appliquer sur le visage et le cou en effectuant des massages circulaires ■ rincer soigneusement à l'eau tiède et sécher délicatement  
**Pour les nouveaux utilisateurs** ■ appliquer le produit sur une petite zone une fois par jour pendant trois jours afin de tester votre sensibilité au produit ■ si aucun inconfort ne survient, couvrir la zone affectée avec une fine couche de produit ■ commencer par une application quotidienne, puis augmenter progressivement à deux ou trois fois par jour si nécessaire ■ en cas de sécheresse ou de desquamation de la peau, réduire l'application à une fois par jour ou à tous les deux jours

#### Autres information

■ Ne pas utiliser si le sceau de sécurité est endommagé ou manquant.  
■ Vous pouvez signaler toute réaction indésirable liée à l'utilisation de ce produit à : Alumier Labs Inc., 550 Cochituate Road, Suite 25, Framingham, MA 01701



### Acne Clarifying Cleanser Nettoyant clarifiant pour acné

ACNE TREATMENT WITH 2%  
SALICYLIC ACID  
TRAITEMENT DE L'ACNÉ AVEC 2 %  
D'ACIDE SALICYLIQUE

NPN: 80064465

177 mL/6 fl oz e

#### Drug Facts

Active Ingredient	Purpose
Salicylic Acid 2.0%	Acne treatment

**Use** ■ for the treatment of acne

#### Warnings

■ For external use only.

#### When using this product

■ avoid contact with eyes. If contact occurs, rinse thoroughly with water.  
■ skin irritation and dryness is more likely to occur if you use another topical acne product at the same time. If irritation occurs, only use one topical acne product at a time.

**Stop use and ask a doctor** if you develop severe irritation, burning or itching of the skin, hives, swelling of eyes and mouth, blistering or difficulty breathing

■ for more severe cases, and/or if improvement is not apparent within 6-8 weeks, consult a health care professional

**Keep out of reach of children.** If swallowed, call a Poison Control Centre or get medical help right away.

**Directions** ■ use twice daily, in the morning and evening ■ moisten face with lukewarm water ■ pump a small amount into your hand ■ add water and apply to your face and neck, massaging in a circular motion ■ rinse thoroughly with lukewarm water and gently pat dry  
**For New Users** ■ apply product to a small area once a day for three days to test if you are sensitive to this product ■ if no discomfort occurs, cover the entire affected area with a thin layer ■ start with one application daily, then gradually increase to two to three times daily, if needed ■ if dryness or peeling occurs, reduce application to once a day or every other day

#### Other Information

■ Do not use if security seal is broken or missing.  
■ You may report a serious adverse reaction event from using this product to: Alumier Labs Inc., 550 Cochituate Road, Suite 25, Framingham, MA 01701

#### Inactive Ingredients/Ingrédients inactifs

1922414 - Water/Aqua/Eau, Sodium C14-16 Olefin Sulfonate, Propanediol, Dimethyl Lauramide/Myristamide, Glycerin, Cocamidopropyl Dimethylamine, Cocamidopropyl Hydroxysulfate, Glyceryl Caprate, Lauryl Lactyl Lactate, Oligopeptide-10, Aloe Barbadosensis Leaf Juice, Boswellia Serrata Gum Extract, Honey Extract (Extrait de Miel), Oryza Sativa (Rice) Bran Extract, Carbomer, Sodium Hydroxide, Disodium EDTA, Caprylyl Glycol, Xylitol, Caprylic Acid, Ethylhexylglycerin.



## ALUMIER MD ACNE CLARIFYING CLEANSER

salicylic acid gel

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69473-001
<b>Route of Administration</b>	TOPICAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>SALICYLIC ACID</b> (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ)	SALICYLIC ACID	20 mg in 1 mL

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>CAPRYLIC ACID</b> (UNII: OBL58JN025)	
<b>PROPANEDIOL</b> (UNII: 5965N8W85T)	
<b>COCAMIDOPROPYL HYDROXYSULTAINE</b> (UNII: 62V75NI93W)	
<b>GLYCERYL CAPRATE</b> (UNII: 197M6VFC1W)	
<b>OLIGOPEPTIDE-10</b> (UNII: Q46328TRNK)	
<b>HONEY</b> (UNII: Y9H1V576FH)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>COCAMIDOPROPYL DIMETHYLAMINE</b> (UNII: L36BM7DG2T)	
<b>ALOE VERA LEAF</b> (UNII: ZY81Z83H0X)	
<b>CAPRYLYL GLYCOL</b> (UNII: 00YIU5438U)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>RICE BRAN</b> (UNII: R60QEP13IC)	
<b>CARBOXYPOLYMETHYLENE</b> (UNII: 0A5MM307FC)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>XYLITOL</b> (UNII: VCQ006KQ1E)	
<b>ETHYLHEXYLGLYCERIN</b> (UNII: 147D247K3P)	
<b>INDIAN FRANKINCENSE</b> (UNII: 4PW41QCO2M)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM C14-16 OLEFIN SULFONATE</b> (UNII: O9W3D3YF5U)	

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:69473-001-01	1 in 1 CARTON	07/01/2019	
1		177 mL in 1 BOTTLE, PUMP; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M006	07/01/2019	

