SUNMARK ACID REDUCER- famotidine tablet, film coated Strategic Sourcing Services LLC

McKesson Acid Reducer Drug Facts

Active ingredient (in each tablet)

Famotidine 10 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water at any time from 15
 to 60 minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture

Inactive ingredients

carnauba wax, hypromellose, iron oxide red, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, pregelatinized starch, talc, titanium dioxide, triacetin

Questions or comments?

1-800-719-9260

Principal Display Panel

COMPARE TO PEPCID[®] AC ACTIVE INGREDIENT See New Warnings famotidine tablets, 10 mg acid reducer Just one tablet prevents & relieves heartburn due to acid indigestion

ORIGINAL STRENGTH

ACTUAL SIZE 30 TABLETS

DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

JUST ONE TABLET prevents and re lieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

Supmarkorand.com Compare to PEPCID® AC ACTIVE INGREDIENT* NDC 49348-128-44

famotidine tablets, 10 mg acid reducer

MSKESSON

Another Quality Product Distributed By McKesson 6565 State Highway 161, Irving, TX 75039 Money Back Guarantee Please visit us at www.sunmar kbrand.com

> Just one tablet prevents & relieves heartburn due to acid indigestion

> > **ORIGINAL STRENGTH**

ACTUAL SIZE

30 TABLETS

	Drug Facts	Drug Facts (continued)	a a a
	Active ingredient (in each tablet) Purpose Famotidine 10 mgAcid reducer	Aska d octo ror pharmacist before use if you a re taking a prescription drug. Acid reducers may interact with certain prescription drugs.	e Avoid Avoid d and cer uit smokin
	Uses: relieves hearburn associated with acid indigestion and sourstomach	Stopuse and aska doctorif your heartburn continues or worsens you need to take thisproduct for more than 14 days	Do not wear fight-fifting of your bed • Avoid affeine, alcohol and cert sweight • Outhsmokin
ę	prevents heartburn as sociated with acid indigestion and sourstomach brought on by eating or drinking certain food and beverages	If pregnant or breast-feed in g, ask a health professional bebreuse. Keepoutof reach of children. In case of overdose,	atting • Do the head of colate, caff sight, lose w 3 TAB
OPEN	Warn ings Allergy al ert: Do not us ei f you are allergic to famotidine	get me dical help or contact a Poison Control Center right away. (1-8 00-222-12 22)	re after ea • Raise th oods, choo ff overweig OSING
OTHER	or other acid reducers Donot use ■ if you have trouble or pain swalkowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ with other acid reducers	Directions adults and children 1 2 years and over: to relie ve symptoms, s wallow 1 tablet with a glass of water. Do not chew. to prevent symptoms, s wallow 1 tablet with a glass of water atany time from 1 5 to 6 0 minutes before	TRC assistant
E	Aska dioctor before use if you have had hear burn over 3 months. This may be a sign of a more serious condition.	eating food or drinking beverages thatcause heartburn III do not use more fran 2 tablets in 24 hours III children under 12 years: ask a doctor	um • Do not lie flat c • Do not eat before th as rich, spicy, fatty slowly and avoid big CONV ENIENT
	 hearburn with lightheadedness, sweating, or dizz iness ches tp ain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadednes s 	Other information read the directions and warningsbefore use keep the carton. It contains important information. store at 20°-25°C (68°-77°F) protect from mois ture	ing Heartburn the sto mach • ing foots such ables • Eat slo CC
	■ frequent ch est pa in ■ frequent whe ezing, particul arly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain ■ kidney disease	In active ingredients canaiba wax, hypromellose, iron oxi de red, iron oxi de yellow, magnesium s barate, microcrystallinec elulose, polydextrose, poly ethylere glycol, pregelatinized s tarch, talc, titanium dioxi de, triacetin	The for Managing Hearthum conting around the stomach hardhum-causing foods such fruits and vegetables • Eat slo CC

14165 S 1 C2





fam	notidine tablet	ACID REI , film coated							
Pr	oduct Infor	mation							
Pr	oduct Type		HUMAN OTC DR	RUG I	tem C	ode (S	ource)	NDC:49	348-128
	ute of Admini	stration	ORAL						
۵c	tive Ingredi	ent/Active	Moiety						
	the ingrea		dient Name				Basis of St	ronath	Strength
FΔ		-	(FAMOTIDINE - U	INII:5070151	278)		FAMOTIDINE	lengti	10 mg
		5920155220		11.5 Q 2 0 1 5 J	2207				10 mg
In	active Ingre	dients							
			Ingredien	t Name				:	Strength
CA	RNAUBA WAX (UNII: R12CBM0	EIZ)						
ΗY	PROMELLOSE,	UNSPECIFIED	(UNII: 3NXW29V3	SWO)					
MA	GNESIUM STEA	RATE (UNII: 70	097M6I30)						
MI	CROCRYSTALLI	NE CELLULOS	E (UNII: OP1R32D	D61U)					
PO	LYDEXTROSE (I	JNII: VH2XOU12	2IE)						
PO	LYETHYLENE G	LYCOL, UNSP	ECIFIED (UNII: 3	WJQ0SDW1A)					
TA	LC (UNII: 7SEV7J4	4R1U)							
тіт	ANIUM DIOXID	E (UNII: 15FIX9)	V2JP)						
TR	ACETIN (UNII: X	HX3C3X673)							
FE	RRIC OXIDE REI	D (UNII: 1K09F3	G675)						
FE	RRIC OXIDE YEI	LOW (UNII: E)	(438O2MRT)						
Pr	oduct Chara	acteristics							
	lor	PIN	к	Score				no score	
	ape		UND	Size				8mm	
	vor			Imprint Co	odo			L141	
	ntains			imprint co	Jue			LI41	
	iitaiiis								
Pa	ckaging								
#	ltem Code	Pa	ickage Descr	iption		Mark	eting Start Date		eting End Date
	NDC:49348-128-	30 in 1 CARTC				10/14/2	000		

M	larketing Marketing Category	Information Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
Μ	larketing	Information		
3		60 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:49348-128- 12	1 in 1 CARTON	09/10/2009	
2		90 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:49348-128- 13	1 in 1 CARTON	11/04/2009	

Revised: 7/2021

Strategic Sourcing Services LLC