

**MUCUS RELIEF MAXIMUM STRENGTH COLD FLU AND SORE THROAT-
acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid
Walgreens**

Drug Facts

Active ingredients (in each 20 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Phenylephrine HCL 10 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - sinus congestion and pressure
 - minor aches and pains
 - nasal congestion
 - headache
 - cough
 - sore throat
 - temporarily reduces fever
 - temporarily promotes nasal and/or sinus drainage
 - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product,

do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 day
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in a 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- dose as follows or as directed by a doctor
- mL = milliliter
- adults and children 12 years of age and older: 20 mL every 4 hours
- children under 12 years of age: do not use

Other information

- **each 20 mL contains:** sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, disodium EDTA, FD&C blue #1, FD&C yellow #10, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Principal Display Panel

Compare to Maximum Strength Mucinex® Fast-Max® Clear & Cool® Cold, Flu, & Sore Throat active ingredients††

Cold, Flu & Sore Throat

ACETAMINOPHEN 650 mg PAIN RELIEVER & FEVER REDUCER

Dextromethorphan HBr 20 mg / Cough Suppressant

GUAIFENESIN 400 mg / EXPECTORANT

PHENYLEPHRINE HCL 10 mg / NASAL DECONGESTANT

MAXIMUM STRENGTH

- Relieves headache & fever
- Controls cough
- Relieves nasal & chest congestion
- Thins & loosens mucus
- 12 years & older

MINT FLAVOR

FL OZ (mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

††This product is no manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Clear & Cool® Cold, Flu & Sore Throat.

DISTRIBUTED BY: WALGREEN CO.

200WILMOT RD., DEERFIELD, IL 60015

walgreens.com

Package Label

Walgreens

Compare to Maximum Strength Mucinex®
Fast-Max® Clear & Cool® Cold,
Flu & Sore Throat active ingredients!†

NDC 0363-0782-06

Cold, Flu & Sore Throat

ACETAMINOPHEN 650 mg /
PAIN RELIEVER & FEVER REDUCER
DEXTROMETHORPHAN HBr 20 mg /
COUGH SUPPRESSANT
GUAIFENESIN 400 mg / EXPECTORANT
PHENYLEPHRINE HCl 10 mg / NASAL DECONGESTANT

MAXIMUM STRENGTH

- Relieves headache & fever
- Controls cough
- Relieves nasal & chest congestion
- Thins & loosens mucus
- 12 years & older

MINT FLAVOR

6 FL OZ (177 mL)

PLD-C452B
LB006897

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2019 Walgreen Co.
ITEM 984979 W10139-0719-F



Walgreens Pharmacist Recommended
Walgreens Pharmacist Survey

†This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Clear & Cool® Cold, Flu & Sore Throat.

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

OR0719-F
PLD-C452B LB006896
PEEL CORNER FOR DRUG FACTS ▶

Drug Facts

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Guaifenesin 400 mg.....	Expectorant
Phenylephrine HCl 10 mg.....	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms
 - sinus congestion and pressure
 - minor aches and pains
 - headache
- temporarily reduces fever
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

Drug Facts (continued)

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease ■ heart disease ■ diabetes
- thyroid disease ■ high blood pressure

Drug Facts (continued)

- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for ▶

Drug Facts (continued)

adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years of age and older: 20 mL every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, disodium EDTA, FD&C blue #1, FD&C yellow #10, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

PEEL CORNER FOR MORE DRUG FACTS ▶

WALGREENS Cold Flu & Sore Throat

MUCUS RELIEF MAXIMUM STRENGTH COLD FLU AND SORE THROAT

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0782
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Acetaminophen	650 mg	

ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYL GALLATE (UNII: 8D45NN7V92)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0782-06	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/31/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/31/2017	

Labeler - Walgreens (008965063)

Revised: 10/2023

Walgreens