BASIC CARE OMEPRAZOLE- omeprazole tablet, orally disintegrating, delayed release

Amazon.com Services LLC

Amazon Omeprazole Drug Facts

Active ingredient (in each tablet)

Omeprazole 20 mg

Purpose

Acid reducer

Use

- treats frequent heartburn (occurs <u>2 or more</u> days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- take 1 tablet before eating in the morning
- do not crush or chew tablets
- place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- do not take this medicine with alcohol

Repeated 14-Day Course (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F); keep product out of high heat and moisture

Inactive ingredients

amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide,

crospovidone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize maltodextrin, mannitol, microcrystalline cellulose, propylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Questions or comments?

1-800-719-9260: weekdays 7:30 AM to 5:00 PM EST

Package/Label Principal Display Panel

amazon

basic care

Acid Reducer

Compare to Prilosec OTC®

Omeprazole

Delayed Release Orally Disintegrating Tablets 20 mg

Acid Reducer

Treats FREQUENT Heartburn!

24 HR

Melts in Your Mouth

Dissolves Without Water

actual size

Strawberry Flavor

14 TABLETS

One 14-day course of treatment

May take 1 to 4 days for full effect

MELTech™

Melts In Your Mouth

Tips for Managing Heartburn

Made in Israel

DISTRIBUTED BY: AMAZON.COM SERVICES LLC
410 TERRY AVENUE N., SEATTLE, WA 981 09

2020 AMAZON.COM, INC. OR ITS AFFILIATES.

ALL RIGHTS RESERVED

- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods.
- Eat slowly and avoid big meals
- If overweight, lose weight Quit smoking

Que stions or comments ? 1-800-719-9260: wækdays 7:30AM to

sodium stearyf fumarate, sorbitol, sucralose, sugar spheres, tale, titanium unicrocrystatine centrose, propylene glycol, silicon dioxide, sodium stearate, hypromellose, hypromellose phthalate, maize maltodextrin, mannifol, cetyl alcohol, colloidal alicon dioxide, crospovidone, ferric oxide, flavor, Inactive ingredients amino methacryate copolymer, ascorbic acid,

■ store at 20-25°C (68-77°F); keep product out of high heat and moisture keep the carton. It contains important information.

Other information = read the directions and warnings before use

children may sometimes be caused by a serious condition. ■ children under 18 years of age: ask a doctor before use. Heartburn in

4 months unless directed by a doctor ■ do not take for more than 14 days or more often than every ■ you may repeat a 14-day course every 4 months

Repeated 14-Day Course (if needed)

- do not take this medicine with alcohol
- do not use for more than 14 days unless directed by your doctor
 - do not take more than 1 tablet a day ■ take every day for 14 days
- The tablets can also be swallowed whole with water. ■ place the tablet on tongue; tablet disintegrates, with or without water.
 - a do not crush or chew tablets eating in the perore eating in the morning
 - 14-Day Course of Treatment
 - symptoms within 24 hours

■ it may take 1 to 4 days for full effect; some people get complete relief of ■ this product is to be used once a day (every 24 hours), every day for 14 days Directions Infor adults 18 years of age and older

Orug Facts (continued)

confact a Poison Control Center right away (1-800-222-1222). Keep out of reach of children. In case of overdose, get medical help or If pregnant or breast-feeding, ask a health professional before use.

- you develop a rash or joint pain
 - wouldet diarrhea
- you need to take more than 1 course of treatment every 4 months
 - you need to take this product for more than 14 days
 - Aont heartburn continues or worsens Stop use and ask a doctor if:

drug. Acid reducers may interact with certain prescription drugs. Ask a doctor or pharmacist before use if you are taking a prescription

- usness or vomiting stomach pain
 - ssol Jrigiaw banisiqxanu
- frequent wheezing, particularly with heartburn
- had heartburn over 3 months. This may be a sign of a more serious condition. Ask a doctor before use if you have:

These may be signs of a serious condition. See your doctor.

- Trequent chest pain
- to arms, neck or shoulders; or lightheadedness cyest or shoulder pain with shortness of breath; sweating; pain spreading
 - m heartburn with lightheadedness, sweating or dizziness
- trouble or pain swallowing food, vomiting with blood, or bloody or black stools Do not use if you have:

Amergy slert: do not use if you are allergic to omeprazole Marnings

days for full effect

■ not intended for immediate relief of heartburn; this drug may take 1 to 4 **OSC** Treats frequent heartburn (occurs 2 or more days a week)

Omeprazole 20 mg Acid reducer Active ingredient (in each tablet) əsodınd

Drug Facts



Safety Feature -Do not use if printed tablet blister unit is open or torn.

basic@care

Acid Reducer

NDC 72288-123-74

Compare to Prilosec OTC®

)meprazole

Delayed Release Orally Disintegrating Tablets 20 mg Acid Reducer

Treats FREQUENT Heartburn!

24 HR

Melts in Your Mouth Dissolves Without Water



actual size

MELTech™ Melts In Your Mouth

Strawberry

14 TABLETS

One 14-day course of treatment May take 1 to 4 days for full effect



BASIC CARE OMEPRAZOLE

omeprazole tablet, orally disintegrating, delayed release

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:72288-123

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OMEPRAZOLE (UNII: KG60484QX9) (OMEPRAZOLE - UNII:KG60484QX9)	OMEPRAZ OLE	20 mg

Inactive Ingredients

Ingredient Name	Strength
DIMETHYLAMINOETHYL METHACRYLATE - BUTYL METHACRYLATE - METHYL METHACRYLATE COPOLYMER (UNII: 905HNO1SIH)	
ASCORBIC ACID (UNII: PQ6CK8PD0R)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSPOVIDONE (15 MPA.S AT 5%) (UNII: 68401960MK)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MANNITOL (UNII: 30WL53L36A)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM STEARATE (UNII: QU7E2XA9TG)	
SODIUM STEARYL FUMARATE (UNII: 7CV7WJK4UI)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	RED (reddish)	Score	no score
Shape	ROUND	Size	9mm
Flavor	STRAWBERRY	Imprint Code	20
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:72288-123- 74	14 in 1 CARTON	08/19/2021		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product			
2	NDC:72288-123- 55	42 in 1 CARTON	04/11/2022		
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA209400	08/19/2021	

Labeler - Amazon.com Services LLC (128990418)

Revised: 4/2022 Amazon.com Services LLC