

CHILDRENS PAIN AND FEVER- acetaminophen tablet, chewable
Topco Associates, LLC

Topcare 44-450

Active ingredient (in each chewable tablet)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - sore throat
 - flu
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if your child has

liver disease.

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- any new symptoms appear

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (chewable tablets)*
Under 24	Under 2	Ask a doctor
24-35	2-3	1 tablet
36-47	4-5	1 1/2 tablets
48-59	6-8	2 tablets
60-71	9-10	2 1/2 tablets
72-95	11	3 tablets

*or as directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid high humidity
- see end flap for expiration date and lot number

Inactive ingredients

citric acid, crospovidone, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrans hydrated, ethylcellulose, flavor, magnesium stearate, mannitol, polyethylene, stearic acid, sucralose

Questions or comments?

1-888-423-0139

Principal display panel

+TopCare®
health

NDC 36800-450-08

**†COMPARE TO THE ACTIVE INGREDIENT IN
CHILDREN'S TYLENOL® CHEWABLES**

children's
Pain & Fever

ACETAMINOPHEN 160 mg,
CHEWABLE TABLETS
PAIN RELIEVER • FEVER REDUCER

• Ibuprofen Free • Aspirin Free

Chew or crush tablets completely before swallowing.

BUBBLE
GUM
FLAVOR

24 CHEWABLE TABLETS

For Ages 2 to 11 Years

actual size

**TAMPER EVIDENT: DO NOT USE IF
IMPRINTED SAFETY SEAL UNDER
CAP IS BROKEN OR MISSING**

QUALITY GUARANTEED

† This product is not manufactured or distributed by
Kenvue Inc., owner of the registered trademark
Children's Tylenol® Chewables. 50844 REV0424C45008

DISTRIBUTED BY TOPCO ASSOCIATES LLC, ITASCA, IL 60143

©TOPCO LNKA1024 QUESTIONS? 1-888-423-0139

topcare@topco.com

www.topcarebrand.com



KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts
Active ingredient (in each chewable tablet)
 Acetaminophen 160 mg..... Pain reliever/fever reducer

Uses
 ■ temporarily relieves minor aches and pains due to:
 ■ headache ■ sore throat ■ the common cold
 ■ toothache ■ flu
 ■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes
 ■ more than 5 doses in 24 hours, which is the maximum daily amount
 ■ with other drugs containing acetaminophen
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ rash ■ skin redness ■ blisters
 If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
 Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
Ask a doctor before use if your child has liver disease.

No Print/No Varnish
 Lot & EXP

0 3680045569 3

QUALITY GUARANTEED

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

TopCare
 health

NDC 36800-450-08

'COMPARE TO THE ACTIVE INGREDIENT IN CHILDREN'S TYLENOL® CHEWABLES

TopCare
 health

children's
Pain & Fever
ACETAMINOPHEN 160 mg,
CHEWABLE TABLETS
 PAIN RELIEVER • FEVER REDUCER
 • Ibuprofen Free • Aspirin Free
 Chew or crush tablets completely before swallowing.



children's
Pain & Fever
ACETAMINOPHEN 160 mg,
CHEWABLE TABLETS
 PAIN RELIEVER • FEVER REDUCER

24 CHEWABLE TABLETS For Ages 2 to 11 Years actual size

24 CHEWABLE TABLETS

Drug Facts (continued)
Other information
 ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
 ■ avoid high humidity
 ■ see end flap for expiration date and lot number

Inactive ingredients
 citric acid, croscollonite, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrates hydrated, ethylcellulose, flavor, magnesium stearate, mannitol, polyethylene, stearic acid, sucralose

Questions or comments? 1-888-423-0139

* This product is not manufactured or distributed by Kenvue Inc., owner of the registered trademark Children's Tylenol® Chewables.
 50844 REV0424C45008

DISTRIBUTED BY TOPCO ASSOCIATES LLC, ITASCA, IL 60143
 ©TOPCO LWA1024 QUESTIONS? 1-888-423-0139
 topcare@topco.com
 www.topcarebrand.com



Scan here for more information or call 1-888-423-0139

B-1910-450-08-R
 REV10424C45008

Drug Facts (continued)
Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.
Stop use and ask a doctor if
 ■ pain gets worse or lasts more than 5 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ any new symptoms appear
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
 ■ this product does not contain directions or complete warnings for adult use
 ■ do not give more than directed
 ■ find the right dose on chart below. If possible, use weight to dose; otherwise, use age.
 ■ chew or crush tablets completely before swallowing; do not swallow tablets whole
 ■ repeat dose every 4 hours while symptoms last
 ■ do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (chewable tablets)*
Under 24	Under 2	Ask a doctor
24-35	2-3	1 tablet
36-47	4-5	1 1/2 tablets
48-59	6-8	2 tablets
60-71	9-10	2 1/2 tablets
72-95	11	3 tablets

*or as directed by a doctor

TopCare 44-450

CHILDRENS PAIN AND FEVER

acetaminophen tablet, chewable

Product Information		
Product Type	HUMAN OTC DRUG	Item Code (Source)
Route of Administration	ORAL	NDC:36800-450

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
CROSPROVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)	
D&C RED NO. 30 ALUMINUM LAKE (UNII: GE75M6YV5W)	
DEXTROSE MONOHYDRATE (UNII: LX22YL083G)	
ETHYLCELLULOSE, UNSPECIFIED (UNII: 7Z8S9VYZ4B)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 3OWL53L36A)	
HIGH DENSITY POLYETHYLENE (UNII: UG00KM4WR7)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	pink	Score	2 pieces
Shape	ROUND	Size	16mm
Flavor	BUBBLE GUM	Imprint Code	44;450
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-450-08	1 in 1 CARTON	02/15/2019	
1		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/15/2019	

Labeler - Topco Associates, LLC (006935977)**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(36800-450)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(36800-450) , pack(36800-450)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(36800-450)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(36800-450)

Revised: 3/2025

Topco Associates, LLC