

ASPIRIN- aspirin tablet, coated
ARMY AND AIR FORCE EXCHANGE SERVICE

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Exchange Select 44-157

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves:

- minor pain of arthritis
- headache
- muscle pain
- menstrual pain
- toothache
- pain and fever of colds

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- asthma (wheezing)
- hives
- facial swelling
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you are taking a diuretic
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients

corn starch, hypromellose, polyethylene glycol, propylene glycol

Questions or comments?

1-800-426-9391

Principal Display Panel

**Compare To The Active Ingredient of
Genuine Bayer® Aspirin†**

exchange select™

ASPIRIN

Pain Reliever/Fever Reducer
(NSAID)

325 mg

Actual Size

Talk to your doctor or other healthcare provider before using this product for your heart.

100 COATED TABLETS

quality

value

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by Bayer AG,
owner of the registered trademark Genuine Bayer® Aspirin.

"SATISFACTION GUARANTEED OR YOUR MONEY BACK"

Manufactured For Your Military Exchanges

Distributed by: LNK International, Inc.,

Hauppauge, NY 11788

1-800-426-9391

50844 REV0318D15712

Compare To The Active Ingredient of
Genuine Bayer® Aspirin[†]

exchange **select**[™]

ASPIRIN

Pain Reliever/Fever Reducer
(NSAID)

325 mg
Actual Size

Talk to your doctor or other healthcare provider before using this product for your heart.

100 COATED TABLETS quality value

50844 RE10318015712

5
1 4 2 9 9 3 9 8 6 6
6

This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Genuine Bayer® Aspirin.

"SATISFACTION GUARANTEED OR YOUR MONEY BACK"
Manufactured For Your Military Exchanges
Distributed by LNK International, Inc.
Hauppauge, NY 11788 1-800-426-9391

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet)
Aspirin 325 mg (NSAID)*Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses temporarily relieves: ■ minor pain of arthritis ■ headache ■ muscle pain ■ menstrual pain ■ toothache ■ pain and fever of colds

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ facial swelling ■ shock ■ hives ■ asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ take more or for a longer time than directed

No print/No varnish
Lot & Exp date

PEEL HERE FOR MORE DRUG FACTS

Adhesive Only

STOP PEELING

Drug Facts (continued)

■ have had stomach ulcers or bleeding problems
■ have 3 or more alcoholic drinks every day while using this product

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ you are taking a diuretic ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ ringing in the ears or a loss of hearing occurs
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present ■ new symptoms occur
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ drink a full glass of water with each dose
■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
■ children under 12 years: do not use unless directed by a doctor

Other information ■ use by expiration date on package ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients corn starch, hypromellose, polyethylene glycol, propylene glycol

Questions or comments? 1-800-426-9391

Exchange Select 44-157

ASPIRIN

aspirin tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55301-157
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Product Characteristics

Color	WHITE	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	ASPIRIN;44;157
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55301-157-12	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/10/1996	
2	NDC:55301-157-14	500 in 1 BOTTLE; Type 0: Not a Combination Product	02/10/1996	
3	NDC:55301-157-17	300 in 1 BOTTLE; Type 0: Not a Combination Product	02/10/1996	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part343	02/10/1996	

Labeler - ARMY AND AIR FORCE EXCHANGE SERVICE (001695568)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(55301-157)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	MANUFACTURE(55301-157)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	PACK(55301-157)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	PACK(55301-157)