

EXTRA STRENGTH PAIN RELIEVER- acetaminophen, aspirin, and caffeine capsule, coated
JHK Inc dba American Safety & First Aid

Extra Strength Pain Reliever

Drug Facts

<i>Active ingredient (In Each Caplets)</i>	<i>Purpose</i>
Acetaminophen 250 mg	pain reliever/fever reducer
Aspirin 250 mg (NSAID)*	pain reliever/fever reducer
Caffeine 65 mg	pain reliever aid

* Nonsteroidal Anti-Inflammatory Drug

Use

temporarily relieves fever and minor aches and pains associated with:

- headache
- colds
- muscular aches
- menstrual discomfort
- toothaches
- minor arthritis pain

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness.

Allergy Alert

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Liver Warning

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning

This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing a prescription or nonprescription NSAID (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning

The recommended dosage of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness and, occasionally, rapid heartbeat.

Do not use

- if you are allergic to aspirin
- with any other pain reliever/fever reducer
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for pain for more than 10 days or for fever for more than 3 days unless directed by a doctor
- with any other product containing acetaminophen (prescription or nonprescription).
- If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask doctor before use if

- stomach bleeding warning applies to you
- you have liver disease, high blood pressure, heart disease, liver cirrhosis, asthma or kidney disease
- you are taking diuretic
- you have a history of stomach problems, such as heartburn

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- anticoagulation (thinning of blood)
- diabetes
- gout
- arthritis

When using this product do not exceed recommended dose.

Stop use and ask doctor if you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- pain or fever persists or gets worse
- New symptoms occur
- Redness or swelling is present

If pregnant or breastfeeding, ask a health professional before use. IT IS ESPECIALLY IMPORTANT NOT TO USE ASPIRIN DURING THE LAST 3 MONTHS OF PREGNANCY UNLESS SPECIFICALLY DIRECTED TO DO SO BY A DOCTOR BECAUSE IT MAY CAUSE PROBLEMS IN THE UNBORN CHILD OR COMPLICATIONS DURING DELIVERY.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions Adults

2 tablets, with water every 6 hours while symptoms persist; not to exceed 8 tablets in 24 hours or as directed by a doctor. Children under 12 years: consult a doctor.

Other Information

- do not use if packet is opened or torn
- store at room temperature 68°-77°F (20°-25°C)
- avoid excessive heat and humidity
- each caplet contains: Magnesium 0.58 mg

Inactive ingredient

Colloidal Silicon Dioxide, Hypromellose, Microcrystalline Cellulose, Polyvinyl Pyrrolidone, Polyethylene Glycol, Pregelatinized Starch, Starch 1500, Stearic Acid, Talc

Questions?

1-866-651-3660 Monday-Friday

8:00am-5:00pm EST

Manufactured for American Safety & First Aid Osceola, IN 46561

PRINCIPAL DISPLAY PANEL - 50 Caplet Packet Box

This product is not manufactured or distributed by Novartis
Owner of the registered trademark Excedrin®.

Product #1372

Extra Strength
Pain Reliever

Pain Reliever/Fever Reducer

Active ingredient:

Acetaminophen, Aspirin

Caffeine

- Temporary relief from pain, fever, and inflammation
- Soothes muscle aches and minor arthritis pain

These quality comfort caplets are formulated and packaged for use in the workplace and meet ANSI Z308. 1-2021 standards. They are sealed in tamper-evident packets for safety and convenience.

50 Caplets • 2 Caplets Per Packet

This product is not manufactured or distributed by Novartis
Owner of the registered trademark Excedrin®.

Product #1372



Extra Strength Pain Reliever

Active ingredient:
Acetaminophen, Aspirin
Caffeine

Pain Reliever/Fever Reducer

- Temporary relief from pain, fever, and inflammation
- Soothes muscle aches and minor arthritis pain

These quality comfort caplets are formulated and packaged for use in the workplace and meet ANSI Z308.1-2021 standards. They are sealed in tamper-evident packets for safety and convenience.

50 Caplets • 2 Caplets Per Packet

Drug Facts (continued)

• anticoagulation (thinning of blood) • diabetes • gout • arthritis

When using this product do not exceed recommended dose.

Stop use and ask doctor if you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • ringing in the ears or loss of hearing occurs • pain or fever persists or gets worse • New symptoms occur • Fizziness or swelling is present

If pregnant or breastfeeding, ask a health professional before use. IT IS ESPECIALLY IMPORTANT TO USE ASPIRIN DURING THE LAST 3 MONTHS OF PREGNANCY UNLESS SPECIFICALLY DIRECTED TO DO SO BY A DOCTOR BECAUSE IT MAY CAUSE PROBLEMS IN THE UNBORN CHILD OR COMPLICATIONS DURING DELIVERY. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions Adults: 2 tablets, with water every 6 hours while symptoms persist, not to exceed 8 tablets in 24 hours or as directed by a doctor. Children under 12 years: consult a doctor.

Other Information • do not use if packet is opened or torn • store at room temperature 68°-77°F (20°-25°C) • avoid excessive heat and humidity • each caplet contains: Magnesium 0.58 mg

Inactive Ingredient Colloidal Silicon Dioxide, Hypromellose, Microcrystalline Cellulose, Polyvinyl Pyrrolidone, Polyethylene Glycol, Prugelatinized Starch, Starch 1500, Stearic Acid, Talc

Questions? 1-866-651-3660 Monday-Friday
8:00am-5:00pm EST
Manufactured for American Safety & First Aid Osceola, IN 46061

Pain Reliever/Fever Reducer

Extra Strength Pain Reliever



Product #1372

50 Caplets • 2 Caplets Per Packet

Drug Facts

Active Ingredient (In Each Caplet)	Purpose
Acetaminophen 250 mg	pain reliever/fever reducer
Aspirin 250 mg (NSAID)*	pain reliever/fever reducer
Caffeine 65 mg	pain reliever aid

*Nonsteroidal Anti-Inflammatory Drug

Use temporarily relieves fever and minor aches and pains associated with: • headache • colds • muscular aches • menstrual discomfort • toothaches • minor arthritis pain


Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness. **Allergy Alert:** Aspirin may cause a severe allergic reaction which may include: hives • facial swelling • asthma (wheezing) • stomach Liver Warning: This product contains acetaminophen. Severe liver damage may occur if you take: • more than 8 tablets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs



Extra Strength Pain Reliever

Pain Reliever/Fever Reducer



EXTRA STRENGTH PAIN RELIEVER

acetaminophen, aspirin, and caffeine capsule, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73598-1372
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

TALC (UNII: 7SEV7J4R1U)

Product Characteristics

Color	WHITE	Score	no score
Shape	RECTANGLE	Size	18mm
Flavor		Imprint Code	S53
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73598-1372-1	25 in 1 BOX	02/02/2000	
1		2 in 1 DOSE PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph drug	M013	02/02/2000	

Labeler - JHK Inc dba American Safety & First Aid (867236309)

Revised: 4/2024

JHK Inc dba American Safety & First Aid