METHYL SALICYLATE- methyl salicylate cream PHARMACURE LLC

Active Ingredient **Purpose**

Methyl Salicylate 10% (NSAID: nonsteroidal anti-inflammatory drug) Analgesic

Topical

Uses

Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:

• strains • sprains • simple backache • arthritis • bruises

Warnings

For external use only

Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding.

The chance is small but higher if you:

- are age 60 or older have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- on the face or rashes on wounds or damaged skin if allergic to aspirin or other **NSAIDs**
- with a heating pad when sweating (such as from exercise or heat)
- right before or after heart surgery

Ask a doctor before use if

- you are allergic to topical products the stomach bleeding warning applies to you
- you have high blood pressure, heart disease, or kidney disease you are taking a diuretic

When using this product

- wash hands after applying or removing cream. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water.
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

• stomach pain or upset gets worse or lasts • you feel faint, vomit blood, or have bloody or black stools

These are signs of stomach bleeding. • rash, itching or skin irritation develops

- condition worsens
- symptoms last for more than 3 days symptoms clear up and occur again within a few days

If pregnant or breast-feeding,

ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant.

Directions

- Use only as directed
- Adults and children 12 years of age and older: Apply to affected area not more than 3 to 4 times daily
- children under 12 years of age: consult a physician

Other information

- Store at 20-25°C (68-77°F) Avoid storing product in direct sunlight
- Protect product from excessive moisture

Inactive ingredients

Aqua (deionized water), arnica montana (arnica) extract, boswellia serrata extract, cetearyl olivate, cetyl alcohol, dimethyl sulfone (MSM), ethylhexylglycerin, glycerin, glyceryl stearate, hemp, sodium laurylglucosides hydroxypropylsulfonate, PEG-100 stearate, phenoxyethanol, polysorbate-20, resveratrol, sodium laurylglucosides hydroxypropylsulfonate, sorbitan olivate, stearic acid

Product label





CONTAINS

- ARNICA
- BOSWELLIA
- MSM
- RESVERATROL

NDC: 79643-009-01





CYLATE CREAM

DRUG FACTS

ACTIVE INGREDIENTS: Purpose
Methyl Salicylate 10% (NSAID: nonsteroidal anti-inflammatory drug).....Topical Analges

Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:

*strains *sprains *simple backache *arthrifis *bruises

- · Use only as directed · Adults and children 12 years of age and older: apply to affected
- area not more than 3 to 4 times daily

 children under 12 years of age: consult a physician

For external use only
Stomach bleeding warning: This product contains an NSAID, which may
cause stomach bleeding.

cause stomach bleeding.

The chance is mall but higher if you:

*are age 60 or older * have had stomach ulcens or bleeding problems

*ake a blood thirming fantiosqualidny of setend drug

*ake other drugs containing an NSAID (saprim, buprofen, naprosen, or others)

*ake one acchoric drivisk every day while using this product

*ake more or for a longer time than directed

Do not use

- Table Thore on the disregard with the research of the face or rathers on wounds or damaged skin if allergic to aspirin or other NSAIDs with a heating pad in when sweating (such as from exercise or heat) right before or after heart surgery
- Ask a doctor before use if
- you are allergic to topical products the stomach bleeding warning applies to you you have high blood pressure, heart disease, or kidney disease you are taking a diuretic

- you not be right blood pressure, near unescale, or hothey disease
 "you are taking a diserlect
 When using this product
 "when hands there applying or removing cream. Avoid contact with eyes.
 If eye contact occurs, mee theroughly with water.
 If eye contact occurs, mee theroughly with water,
 the risk of heart attack or stoke may increase if you use more than
 directed or for longer than directed
 Stop use and ask a doctor If
 "attorned pann or upset gets worse or lasts "you feel faint, vomit blood,
 or have bloody or block stoods. These are signs of stomach bleeding.
 "rash, fathing or slin initiation develops "condition worsens
 "symptoms last for more than 3 days
 "symptoms clear no met han 3 days
 "symptoms clear no met han 3 days
 "symptoms clear for more than 3 days
 The great may be reast-feedling,
 ask a doctor before use while breast-feeding and during the first 6 months of pregnancy.
 Do not use during the last 3" months of pregnancy because it may cause problems in the
 unborn child or complications during delivery. Keep out of reach of children. If
 If put in month, get medical help or contact a Poison Control Center right away.
 Package not child resistant.

- Store at 20-25°C (68-77°F) · Avoid storing product in direct sunlight Protect product from excessive moisture

IRINAL IVE INVICEURENTS:
Aqua (deionized water), amicia montana (amica) extract, bosvellia serrata extract, ceteary olivate, cetyl alcohol, dimethyl sulfone (MSM), ethylhexylgycerin, glycerin, gly



Manufactured for: PHARMACURE

For questions or comments



methyl salicylate cream

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79643-009
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
METHYL SALICYLATE (UNII: LAV5U5022Y) (SALICYLIC ACID - UNII: 0414PZ4LPZ)	METHYL SALICYLATE	100 mg in 1 g	

Inactive Ingredients		
	Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)		
ARNICA MONTANA FLOWER (U	NII: OZ0E5Y15PZ)	
BOSWELLIA SERRATA WHOLE	(UNII: X7B7P649WQ)	
CETEARYL OLIVATE (UNII: 5886	9Q84JO)	
CETYL ALCOHOL (UNII: 936JST6	JCN)	
DIMETHYL SULFONE (UNII: 9H4	PO4Z4FT)	
ETHYLHEXYLGLYCERIN (UNII: 1	47D247K3P)	
GLYCERIN (UNII: PDC6A3C0OX)		
GLYCERYL MONOSTEARATE (U	NII: 2300U9XXE4)	
HEMP (UNII: TD1MUT01Q7)		
PEG-100 STEARATE (UNII: YD0	.N1999R)	
PHENOXYETHANOL (UNII: HIE49	92ZZ3T)	
POLYSORBATE 20 (UNII: 7T1F3	OV5YH)	
RESVERATROL (UNII: Q3690892	6L)	
SODIUM LAURYLGLUCOSIDES	HYDROXYPROPYLSULFONATE (UNII: Z6GFR7R72Y)	
SORBITAN OLIVATE (UNII: MDL2		
STEARIC ACID (UNII: 4ELV7Z65A	AP)	

F	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79643-009- 01	1 in 1 BOX	02/01/2025	
1		100 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	02/01/2025	

Labeler - PHARMACURE LLC (055983858)

Revised: 2/2025 PHARMACURE LLC