

**4146 FIRST AID KIT- 4146 first aid
Honeywell Safety Products USA, Inc.**

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

0498-4146: First Aid Kit (Burn Jel, BZK wipe, alcohol wipe, sting relief-SF00004042)

**Burn Jel
Active ingredient**

Lidocaine HCl 2.0%

**Burn Jel
Purpose**

External analgesic

**Burn Jel
Uses**

- temporarily relieves pain due to minor burns

**Burn Jel
Warnings**

For external use only

Do not use

- on large areas of the body, particularly over raw surfaces or blistered areas

When using this product

- avoid contact with eyes

Stop use and ask a doctor if

- the condition gets worse
- symptoms persist for more than 7 days
- condition clears up and recurs within a few days

Keep out of reach of children

- If swallowed, get medical help or contact a Poison Control Center right away.

**Burn JEl
Directions**

- adults and children 2 years of age and older; apply to affected area not more than 3

to 4 times daily

- children under 2 years of age: ask a doctor
- you may report a serious reaction to this product to 800-430-5490

Burn Jel

Other information

- store at room temperature - do not use if opened or torn

Burn Jel

Inactive ingredients

carbopol 940, carbopol 1342, diazolidinyl urea, glycerin, melaleuca alternifolia (tea tree) leaf oil, methylparaben, octoxynol-9, propylene glycol, propylparaben, trolamine, water

Burn Jel

Questions

1-800-430-5490

BZK Wipe

Active ingredient

Benzalkonium chloride 0.13% w/v

BZK Wipe

Purpose

First aid antiseptic

BzK Wipe

Uses

Antiseptic cleansing of face, hands, and body without soap and water

BZK Wipe

Warnings

For external use only

Do not use

- in the eyes or over large areas of the body
- on mucous membranes
- on irritated skin
- in case of deep puncture wounds, animal bites or serious burns, consult a doctor
- longer than 1 week unless directed by a doctor

Stop use and ask a doctor if

- if irritation, redness or other symptoms develop
- the condition persists or gets worse

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

BZK Wipe***Directions***

tear open packet and use as a washcloth

BZK Wipe***Other information***

- store at room temperature 15⁰ to 30⁰ C (59⁰ - 86⁰ F)
- do not reuse towelette

BZK Wipe***Inactive ingredient***

water

BZK Wipe***Questions***

1-800-430-5490

Sting Relief***Active ingredient (in each wipe)***

Ethyl alcohol 50.0%

Lidocaine HCl 2.0%

Sting Relief***Purpose***

Antiseptic

Topical pain relief

Sting Relief***Uses***

- prevent infection in minor scrapes, and temporary relief of itching of insect bites

Sting Relief

Warnings

For external use only

Flammable, keep away from open fire or flame

Do not use

- over large areas of the body
- in eyes
- over raw or blistered areas

Stop use and ask a doctor

- if conditions worsen or persist for more than 7 days or clear up and occur again within a few days

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Sting Relief

Directions

- adults and children 2 years and older: Apply to cleaned affected area not more than 3 times daily.
- children under 2 years of age: consult a doctor.

Sting Relief

Inactive ingredients

benzalkonium chloride, menthol, and purified water

Sting Relief

Questions or Comments?

1-800-430-5490

Alcohol

Active ingredient

Isopropyl alcohol 70%

Alcohol

Purpose

First aid antiseptic

Alcohol

Uses

- first aid to help prevent infection in minor cuts, scrapes, and burns

Alcohol

Warnings

For external use only

Flammable, keep away from fire and flame

Do not use

- in or near eyes
- over large areas of the body

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

When using this product

- do not use longer than 1 week unless directed by a doctor

Stop use and consult a doctor if

- condition persists or gets worse

Keep out of the reach of children

If swallowed, get medical help or contact a Poison Control center right away

Alcohol

Directions

- clean the affected area
- may be covered with a sterile bandage
- apply wipe to affected area 1 to 3 times daily
- discard wipe after single use

Alcohol

Other information

- store at room temperature 15 ° to 25 ° C (59 ° to 77 ° F)
- do not use if packet is torn or

**Alcohol
Questions**


1-800-430-5490

**4146
SF00004042 kit contents**



1 KNUCKLE BAND 8 PER
1 TRIANGULAR BDG, NON-STERILE
1 GAUZE PADS, 3" X 3", 4 PER
1 ADH TAPE, .5" X 2.5 YD, 2 PER
1 GAUZE COMPRESS, 1728 SQ IN 1
1 FORCEPS & SCISSORS, 1 EA
1 INSTANT COLD PACK 4" X 6"
1 ADHESIVE BDG, PLSTIC, 1" X 3" 16 PER
1 BURN JEL 1/8 OZ, 6 PER
1 NITRILE GLOVES 2PR BBP
1 ANTIMCRBL ANTSPTC TWLETTS
1 MICROSHIELD W/VNL GLV/ALCL
1 F. A. INST CHART SM (INDIVIDUAL LBL)
1 BANDAGE COMP 4" W/TELFAPAD 1
LBL STOCK 6-3/8" X 4"
LBL STOCK 4" X 2-7/8"
1 LBL STOCK 3" X 1-7/8"
1 KIT STL 16 UN (VERTICAL)
1 STING Relief SWAB 10

**Burn Jel
Principal Display Panel**


796353 Rev. E Unit Carton Printing Plate for "B" size cartor



Principal Display Panel



Sting Relief Principal Display Panel



825366 Rev B

Honeywell

032043P

Sting Relief Wipes

Ethyl alcohol
First aid antiseptic
Lidocaine HCl
Topical analgesic

Use for:
Minor Cuts • Scrapes • Insect Bites

Single Use Pouches
Saturated Wipes

100 wipes

Honeywell

032043P

Sting Relief Wipes

Ethyl alcohol
First aid antiseptic
Lidocaine HCl
Topical analgesic

Use for:
Minor Cuts • Scrapes • Insect Bites

Single Use Pouches
Saturated Wipes

100 wipes

Drug Facts

Active Ingredients
Ethyl alcohol 50.0%
Lidocaine HCl 2.0%

Purpose
First aid antiseptic
Topical analgesic

Uses first aid to help prevent infection in minor scrapes and temporary relief of itching of insect bites.

Warnings
For external use only.
Flammable, keep away from fire or flame.

Do not use • over large areas of the body • in eyes • over raw or blistered areas

Stop use and ask a doctor • If conditions worsen or persist for more than 7 days

Keep out of reach of children.
If swallowed, get medical help or contact Poison Control Center right away.

Directions
• adults and children 2 years and older: apply to cleaned affected area not more than 3 times daily.
• children under 2 years of age: consult a doctor.

Inactive Ingredients
Benzalkonium chloride, menthol, purified water

Questions or comments? 1-800-430-5490

Distributed by
Honeywell Safety
Products, USA, Inc.
Smithfield, RI 02917


USA
www.honeywellsafety.com

1-800-430-5490
0-401-943-4400



Made in USA
Packaged in Mexico

Alcohol Principal Display Panel



4146 Kit Label
SF00004042





CITY OF SEATTLE

FIRST AID KIT

Distributed by Honeywell Safety Products USA, Inc. Smithfield, RI 02917

4146 FIRST AID KIT

4146 first aid kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0498-4146
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-4146-01	1 in 1 KIT	09/13/2018	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	6 PACKET	21 g

Part 2	2 PACKET	2.8 mL
Part 3	4 POUCH	1.6 mL
Part 4	10 POUCH	4 mL

Part 1 of 4

BURN JEL

gel for bums gel

Product Information

Item Code (Source) NDC:0498-0203

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE ANHYDROUS	2 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
TEA TREE OIL (UNII: VIF565UC2G)	
DIAZOLIDINYL UREA (UNII: H5RIZ3MPW4)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
GLYCERIN (UNII: PDC6A3C0OX)	
TROLAMINE (UNII: 9O3K93S3TK)	
CARBOMER HOMOPOLYMER TYPE C (ALLYL PENTAERYTHRITOL CROSSLINKED) (UNII: 4Q93RCW27E)	
CARBOMER HOMOPOLYMER TYPE B (ALLYL PENTAERYTHRITOL CROSSLINKED) (UNII: HHT01Z NK31)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
OCTOXYNOL-9 (UNII: 7JPC6Y25QS)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0203-00	3.5 g in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
unapproved drug other		09/19/2018	

Part 2 of 4

ANTISEPTIC TOWELETTE

benzalkonium chloride liquid

Product Information

Item Code (Source) NDC:0498-0501

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII: 7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	1.3 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KOOR)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0501-00	1.4 mL in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/22/2017	

Part 3 of 4

ALCOHOL WIPE

isopropyl alcohol swab

Product Information**Item Code (Source)** NDC:0498-0143**Route of Administration** TOPICAL**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	0.7 mL in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0143-04	0.4 mL in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	

Part 4 of 4**STING RELIEF PAD**

ethyl alcohol, lidocaine swab

Product Information**Item Code (Source)** NDC:0498-0733**Route of Administration** TOPICAL**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE ANHYDROUS	20 mg in 1 mL
ALCOHOL (UNII: 3K9958V90M) (ALCOHOL - UNII:3K9958V90M)	ALCOHOL	0.5 mL in 1 mL

Inactive Ingredients

Ingredient Name	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	
MENTHOL (UNII: L7T10EIP3A)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0733-00	0.4 mL in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/23/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/13/2018	

Labeler - Honeywell Safety Products USA, Inc. (118768815)

Revised: 1/2024

Honeywell Safety Products USA, Inc.