DEXTROMETHORPHAN HYDROBROMIDE AND GUAIFENESINdextromethorphan hydrobromide and guaifenesin syrup KJD Pharma Private Limited

Cough & Chest CongestionDextromethorphan Hbr - Guaifenesin CHILDREN Syrup 118 mL, 236 mL and 473 mL and measuring cup

Active Ingredient (in each 5 mL) Purpose

Dextromethorphan HBr USP 10 mg.	Cough suppressant
Guaifenesin, USP 100 mg	Expectorant

Relieves:

- Chest congestion / Mucus
- Cough

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps to loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

• **Do not use**if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor ifcough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feedingask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a

Poison Control Center right away.

Directions

- take evey 4 hours
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product

Age	Dose
Adults and children 12 years and over	2 teaspoonful (10 mL)
Children 6 to under 12 years	1 teaspoonful (5 mL)
Children under 6 years	Ask a doctor

Other information

- Store at 20°-25°C (68° -77°F).
- Do not refrigerate.
- Retain carton for future reference on full labeling

Inactive ingredients

citric acid, flavor, glycerin, menthol, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium saccharin, sorbitol.

Manufactured by

KJD Phanna Pvt lid, 148, 3 Kiadb, Naubad, Bidar, Kamataka 585401 , India

NDC 85722-0001-4 Cough & Chest Congestion Children Syrup Dextromethorphan Hbr - Guaifenesin	lamper evident: Do not use if inner seal under cap is broken or missing.	Purpose Cough suppressant Exnectorant	Martings Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor or pharmacist before taking this product. Ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have a cough that loccurs with two much phegm (mucus) = cough that losts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.	Take every 4 hours = do not take more than 6 doses in any 24-hour period take every 4 hours = do not take more than 6 doses in any 24-hour period the measure only with dosing cup provided the keep dosing cup with product Add the measure only with dosing and over 2 teaspoonful (10 m) Add the and children 12 years and over 2 teaspoonful (5 m) Children under 6 years Advector Ask a doctor	thylene glycol, propylene glycol, saccharin, sorbitol		T HERE BATCH CODE
Helps relieve cough Promotes phlegm expulsion Helps to decongest the respiratory tract	ot use if inner seal u	L LABELLING 1 each 5 ml) USP 10 mg	not use if you are now taking a prescr (MAOI) (certain drugs for depression ff you do not know if your prescription pharmacist before taking this product efore use if you have cours with too much phlegm (mucus) sts or is chronic such as occurs with si its or emphysema act of the er, rash, or persistent headac us condition. Is condition act of children, ask a health profession act of children in case of overdose in Control Center right away.	do not take more thar sing cup provided 2 years and over 5 s	menthol, polye zoate, sodium	.B, 3 Kiadb, ka 585401, India	DO NOT PRIN' NON-VARNISH AREA FOR & EXP DATE
Administration way: Oral Boxed bottle of 4 fl oz 118 mL and measuring cup	Tamper evident: Do n	SEE CARTON FOR FULL LABELLIN Active ingredients (in each 5 ml) Dextromethorphan HBr USP 10 mg Guaitenesin USP 100 mr	Warmings Do not use if you are now taking a presci oxidase inhibitor (MAOI) (certain drugs for depression emotional conditions, or Parkinson's disease), or for the MAOI drug. If you do not know if your prescription ask a doctor before use if you have – cough that lasts or is chronic such as occurs with a cough that lasts or is chronic such as occurs with chronic bronchitis or emphysema fromic bronchitis or emphysema Stop use and ask a doctor if ough lasts more that or is accompanied by fever, rash, or persistent heada signs of a serious condition. If pregnant or breaset feeding, ask a health professi Keep out of reach of children. In case of overdose contact a Poison Control Center right away. Directions	■ take every 4 hours ■ do not take more	Inactive ingredients citric acid, flavor, glycerin, ment purified water, sodium benzoate,	Manufactured by KJD Pharma Pvt Ltd, 14B, Naubad, Bidar, Karmataka	Exp NON-VARNISH /

NDC 85722-0001-8 Cough & Chest Congestion Children Syrup Dextromethorphan Hbr - Guaifenesin Helps relieve cough Promotes phlegm expulsion Helps to decongest the respiratory tract Administration way: Oral Boxed bottle of 8 fl oz 236 mL and measuring cup	Tamper evident: Do not use if inner seal under cap is broken or missing. SEF CADTAN FOD FIULL ABELLING	Active ingredients (in each 5 m) Purpose Dextromethorphan HBr USP 10 mg. Cough suppressant Guaitenesin, USP 100 mg.	Warnings Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have Lave a cough that occurs with too much phegm (mucus)	cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema Stop use and ask a doctor If cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be siens of a serious condition.	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	Directions ■ take every 4 hours = do not take more than 6 doses in any 24-hour period ■ take every 4 hours = do not take more than 6 doses in any 24-hour period ■ measure only with dosing cup provided ⊕ keep dosing cup with product ■ measure only with dosing cup provided ⊕ keep dosing cup with product ■ Measure only with dosing cup provided ⊕ keep dosing cup with product ■ Measure only with dosing cup provided ⊕ keep dosing cup with product ■ Measure only with dosing cup provided ⊕ keep dosing cup with product ■ Measure only with dosing cup provided ■ Measure only with product Adults and children 12 years and over 2 teaspoonful (10 m) Children 6 to under 12 years Ask a doctor	Inactive ingredients citric acid, flavor, glycerin, menthol, polyethylene glycol, purified water, sodium benzoate, sodium saccharin, sorbitol	Manufactured by KJD Pharma Pvf Ltd. 14B, 3 Kladb, Naubad, Bidar, Kamataka 585401, India	Lot DO NOT PRINT HERE NON-VARNISH AREA FOR BATCH CODE & EXP DATE
NDC 85722-0001-6	ing	cose sant rant	NG.		-	Bt	ycol,		

NDC 85722-0001-8 Cough & Chest Congestion Children Syrup Dextromethorphan Hbr - Guaifenesin	nder cap is broken or missing	Purpose Cough suppressant Expectorant	iption mono , psychiatrio weeks after drug contai	k a doctor before use if you have cough that occurs with too much phegm (mucus) cough that lasts or is chronic such as occurs with smoking, asthma,	ore than 7 days, comes back, t headache. These could be	nt or breast-leeding, ask a health professional before use. of reach of children. In case of overdose, get medical help or Poison Control Center right away.	Directions ■ take every 4 hours ■ do not take more than 6 doses in any 24-hour period Ф measure only with dosing cup provided	DOSE 2 teaspoonful (10 ml) 1 teaspoonful (5 ml) Ask a chortor	polyethylene glycol, propylene glycol ilum saccharin, sorbitol		T HERE BATCH CODE
Helps relieve cough Promotes phlegm expulsion Helps to decongest the respiratory tract	Do not use if inner seal under	380	if you are now taki (certain drugs for Parkinson's disea not know if your p	use if you have with too much phegm (i is chronic such as occur	or eniprysenta a doctor if cough lasts more than 7 d by fever, rash, or persistent headache. condition	feeding, ask a health children. In case of o of Center right away.	do not take more than using cup provided \oplus	2 years and over 2 years	n, menthol, enzoate, soo	.B, 3 Kladb, ka 585401, India	DO NOT PRIN NON-VARNISH AREA FOR & EXP DATE
Administration way: Oral Boxed bottle of 16 fl oz 473 mL and measuring cup	Tamper evident: Do no	Active ingredients (in Dextromethorphan HBr U Guaifenesin, USP 100 mg.	Warmings Do not use if you are oxidase inhibitor (MAOI) (certain to emotional conditions, or Parkinso the MAOI drug. If you do not know ask a dorchor or orbarmacish before ask a dorchor orbarmacish before a dorchor orbarmacish before a dorchor orbarmacish before a dor a dorchor orbarmacish before a dorchor orbarmacish before	Ask a doctor before use if you have acough that occurs with too much phegm (mucus) acough that lasts or is chronic such as occurs with a	curronic providuals or emproyeemed Stop use and ask a doctor if cough lasts more than 7 days, or is accompanied by fewer, rash, or persistent headache. The siens of a serious condition.	If pregnant or breast- Keep out of reach of contact a Poison Contr	Directions ■ take every 4 hours ⊕ measure only with do	AGE Adults and chikiren 12 years and over Children 6 to under 12 years Children under 6 years	Inactive ingredients citric acid, flavor, glycerin, i purified water, sodium benz	Manufactured by KJD Pharma Pvt Ltd, 14B, Naubad, Bidar, Karmataka t	Lot DO NOT Exp NON-VARNISH, &

DEXTROMETHORPHAN HYDROBROMIDE AND GUAIFENESIN

dextromethorphan hydrobromide and guaifenesin syrup

Product Information						
Product Type	HUMAN OTC DRUG	Item Code (Source) NDC:85722-00				
Route of Administration	ORAL					
Active Ingredient/Active	Moiety					
Ingre	dient Name		Basis of St	rength	Strength	
Ingre DEXTROMETHORPHAN HYDROB (DEXTROMETHORPHAN - UNII:7355)	ROMIDE (UNII: 9D2RTI9KYH	1)	Basis of St DEXTROMETHOR HYDROBROMIDE	•	Strength 10 mg in 5 mL	
DEXTROMETHORPHAN HYDROB	ROMIDE (UNII: 9D2RTI9KYH (3ROTS)		DEXTROMETHOR	•	10 mg	

	edients				
		Ingredient Name	e		Strength
ANHYDROUS CITE	RIC ACID (UNII:	XF417D3PSL)			
GLYCERIN (UNII: P	DC6A3C0OX)				
MENTHOL (UNII: L	7T10EIP3A)				
POLYETHYLENE O	GLYCOL, UNSP	ECIFIED (UNII: 3WJQ0SDV	V1A)		
PROPYLENE GLYC	COL (UNII: 6DC9	Q167V3)			
WATER (UNII: 0590	QF0KO0R)				
SODIUM BENZOA	TE (UNII: 0J245	FE5EU)			
SACCHARIN SODI	UM (UNII: SB8Z	(UX40TY)			
SORBITOL (UNII: 5	06T60A25R)				
Product Char	acteristics				
Color			Score		
Shape			Size		
Flavor		CHERRY	Imprint	Code	
Contains					
Packaging					
	Pa	ackage Description		Marketing Start Date	Marketing End Date
# Item Code		ackage Description OTTLE; Type 0: Not a Cor	nbination	-	
 # Item Code 1 NDC:85722- 0001-4 NDC:95722 	118 mL in 1 B Product			Date	
 # Item Code 1 NDC:85722- 0001-4 2 NDC:85722- 0001-8 NDC:85722- 0001-8 	118 mL in 1 B Product 236 mL in 1 B Product	OTTLE; Type 0: Not a Cor	nbination	Date 02/01/2021	
 # Item Code 1 NDC:85722- 0001-4 2 NDC:85722- 0001-8 3 NDC:85722- 	118 mL in 1 B Product 236 mL in 1 B Product 473 mL in 1 B	OTTLE; Type 0: Not a Cor OTTLE; Type 0: Not a Cor	nbination	Date 02/01/2021 02/01/2021	
 # Item Code 1 NDC:85722- 0001-4 2 NDC:85722- 0001-8 3 NDC:85722- 0001-6 	118 mL in 1 B Product 236 mL in 1 B Product 473 mL in 1 B Product	OTTLE; Type 0: Not a Cor OTTLE; Type 0: Not a Cor OTTLE; Type 0: Not a Cor	nbination	Date 02/01/2021 02/01/2021	
 NDC:85722- 0001-4 NDC:85722- 0001-8 NDC:85722- 	118 mL in 1 B Product 236 mL in 1 B Product 473 mL in 1 B Product	OTTLE; Type 0: Not a Cor OTTLE; Type 0: Not a Cor OTTLE; Type 0: Not a Cor	nbination	Date 02/01/2021 02/01/2021	

Labeler - KJD Pharma Private Limited (877256314)

Revised: 6/2025

KJD Pharma Private Limited