

**PEPTO BISMOL FAST MELTS WINTER FROST FLAVORED- bismuth subsalicylate tablet, chewable**  
**The Procter & Gamble Manufacturing Company**

-----  
**Pepto Bismol Fast Melts Winter Frost Flavored**

***Drug Facts***

**Active ingredient (in each soft chewable tablet)**

Bismuth subsalicylate 262 mg

**Purpose**

Upset stomach reliever and antidiarrheal

**Uses**

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - heartburn
  - indigestion
  - nausea
  - gas
  - belching
  - fullness

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use if**

you have

- an ulcer
- a bleeding problem

- bloody or black stool

### **Ask a doctor before use if you have**

- fever
- mucus in the stool

### **Ask a doctor or pharmacist before use if you are**

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

### **When using this product**

a temporary, but harmless, darkening of the stool and/or tongue may occur

### **Stop use and ask a doctor if**

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast feeding,** ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

- fully chew before swallowing
- adults and children 12 years and over:
  - 2 soft chewable tablets every 1/2 hour or 4 soft chewable tablets every hour as needed for diarrhea
  - 2 soft chewable tablets every 1/2 hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (16 soft chewable tablets) in 24 hours • use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

### **Other information**

- **each soft chewable tablet contains:** • calcium 55 mg • salicylate 100 mg • low sodium
- Store at no greater than 25°C. Avoid excessive heat (over 40°C).

**Inactive ingredients**

calcium carbonate, carrageenan, corn syrup, FD&C Blue No. 1, FD&C Red No. 40, flavor, hydrogenated coconut oil, soy lecithin, sucrose, water

**Questions?****1-800-717-3786****TAMPER EVIDENT: DO NOT USE IF INNER PRINTED SEAL ON BOTTLE IS  
BROKEN OR MISSING.****DIST. BY: PROCTER & GAMBLE,  
CINCINNATI, OH 45202****PRINCIPAL DISPLAY PANEL - 24 SOFT CHEWABLE TABLETS****Pepto Bismol®**

Bismuth Subsalicylate

Upset Stomach Reliever/Antidiarrheal

FAST MELTS

COATS & COOLS

WINTER FROST FLAVORED

24 SOFT CHEWABLE TABLETS



DIST. BY: PROCTER & GAMBLE, CINCINNATI, OH 45202

P&G Patents: www.pg.com/patents  
www.pg.com/pepto-bismol.com

21140427

# Pepto Bismol®

Bismuth Subsalicylate - Upset Stomach Reliever/Antidiarrheal



COATS & COOLS

WINTER FROST FLAVORED

24 SOFT CHEWABLE TABLETS

TAMPER EVIDENT: DO NOT USE IF INNER PRINTED SEAL ON BOTTLE IS BROKEN OR MISSING.

Drug Facts	Purpose
<b>Active ingredient (in each soft chewable tablet)</b> Bismuth subsalicylate 262 mg.....Upset stomach reliever and antidiarrheal	
<b>Uses</b> relieves <ul style="list-style-type: none"> <li>• upset stomach due to overindulgence in food and drink, including:</li> <li>• heartburn</li> <li>• gas</li> </ul>	<ul style="list-style-type: none"> <li>• travelers' diarrhea</li> <li>• indigestion</li> <li>• belching</li> <li>• diarrhea</li> <li>• nausea</li> <li>• fullness</li> </ul>
<b>Warnings</b> Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Contains salicylate. Do not take if you are <ul style="list-style-type: none"> <li>• allergic to salicylates (including aspirin)</li> <li>• taking other salicylate products</li> </ul> Do not use if you have <ul style="list-style-type: none"> <li>• an ulcer</li> <li>• a bleeding problem</li> <li>• bloody or black stool</li> </ul>	

PEEL BACK FOR DRUG FACTS ▶

<b>Drug Facts (continued)</b>
Ask a doctor before use if you have <ul style="list-style-type: none"> <li>• fever</li> <li>• mucus in the stool</li> </ul>
Ask a doctor or pharmacist before use if you are taking any drug for <ul style="list-style-type: none"> <li>• anticoagulation (thinning the blood)</li> <li>• diabetes</li> <li>• gout</li> <li>• arthritis</li> </ul>
When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur
Stop use and ask a doctor if <ul style="list-style-type: none"> <li>• symptoms get worse or last more than 2 days</li> <li>• ringing in the ears or loss of hearing occurs</li> <li>• diarrhea lasts more than 2 days</li> </ul>
If pregnant or breast feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
<b>Directions</b> <ul style="list-style-type: none"> <li>• fully chew before swallowing</li> <li>• adults and children 12 years and over:               <ul style="list-style-type: none"> <li>• 2 soft chewable tablets every ½ hour or 4 soft chewable tablets every hour as needed for diarrhea</li> <li>• 2 soft chewable tablets every ½ hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)</li> </ul> </li> <li>• do not exceed 8 doses (16 soft chewable tablets) in 24 hours</li> <li>• use until diarrhea stops but not more than 2 days</li> <li>• children under 12 years: ask a doctor</li> <li>• drink plenty of clear fluids to help prevent dehydration caused by diarrhea</li> </ul>
<b>Other information</b> <ul style="list-style-type: none"> <li>• each soft chewable tablet contains:               <ul style="list-style-type: none"> <li>• calcium 55 mg</li> <li>• salicylate 110 mg</li> </ul> </li> <li>• low sodium</li> <li>• Store at no greater than 25°C. Avoid excessive heat (over 40°C).</li> </ul>
<b>Inactive ingredients</b> calcium carbonate, carrageenan, corn syrup, FD&C Blue No. 1, FD&C Red No. 40, flavor, hydrogenated coconut oil, soy lecithin, sucrose, water
<b>Questions?</b> 1-800-717-3786

## 5 SYMPTOM RELIEF



NAUSEA      HEARTBURN      INDIGESTION      UPSET STOMACH      DIARRHEA

PEPTO BISMOL FAST MELTS WINTER FROST FLAVORED			
bismuth subsalicylate tablet, chewable			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69423-832
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength

<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	262 mg
--	-----------------------	--------

<b>Inactive Ingredients</b>	
Ingredient Name	Strength
<b>CARRAGEENAN</b> (UNII: 5C69YCD2YJ)	
<b>CORN SYRUP</b> (UNII: 9G5L16BK6N)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>HYDROGENATED COCONUT OIL</b> (UNII: JY81OXM1OM)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>CALCIUM CARBONATE</b> (UNII: H0G9379FGK)	

<b>Product Characteristics</b>			
<b>Color</b>	blue	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	22mm
<b>Flavor</b>	MINT	<b>Imprint Code</b>	C
<b>Contains</b>			

<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-832-24	24 in 1 BOTTLE; Type 0: Not a Combination Product	03/05/2024	

<b>Marketing Information</b>			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	03/05/2024	

**Labeler** - The Procter & Gamble Manufacturing Company (004238200)

Revised: 10/2025

The Procter & Gamble Manufacturing Company