

SIGNATURE CARE ALLERGY RELIEF D- cetirizine hcl, pseudoephedrine hcl tablet, extended release

Safeway

Better Living Brands LLC Allergy Relief-D Drug Facts

Active ingredients (in each extended release tablet)

Cetirizine HCl 5 mg

Pseudoephedrine HCl 120 mg

Purpose

Antihistamine

Nasal Decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- nasal congestion
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- thyroid disease
- diabetes
- glaucoma
- high blood pressure
- trouble urinating due to an enlarged prostate gland

- liver or kidney disease. Your doctor should determine if you need a different dose.

Ask a doctor or pharmacist before use if you are

taking tranquilizers or sedatives.

When using this product

- do not use more than directed
- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- you get nervous, dizzy, or sleepless
- symptoms do not improve within 7 days or are accompanied by fever

If pregnant or breast-feeding:

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not break or chew tablet; swallow tablet whole

adults and children 12 years and over	take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.
adults 65 years and over	ask a doctor
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- store between 20° to 25°C (68° to 77°F)

Inactive ingredients

colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, talc, yellow iron oxide

Questions or comments?

1-888-723-3929

Principal Display Panel

12 HOUR | ORIGINAL PRESCRIPTION STRENGTH

Allergy Relief-D

Cetirizine Hydrochloride and Pseudoephedrine

Hydrochloride Extended Release Tablets 5 mg/120 mg

Antihistamine/Nasal Decongestant

Compare to Zyrtec-D[®] active ingredients

Allergy & congestion

Indoor & outdoor allergies

12 Hour relief of:

sneezing; runny nose; sinus pressure; itchy, watery eyes; itchy throat or nose; nasal congestion

Actual Size

12 Extended Release Tablets



Drug Facts**Active ingredients (in each extended release tablet) Purpose**

Cetirizine HCl 5 mg.....Antihistamine
 Pseudoephedrine HCl 120 mg.....Nasal Decongestant

Uses

temporarily relieves these symptoms due to hay

Drug Facts (continued)

fever or other upper respiratory allergies:
 ■ runny nose ■ sneezing ■ itchy, watery eyes
 ■ itching of the nose or throat ■ nasal congestion
 ■ reduces swelling of nasal passages
 ■ temporarily relieves sinus congestion and pressure
 ■ temporarily restores freer breathing through the nose

Drug Facts (continued)**Warnings****Do not use**

- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease ■ thyroid disease ■ diabetes
- glaucoma ■ high blood pressure
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.**When using this product**

- do not use more than directed
- drowsiness may occur ■ avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- an allergic reaction to this product occurs. See medical help right away.

Drug Facts (continued)

- you get nervous, dizzy, or sleepless
- symptoms do not improve within 7 days or are accompanied by fever

If pregnant or breast-feeding:

- if breast-feeding: not recommended
 - if pregnant: ask a health professional before use.
- Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

do not break or chew tablet; swallow tablet whole	
adults and children 12 years and over	take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.
adults 65 years and over	ask a doctor
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- store between 20° to 25°C (68° to 77°F)

Inactive ingredients

colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, talc, yellow iron oxide

Drug Facts (continued)

Questions or comments? 1-888-723-3929



17553 L J C1

DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

OUR PROMISE
 QUALITY & SATISFACTION
 100%
 GUARANTEED
 OR YOUR MONEY BACK

MADE IN INDIA
 DISTRIBUTED BY: BETTER LIVING BRANDS LLC
 P.O. BOX 99, PLEASANTON, CA 94566-0099
 1-888-723-3929/www.betterlivingbrandsllc.com
 *This product is not manufactured or distributed by McNeil Consumer Healthcare, division of McNeil-PPC, Inc., distributor of Zyrtec-D®.

SIGNATURE CARE ALLERGY RELIEF D

cetirizine hcl, pseudoephedrine hcl tablet, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-176
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CETIRIZINE HYDROCHLORIDE (UNII: 64O047KTOA) (CETIRIZINE - UNII:YO7261ME24)	CETIRIZINE HYDROCHLORIDE	5 mg
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DD19F)	PSEUDOEPHEDRINE HYDROCHLORIDE	120 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYDROXYPROPYL CELLULOSE (1600000 WAMW) (UNII: RFW2ET671P)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
TALC (UNII: 7SEV7J4R1U)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	

Product Characteristics

Color	WHITE (one side white one side light yellow)	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	5029;5;120
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-176-53	12 in 1 CARTON	09/12/2018	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:21130-176-62	24 in 1 CARTON	10/15/2015	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077170	10/15/2015	

Labeler - Safeway (009137209)