

**4332 FIRST AID KIT- 4332 first aid kit
Honeywell Safety Products USA, INC**

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

0498-4332: First Aid Kit (Alcohol wipe, Eye Wash, FABC, BZK, sting rel, neomycinASA- Z019816)

Active ingredient

Sterile Water 99%

Purpose

Eyewash

Uses

- For flushing the eye to remove loose foreign material, air pollutants, or chlorinated water

Warnings

For external use only- Obtain immediate medical treatment for all open wounds in or near the eyes. To avoid contamination, do not touch tip of container to any surface. Do not reuse. Once opened, discard.

Do not use

- if solution changes color or becomes cloudy
- if you have open wounds in or near the eyes, get medical help right away

Stop use and ask a doctor if

- you experience eye pain
- changes in vision
- continued redness or irritation of the eye
- condition worsens or persists

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- remove contacts before using
- twist top to remove

- flush the affected area as needed
- control rate of flow by pressure on the bottle
- if necessary, continue flushing with emergency eyewash or shower

Inactive Ingredients

sodium chloride, sodium phosphate dibasic, sodium phosphate monobasic

Questions?

Call **1-800-430-5490**

Honeywell Safety Products USA, Inc. Smithfield, RI 02917

First Aid Burn Cream

Active ingredient

Benzalkonium chloride 0.13%

Lidocaine HCl 0.5%

First Aid Burn Cream

Purpose

First aid antiseptic

External analgesic

First Aid Burn Cream

Uses

- prevent skin infection
- for temporary relief of pain associated with minor burns

First Aid Burn Cream

Warnings

For external use only

Do not use

- in or near the eyes
- if you are allergic to any of the ingredients
- in large areas of the body, particularly over raw surfaces or blistered areas
- for more than 10 days

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites

- serious burns

Stop use and ask a doctor if

- condition worsens
- symptoms persist for more than 7 days or clear up and occurs again within a few days

First Aid Burn Cream

Directions

- **adults and children 2 years of age and older:**
- clean the affected area
- apply a small amount of this product (equal to the surface area of the tip of a finger) onto affected area 1 to 3 times daily
- may be covered with a sterile bandage
- children under 2 years of age: consult a doctor

Other information

- tamper evident sealed packets
- do not use if packet is opened or torn

Inactive ingredients

aloe barbadensis juice, cetyl alcohol, diazolidinyl urea, edetate disodium, glycerin, glyceryl stearate SE, methylparaben, mineral oil, PEG-100, propylene glycol, propylparaben, stearic acid, trolamine, water

Questions

1-800-430-5490

BZK Antiseptic Wipe

Active ingredient

Benzalkonium chloride 0.13%

BZK

Purpose

First aid antiseptic

BZK

Uses

Antiseptic cleansing of face, hands, and body without soap and water

BZK**Warnings**

For external use only

BZK**Do not use**

- in the eyes or over large areas of the body
- on mucous membranes
- on irritated skin
- in case of deep puncture wounds, animal bites or serious burns, consult a doctor
- longer than 1 week unless directed by a doctor

Stop use and ask a doctor if

- if irritation, redness or other symptoms develop
- the condition persists or gets worse

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

BZK**Directions**

- .tear open packet and use as a washcloth

BZK**Other information**

- store at room temperature 15 ° to 30 ° C (59 ° - 86 °F)
- do not reuse towelette

BZK**Inactive ingredients**

water

BZK**Questions**

1-800-430-5490

Sting Relief**Active ingredient**

Ethyl alcohol 50.0%

Lidocaine HCl 2.0%

Sting Relief

Purpose

Antiseptic

Topical pain relief

Sting Relief

Uses

- prevent infection in minor scrapes, and temporary relief of itching of insect bites

Sting Relief

Warnings

For external use only

Flammable, keep away from open fire or flame

Do not use

- over large areas of the body
- in eyes
- over raw or blistered areas

Stop use and ask a doctor

- if conditions worsen or persist for more than 7 days or clear up and occur again within a few days

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away

Sting Relief

Directions

- **adults and children 2 years and older:** Apply to cleaned affected area not more than 3 times daily.
- children under 2 years of age: consult a doctor.

Sting Relief

Inactive ingredients

benzalkonium chloride, menthol, and purified water

Sting relief

Questions or Comments

1-800-430-5490

Neomycin Antibiotic Ointment

Active ingredient

Neomycin sulfate (5 mg equivalent to 3.5 mg Neomycin base)

Neomycin Antibiotic Ointment

Purpose

First aid antibiotic

Neomycin Antibiotic Ointment

Uses

first aid to help prevent infection in - minor cuts - scrapes - burns

Neomycin Antibiotic Ointment

Warnings

For external use only

Do not use

- in the eyes
- over large areas of the body

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Stop use and ask a doctor if

- the condition persists or gets worse
- a rash or other allergic reaction develops
- you need to use longer than 1 week

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away

Neomycin Antibiotic Ointment

Directions

- clean the affected area
- apply a small amount of the product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage

Neomycin Antibiotic Ointment

Other information

store at 15 ° to 25 °C (59 ° to 77 °F)

Neomycin Antibiotic Ointment
Inactive ingredient

petrolatum

Neomycin Antibiotic Ointment
Questions

1-800-430-5490

Alcohol Wipes
Active ingredient

Isopropyl alcohol 70%

Alcohol
Purpose

First aid antiseptic

Alcohol
Uses

- first aid to help prevent infection in minor cuts, scrapes, and burns

Alcohol
Warnings

For external use only

Flammable, keep away from fire and flame

Do not use

- in or near eyes
- over large areas of the body

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

When using this product

- do not use longer than 1 week unless directed by a doctor

Stop use and consult a doctor if

- condition persists or gets worse

Keep out of the reach of children

If swallowed, get medical help or contact a Poison Control center right away

Alcohol

Directions

- clean the affected area
- may be covered with a sterile bandage
- apply wipe to affected area 1 to 3 times daily
- discard wipe after single use

Alcohol

Other information

- store at room temperature 15 ° to 25 ° C (59 ° to 77 ° F)
- do not use if packet is torn or opened

Alcohol

Inactive ingredient

water

Alcohol

Questions

1-800-430-5490

Aspirin

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Aspirin

Purpose

Pain reliever/fever reducer

Aspirin

Uses

temporarily reduces fever and relieves minor aches and pains associated with:

- a cold
- headache
- toothache

- muscular aches
- backache
- minor pain of arthritis
- premenstrual and menstrual periods

Aspirin

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools

- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- ringing in the ears or loss of hearing occurs
- any new symptoms appear

If pregnant or breast-feeding,

- If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last three months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact Poison Control Center right away.

Aspirin

Directions

- drink a full glass of water with each dose
- adults and children 12 years of age and older: take 1 or 2 tablets every 4 hours while symptoms last, not more than 12 tablets in 24 hours
- children under 12 years of age: consult a doctor

Aspirin

Other information

store at room temperature 15° - 30°C (59° - 86°F)

TAMPER EVIDENT PACKETS - DO NOT USE IF OPEN OR TORN

Aspirin

Inactive ingredient

corn starch, croscarmellose sodium*, hypromellose*, microcrystalline cellulose*, mineral oil*, polyethylene glycol*, povidone, propylene glycol, silicon dioxide, stearic acid*, titanium dioxide*

*may contain these ingredients

Aspirin

Question

1-800-430-5490

4332


Z019816 Kit Contents

1 ADHESIVE BDG,PLSTIC,1"X3"16PER
1 TWEEZER PLASTICS 4"
1 FIRST AID GUIDE ASHI
1 TAPE ADHESIVE 1"X 5 YD PLSTC
1 GAUZE CLEAN-WRAP BDGE N/S 2"
1 GAUZE CLEAN-WRAP BDGE N/S 3"
1 ABD COMBINE PAD 5" X 9"
1 MICROSHIELD W/VNL GLV/ALCL
1 4OZ BFS EYEWASH TRILINGUAL BOTTLE
1 SCISSOR BDGE 4" RED PLS HDL
2 WIPE GERM, SANIZIDE PLUS
1 IMPERVIOUS GOWN ONE SIZE EA
1 SHOE COVERS LARGE (1-PAIR)
LBL STOCK 6-3/8"X4"
LBL STOCK 4"X2-7/8"
1 LBL STOCK 3"x1-7/8"
1 LBL CONTS 6 3/4"X3 1/2" ID B
15 BZK ANTISEPTIC WIPE, BULK
1 LABEL COVER, GRAINGER Z019816
4 x 1 PR LRG NITRILE GLVES ZIP BAG
1 PICK-UP SCOOP W/SCRAPER
1 RED-Z FLUID CONTROL PCH 3/4OZ
6 FIRST AID BURN CREAM 1.0GR PKT EACH
6 POUCH NEOMYCIN ANTIBIOTIC .9 G
6 WIPE ALCOHOL PREP IPA 70% (DUKAL)
1 KIT 36 UNIT PLASTIC
1 BAG BIOHAZARD 24 X 24 RED
3 TOWEL 3-FOLD WHITE
6 SAFETEC STING RELIEF WIPES BULK
1 TRI BNDG NON WOVEN 40"X40"X56"
1 COLD PACK UNIT 4"X6" BULK
2 EYE PADS STD OVAL STERILE
2 GAUZE PADS 3"X3" 12PLY

- 2 GAUZE PADS 4"X4" 12PLY
- 2 WOVEN FINGERTIP BANDAGE 2"
- 3 WOVEN KNUCKLE BANDAGE
- 20 WOVEN BANDAGE 1" X 3"
- 1 BOUFFANT CAP 24"
- 2 TWIST TIE
- 1 ZIP LOCK BAG 8 X 10" 2 MIL
- 1 FACE MASK/EYE SHIELD
- 3 ASPIRIN BULK 2/PK

Eye Wash Package label

#32-004513 Rev. H




Top Panel
3/32" from die edges



Back
1/8" from die edges



Label Patent Number
Can be anywhere on
the label at 3.5pt minimum
(use any dark color on that layer)





Base
3/32" from die edges

Printable Text Area

First Aid Burn Cream Principal Display Panel



Principal Display Panel



Sting Relief Principal Display Panel

Honeywell

825366 Rev B

Honeywell

032043P

Honeywell

032043P

Sting Relief Wipes



Use for:
Minor Cuts • Scrapes • Insect Bites

Single Use Pouches
Saturated Wipes

100 wipes

Sting Relief Wipes



Use for:
Minor Cuts • Scrapes • Insect Bites

Single Use Pouches
Saturated Wipes

100 wipes

POUR
HERE &
PULL TO
OPEN

Drug Facts

Active Ingredients	Purpose
Ethyl alcohol 50.0%	First aid antiseptic
Lidocaine HCl 2.0%	Topical anesthetic

Uses First aid to help prevent infection in minor scrapes and temporary relief of itching of insect bites.

Warnings

For external use only.

Flammable, keep away from fire or flame.

Do not use • over large areas of the body • in eyes • over raw or blistered areas

Stop use and ask a doctor • if conditions worsen or persist for more than 7 days or clear up and occur again within a few days.

Keep out of reach of children.

If swallowed get medical help or contact Poison Control center right away.

Directions

• adults and children 2 years and older: apply to cleaned affected area not more than 3 times daily.

• children under 2 years of age: consult a doctor.

Inactive Ingredients

benzalkonium chloride, menthol, purified water

Questions or comments?

1-800-430-5490

Distributed by
Honeywell Safety
Products, USA, Inc.
Smithfield, WI 02917

www.honeywellsafety.com

USA
1-800-430-5490
0-401-343-4400



Made in USA
Packaged in Mexico

Neomycin Antibiotic Ointment Principal Display Panel

822568-25

**Neomycin First
aid antibiotic**

020126-25

Neomycin
First aid antibiotic
Neomycin sulfate
First aid antibiotic

Distributed by:
Honeywell Safety Products
USA, Inc. Smithfield,
RI 02917

25 Packets, Net Wt 1/32 oz (0.9 g) each

Neomycin First aid antibiotic

822568-25

Drug Facts

Active ingredient (in each gram)	Purpose
Neomycin sulfate (equivalent to 3.5 mg neomycin)	First aid antibiotic

Use first aid to help prevent infection in minor • cuts • scrapes • burns

Warnings

For external use only

Allergy alert • do not use if you are allergic to any of the ingredients

Do not use • in the eyes • over large areas of the body

Ask a doctor before use if you have • deep or puncture wounds • animal bites • serious burns

Stop use and ask a doctor if • conditions persists or gets worse • rash or other allergic reaction develops
• you need to use longer than one week

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions


• clean the affected area
• apply a small amount of product (equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
• may be covered with a sterile bandage

Other information store at 15° to 25°C (59° to 77°F)

Inactive ingredient petrolatum

Questions or comments? 1-800-430-5490

Alcohol
Principal Display Panel



**Aspirin
Principal Display Panel**



4332 Kit Label
Z019816

FIRST AID

FIRST AID & BBP, ANSI
25 PERSON



GRAINGER®

FOR THE ONES WHO GET IT DONE

GRAINGER.COM®

Distributed by Honeywell Safety Products USA, Inc. Smithfield, RI 02917

470017065B

4332 FIRST AID KIT

4332 first aid kit kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0498-4332
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-4332-01	1 in 1 KIT; Type 0: Not a Combination Product	10/18/2018	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	118 mL
Part 2	6 POUCH	2.4 mL
Part 3	6 PACKET	5.4 g
Part 4	6 PACKET	5.4 g
Part 5	15 PACKET	21 mL
Part 6	10 POUCH	4 mL
Part 7	3 PACKET	6

Part 1 of 7

EYESALINE EMERGENCY EYEWASH

purified water liquid

Product Information

Item Code (Source)	NDC:0498-0100
---------------------------	---------------

Route of Administration	OPHTHALMIC
--------------------------------	------------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
WATER (UNII: 059QF0KO0R) (WATER - UNII:059QF0KO0R)	WATER	98.6 mL in 100 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM PHOSPHATE, DIBASIC (UNII: GR686LBA74)	
SODIUM PHOSPHATE, MONOBASIC, MONOHYDRATE (UNII: 593YOG76RN)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0100-02	118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M018	12/18/2018	

Part 2 of 7

STING RELIEF PAD

ethyl alcohol, lidocaine swab

Product Information

Item Code (Source)	NDC:0498-0733
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE ANHYDROUS	20 mg in 1 mL
ALCOHOL (UNII: 3K9958V90M) (ALCOHOL - UNII:3K9958V90M)	ALCOHOL	0.5 mL in 1 mL

Inactive Ingredients

Ingredient Name	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	
MENTHOL (UNII: L7T10EIP3A)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		0.4 mL in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/23/2017	

Part 3 of 7

FIRST AID BURN

benzalkonium chloride, lidocaine hydrochloride cream

Product Information

Item Code (Source)	NDC:0498-0903
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	0.13 g in 100 g
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE	0.5 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
DIAZOLIDINYL UREA (UNII: H5RIZ3MPW4)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
ALOE VERA LEAF (UNII: ZY81Z83H0X)	
WATER (UNII: 059QF0KO0R)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
PEG-100 STEARATE (UNII: YD01N1999R)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
TROLAMINE (UNII: 9O3K93S3TK)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		0.9 g in 1 PACKET; Type 0: Not a Combination		

Product

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/20/2017	

Part 4 of 7

NEOMYCIN

antibiotic ointment

Product Information

Item Code (Source)	NDC:0498-0730
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN SULFATE	3.5 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
PETROLATUM (UNII: 4T6H12BN9U)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0730-01	0.9 g in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		03/31/2010	

Part 5 of 7

ANTISEPTIC TOWELETTE

benzalkonium chloride liquid

Product Information

Item Code (Source) NDC:0498-0501

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	1.3 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		1.4 mL in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/21/2017	

Part 6 of 7

ALCOHOL WIPE

isopropyl alcohol swab

Product Information

Item Code (Source) NDC:0498-0143

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	0.7 mL in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0143-04	0.4 mL in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	

Part 7 of 7

ASPIRIN

aspirin tablet

Product Information

Item Code (Source)	NDC:0498-0114
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POVIDONE (UNII: FZ989GH94E)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE 2208 (100 MPA.S) (UNII: B1QE5P712K)	

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
MINERAL OIL (UNII: T5L8T28FGP)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	FR21
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0114-01	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		10/18/2018	

Labeler - Honeywell Safety Products USA, INC (118768815)

Revised: 1/2024

Honeywell Safety Products USA, INC