

**4333 FIRST AID KIT- 4333 first aid kit
Honeywell Safety Products USA, INC**

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

0498-4333: First Aid Kit (FABC, EW, BZK wipe, neomycin, alcohol wipes, sting relief, ASA- Z019818)

**First Aid Burn Cream
Active ingredient**

Benzalkonium chloride 0.13%

Lidocaine HCl 0.5%

**First Aid Burn Cream
Purpose**

First aid antiseptic

External analgesic

**First Aid Burn Cream
Uses**

- prevent skin infection
- for temporary relief of pain associated with minor burns

**First Aid Burn Cream
Warnings**

For external use only

Do not use

- in or near the eyes
- if you are allergic to any of the ingredients
- in large areas of the body, particularly over raw surfaces or blistered areas
- for more than 10 days

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Stop use and ask a doctor if

- condition worsens
- symptoms persist for more than 7 days or clear up and occurs again within a few days

First Aid Burn Cream

Directions

- **adults and children 2 years of age and older:**
- clean the affected area
- apply a small amount of this product (equal to the surface area of the tip of a finger) onto affected area 1 to 3 times daily
- may be covered with a sterile bandage
- children under 2 years of age: consult a doctor

First Aid Burn Cream

Other information

- tamper evident sealed packets
- do not use if packet is opened or torn

First Aid Burn Cream

Inactive ingredients

aloe barbadensis juice, cetyl alcohol, diazolidinyl urea, edetate disodium, glycerin, glyceryl stearate SE, methylparaben, mineral oil, PEG-100, propylene glycol, propylparaben, stearic acid, trolamine, water

First Aid Burn Cream

Questions

1-800-430-5490

Alcohol Wipe

Active ingredient

Isopropyl alcohol 70%

Alcohol Wipe

Purpose

First aid antiseptic

Alcohol Wipe

Uses

- first aid to help prevent infection in minor cuts, scrapes, and burns

Alcohol Wipe

Warnings

For external use only

Do not use

- in the eyes
- over large areas of the body

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burn

When using this product

- do not use longer than one week unless directed by a doctor

Stop use and consult a doctor

- if condition persists or gets worse

Keep out of reach of children

- If swallowed, get medical help or contact a Poison Control Center right away.

Alcohol Wipe

Directions

- clean the affected area
- apply wipe to affected area 1 to 3 times daily
- may be covered with a sterile bandage
- discard wipe after single use

Alcohol Wipe

Other information

store at room temperature 15 ° to 25 ° C (59 ° to 77 °F)

Alcohol Wipe

Inactive ingredient

water

Alcohol Wipe

Questions

1-800-430-5490

Eyewash

Active ingredient

Sterile Water 99%

Eyewash

Purpose

Eyewash

Eyewash

Uses

- for flushing the eye to remove loose foreign material, air pollutants or chlorinated water

Eyewash

Warnings

For external use only Obtain immediate medical treatment for all open wounds in or near eyes. To avoid contamination, do not touch tip of container to any surface. Do not reuse. Once opened, discard.

Do not use

- if solution changes color or becomes cloudy
- if you have open wounds in or near the eyes, get medical help right away.

Stop use and ask a doctor if

- you experience eye pain
- changes in vision
- continued redness or irritation of the eye
- condition worsens or persists

Keep out of reach of children

- If swallowed, get medical help or contact a Poison Control Center right away.

Eyewash

Directions

- remove contacts before using
- twist top to remove
- flush the affected area as needed
- control rate of flow by pressure on the bottle
- if necessary, continue flushing with emergency eyewash or shower

Eyewash

Inactive ingredients

sodium chloride, sodium phosphate dibasic, sodium phosphate monobasic

Eyewash

Questions

1-800-430-5490

Aspirin

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)* *nonsteroidal anti-inflammatory drug

Aspirin

Purpose

Pain reliever/fever reducer

Aspirin

Uses

temporarily reduces fever and relieves minor aches and pains associated with:

- a cold
- headache
- toothache
- muscular aches
- backache
- minor pain of arthritis
- premenstrual and menstrual periods

Aspirin

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:are:

- age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - ringing in the ears or loss of hearing occurs
 - any new symptoms appear

If pregnant or breast-feeding,

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last three months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

- In case of overdose, get medical help or contact Poison Control Center right away.

Aspirin**Directions**

- drink a full glass of water with each dose
- adults and children 12 years of age and older: take 1 or 2 tablets every 4 hours while symptoms last, not more than 12 tablets in 24 hours
- children under 12 years of age: consult a doctor

Aspirin

Other information

- store at room temperature 15° - 30°C (59° - 86°F)
- TAMPER EVIDENT PACKETS
- DO NOT USE IF OPEN OR TORN

Aspirin

Inactive ingredients

corn starch, croscarmellose sodium*, hypromellose*, microcrystalline cellulose*, mineral oil*, polyethylene glycol*, povidone, propylene glycol, silicon dioxide, stearic acid*, titanium dioxide*

*may contain these ingredients

Aspirin

Questions or Comments

1-800-430-5490

Sting Relief

Active ingredient

Ethyl alcohol 50.0%

Lidocaine HCl 2.0%

Sting Relief

Purpose

Antiseptic

Topical pain relief

Sting Relief

Uses

- prevent infection in minor scrapes, and temporary relief of itching of insect bites

Sting Relief

Warnings

For external use only

Flammable, keep away from open fire or flame

Do not use

- over large areas of the body
- in eyes

- over raw or blistered areas

Stop use and ask a doctor

- if conditions worsen or persist for more than 7 days or clear up and occur again within a few days

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

sting Relief***Directions***

- adults and children 2 years and older: Apply to cleaned affected area not more than 3 times daily.
- children under 2 years of age: consult a doctor.

Sting Relief***Inactive ingredients***

benzalkonium chloride, menthol, and purified water

Sting Relief***Questions or Comments?***

1-800-430-5490

BZK***Active ingredient***

Benzalkonium chloride 0.13% w/v

BZK***Purpose***

First aid antiseptic

BZK***Uses***

Antiseptic cleansing of face, hands, and body without soap and water

BZK***Warnings***

For external use only

Do not use

- in the eyes or over large areas of the body
- on mucous membranes
- on irritated skin
- in case of deep puncture wounds, animal bites or serious burns, consult a doctor
- longer than 1 week unless directed by a doctor

Stop use and ask a doctor if

- if irritation, redness or other symptoms develop
- the condition persists or gets worse

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

BZK

Directions

- tear open packet and use as a washcloth

BZK

Other information

- store at room temperature 15⁰ to 30⁰ C (59⁰ - 86⁰ F)
- do not reuse towelette

BZK

Inactive ingredient

water

BZK

Questions

1-800-430-5490

Neomycin

Active ingredient (each gram contains)

Neomycin sulfate (5 mg equivalent to 3.5 mg Neomycin base)

Neomycin

Purpose

First aid antibiotic

Neomycin

Uses

- first aid to help prevent infection in

- minor cuts
- scrapes
- burns

Do not use

- in the eyes
- over large areas of the body

Neomycin***Warnings*****For external use only****Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burns

Stop use and ask a doctor if

- the condition persists or gets worse
- a rash or other allergic reaction develops
- you need to use longer than 1 week

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away

Neomycin***Directions***

- clean the affected area
- apply a small amount of the product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage

Neomycin***Other information***

- store at 15⁰ to 25⁰ C (59⁰ to 77⁰ F)

Neomycin***Inactive ingredient***

petrolatum

Neomycin***Questions***



1-800-430-5490

4333


Z 019818 KIT CONTENTS

1 ADHESIVE BDG,PLSTIC,1"X3"16PER
1 TWEEZER PLASTICS 4"
1 FIRST AID GUIDE ASHI 1
1 TAPE ADHESIVE 1"X 5 YD PLSTC
1 GAUZE CLEAN-WRAP BDGE N/S 2"
1 GAUZE CLEAN-WRAP BDGE N/S 3"
1 ABD COMBINE PAD 5" X 9"
1 4OZ BFS EYEWASH TRILINGUAL BOTTLE
1 SCISSOR BDGE 4" RED PLS HDL
LBL STOCK 6-3/8"X4"
LBL STOCK 4"X2-7/8"
1 LBL STOCK 3"x1-7/8"
1 LBL CONTS 6 3/4"X3 1/2" ID B
RIBBON WAX BLACK 8" ZEBRA
12 BZK ANTISEPTIC WIPE, BULK
1 LABEL COVER, GRAINGER Z019818
2 1 PR LRG NITRILE GLVES ZIP BAG
6 FIRST AID BURN CREAM 1.0GR PKT EACH
6 POUCH NEOMYCIN ANTIBIOTIC .9 G
6 WIPE ALCOHOL PREP IPA 70% (DUKAL)
1 KIT, PP 16 UNIT FA 1 2
SHRINK FILM 12" 100 GA
6 SAFETEC STING RELIEF WIPES BULK
1 TRI BNDG NON WOVEN 40"X40"X56"
1 COLD PACK UNIT 4"X6" BULK
2 EYE PADS STD OVAL STERILE
2 GAUZE PADS 3"X3" 12PLY
2 GAUZE PADS 4"X4" 12PLY
2 WOVEN FINGERTIP BANDAGE 2"
3 WOVEN KNUCKLE BANDAGE
20 WOVEN BANDAGE 1" X 3"
BOX CORR. PLAIN, REV C
3 ASPIRIN BULK 2/PK

**First Aid Burn Cream
Principal Display Panel**



825362T Rev. A
 white printed
 Pms 072 blue, Pms 186 red,
 Pms 7459 blue and black



Eyewash
Principal Display Panel

Honeywell

TAMPER-EVIDENT CAP.
TAPA CON SELLO DE SEGURIDAD.
BOUCHON INDICATEUR D'INFRACTION.

eyesaline®

LAVAOJOS
EYESALINE

EYESALINE
EYEWASH

LAVAGE
OCULAIRE
EYESALINE

Solución
Isotónico Estéril

Sterile
Isotonic Solution

La Solution
Isotonique Stérile

16 fl. oz. (473 mL)

NPN: 80057528
64809 1 45033 3

Drug Facts (for USA only)

Active ingredient Sterile water 99%
Purpose Eyewash
Uses for flushing the eye to remove loose foreign material, air pollutants, or chlorinated water.
Warnings
For external use only - Obtain immediate medical treatment for all open wounds in or near the eyes. To avoid contamination, do not touch tip of container to any surface. Do not reuse. Once opened, discard.
Do not use
• if solution changes color or becomes cloudy
• if you have open wounds in or near the eyes, get medical help right away
Stop use and consult a doctor if:
• you experience eye pain • changes in vision
• continued redness or irritation of the eye
• condition worsens or persists
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.
Directions
• remove contacts before using • twist top to remove
• flush the affected area as needed
• control rate of flow by pressure on the bottle
• if necessary, continue flushing with emergency eyewash or shower
Inactive ingredients sodium chloride, sodium phosphate dibasic, sodium phosphate monobasic
Questions? Call 1-800-430-5490
Honeywell Safety Products USA, Inc. Smithfield, RI. 02917

LABEL #32-0045/0 Rev. J REORDER / NUEVO PEDIDO / REAPPROVISIONNEMENT #32-000454-0000

space for lot code and supplier part number

PEEL / PELAR / PELEL

Datos de medicamento (Para EE.UU. solamente)

Ingrediente Activo Agua estéril 99%
Propósito Lavaojos
Usos para el lavado de ojo para quitar las partículas sueltas y extrañas, los contaminantes aeros, o agua de cloruro
Advertencias
Para el uso externo sólo - Obtenga tratamiento médico inmediato para todas las heridas abiertas en o cerca de los ojos. Para evitar la contaminación, no toque la punta del envase con ninguna superficie. No vuelva a usar. Vez abierto, descarte.
No se use • si la solución se enturbia o cambia de color
• si tiene heridas abiertas en o cerca del ojo, obtenga ayuda medica de inmediato
Deje de usar y consulte a un médico si:
• experimenta dolor de ojo • cambio de visión
• rojez continuo o irritación del ojo
• la condición empeora o persiste
Manténgase fuera del alcance de los niños.
En caso de ingestión accidental, obtenga atención médica o llame a un Centro de control de envenenamiento inmediatamente.
Instrucciones
• quite los lentes de contacto antes de usar la solución
• tuerza la tapa para quitar
• enjuague el área afectada según se necesite
• controle el chorro haciendo presión el la botella
• si es necesario, sigue enjuagado con un lavaojos o ducha de emergencia
Ingredientes inactivos cloruro de sodio, fosfato de sodio dibásico, fosfato de sodio monobásico
¿Preguntas? Llame al 1-800-430-5490
Honeywell Safety Products USA, Inc. Smithfield, RI. 02917

Information

Usages
Pour le rinçage des yeux afin d'enlever un corps étranger, des polluants atmosphériques ou de l'eau chlorée.
Advertissements
Pour usage externe seulement - Obtenir immédiatement des soins médicaux pour toutes les plaies ouvertes dans ou près des yeux. Pour éviter toute contamination, ne pas toucher la pointe du récipient à n'importe quelle surface. Ne pas réutiliser. Une fois ouvert, jeter.
Ne pas utiliser
• si la solution a changé de couleur ou si elle est brouillée
• si vous avez des plaies ouvertes aux yeux ou à proximité, consultez immédiatement un médecin
Cesser d'utiliser la solution et consulter un médecin
• vous ressentez une douleur oculaire • si votre vision change
• rougeur ou irritation persistante des yeux
• condition empire ou persiste
Garder hors de la portée des enfants.
En cas d'ingestion, communiquer immédiatement avec un médecin ou avec un centre antipoison.
Mode d'emploi
• enlever les verres de contact avant l'utilisation • dévisser le bouchon pour l'enlever • rincer la zone touchée selon les besoins • ajuster le débit d'écoulement de la solution en augmentant ou en réduisant la pression exercée sur le contenant
• si nécessaire, continuer de rincer avec une solution de rinçage oculaire d'urgence ou une douche
Ingédients eau stérile, chlorure de sodium, phosphate dibasique de sodium, phosphate monobasique de sodium
Des questions? Faites le 1-800-430-5490
Honeywell Safety Products USA, Inc. Smithfield, RI. 02917

Aspirin
Principal Display Panel



Sting Relief Principal Display Panel

825366 Rev B

Honeywell

032043P

032043P

Sting Relief Wipes

Sting Relief Wipes

Use for:
Minor Cuts • Scrapes • Insect Bites

Use for:
Minor Cuts • Scrapes • Insect Bites

Single Use Pouches
Saturated Wipes

100 wipes

Single Use Pouches
Saturated Wipes

100 wipes

Drug Facts

Active Ingredients	Purpose
Ethyl alcohol 50.0%	First aid antiseptic
Lidocaine HCl 2.0%	Topical analgesic

Uses - first aid to help prevent infection in minor scrapes and temporary relief of itching of insect bites.

Warnings
For external use only.
Flammable, keep away from fire or flame.

Do not use • over large areas of the body • in eyes • over raw or blistered areas

Stop use and ask a doctor • if conditions worsen or persist for more than 7 days or clear up and occur again within a few days.

Keep out of reach of children.
If swallowed get medical help or contact Poison Control center right away.

Directions
• adults and children 2 years and older: apply to cleaned affected area not more than 3 times daily
• children under 2 years of age: consult a doctor.

Inactive Ingredients
benzalkonium chloride, menthol, purified water

Questions or comments? 1-800-430-5490

Distributed by
Honeywell Safety
Products, USA, Inc.
Smithfield, RI 02917

www.honeywellsafety.com

USA
1-800-430-5490
0-401-943-5490

8 21812 01284 1

Made in USA,
Packaged in Mexico

BZK Principal Display Panel

47001083
Rev B

Antiseptic Towelettes

Honeywell

02-16-35MD

Antiseptic Towelettes

Benzalkonium chloride
First aid antiseptic

Six-Saturated Towelettes

Distributed by
Honeywell Safety
Products USA, Inc.
Smithfield, RI 02917

47001083
Rev B

Antiseptic Towelettes

Honeywell

Drug Facts

Active Ingredient	Purpose
Benzalkonium chloride 0.133% w/v	First aid antiseptic

Uses • antiseptic cleaning of face, hands and body without soap and water. • air dries in seconds

Warnings
For external use only

When using this product • do not use in the eyes or apply over large areas of the body

Ask a doctor before use • in case of deep or puncture wounds, animal bites, or serious burns

Stop use and consult a doctor if

• irritation, redness or other symptoms develop • condition persists or gets worse

Do not use • longer than 1 week unless directed by doctor

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions • tear open packet, unfold and use as washcloth

Other information



• store at room temperature 15° -30° C (59° -86° F) • do not reuse towelette

Inactive ingredient water

Questions or comments 1-800-430-5490

Neomycin
Principal Display Panel

796041-25 Rev A Unit Carton Printing Plate for "C" size carton.



Z019818

FIRST AID

COMPLIANCE PACKAGE, BULK
25 PERSON



GRAINGER
FOR THE ONES WHO GET IT DONE

GRAINGER.COM®

Distributed by Honeywell SafetyProducts USA, Inc. Smithfield, RI 02917

47001707RA

4333 FIRST AID KIT

4333 first aid kit kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0498-4333
--------------	----------------	--------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-4333-01	1 in 1 KIT; Type 0: Not a Combination Product	10/18/2018	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	6 PACKET	5.4 g
Part 2	1 BOTTLE	118 mL
Part 3	6 POUCH	2.4 mL
Part 4	3 PACKET	6
Part 5	6 PACKET	5.4 g
Part 6	6 POUCH	2.4 mL

Part 7 12 PACKET

16.8 mL

Part 1 of 7**FIRST AID BURN**

benzalkonium chloride, lidocaine hydrochloride cream

Product Information

Item Code (Source)	NDC:0498-0903
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	0.13 g in 100 g
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE	0.5 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
ALOE VERA LEAF (UNII: ZY81Z83H0X)	
WATER (UNII: 059QF0KO0R)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
PEG-100 STEARATE (UNII: YD01N1999R)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
TROLAMINE (UNII: 9O3K93S3TK)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
DIAZOLIDINYL UREA (UNII: H5RIZ3MPW4)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		0.9 g in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/20/2017	

Part 2 of 7

EYESALINE EMERGENCY EYEWASH

purified water liquid

Product Information

Item Code (Source) NDC:0498-0100

Route of Administration OPHTHALMIC

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
WATER (UNII: 059QF0KO0R) (WATER - UNII:059QF0KO0R)	WATER	98.6 mL in 100 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM PHOSPHATE, MONOBASIC, MONOHYDRATE (UNII: 593YOG76RN)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM PHOSPHATE, DIBASIC (UNII: GR686LBA74)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0100-02	118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M018	12/18/2018	

Part 3 of 7

ALCOHOL WIPE

isopropyl alcohol swab

Product Information**Item Code (Source)** NDC:0498-0143**Route of Administration** TOPICAL**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	0.7 mL in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0143-04	0.4 mL in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	

Part 4 of 7**ASPIRIN**

aspirin tablet

Product Information**Item Code (Source)** NDC:0498-0114**Route of Administration** ORAL**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE 2208 (100 MPA.S) (UNII: B1QE5P712K)	
MINERAL OIL (UNII: T5L8T28FGP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	FR21
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0114-01	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	

Part 5 of 7

NEOMYCIN

antibiotic ointment

Product Information

Item Code (Source)	NDC:0498-0730
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN SULFATE	3.5 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
PETROLATUM (UNII: 4T6H12BN9U)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0730-01	0.9 g in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		03/31/2010	

Part 6 of 7**STING RELIEF PAD**

ethyl alcohol, lidocaine swab

Product Information

Item Code (Source)	NDC:0498-0733
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE ANHYDROUS	20 mg in 1 mL
ALCOHOL (UNII: 3K9958V90M) (ALCOHOL - UNII:3K9958V90M)	ALCOHOL	0.5 mL in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
MENTHOL (UNII: L7T10EIP3A)	

BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0733-00	0.4 mL in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/23/2017	

Part 7 of 7

ANTISEPTIC TOWELETTE

benzalkonium chloride liquid

Product Information

Item Code (Source) NDC:0498-0501

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	1.3 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0501-00	1.4 mL in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/22/2017	
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		10/18/2018	

Labeler - Honeywell Safety Products USA, INC (118768815)

Revised: 1/2024

Honeywell Safety Products USA, INC