

**MEDI-FIRST PEP-T-MED- bismuth subsalicylate tablet, chewable**  
**Unifirst First Aid Corporation**

-----  
**MEDI-FIRST PEP-T-MED**

***Drug Facts***

***Active ingredient (in each tablet)***

Bismuth Subsalicylate 262 mg  
(each tablet contains 102 mg salicylate)

***Purpose***

Upset stomach reliever/antidiarrheal

***Uses***

relieves

- traveler's diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
  - indigestion
  - nausea
  - gas
  - belching
  - fullness

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are:**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use if you have**

- bloody or black stool
- an ulcer
- a bleeding problem

**Ask a doctor before use if you have**

- fever
- mucus in the stool

**Ask a doctor or pharmacist if you are taking any drug for**

- anticoagulation (thinning of the blood)
- diabetes

- gout
- arthritis

**When using this product** a temporary and harmless darkening of the tongue and/or stool may occur.

**Stop use and ask a doctor if**

- symptoms get worse or lasts for more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

In case of accidental overdose, contact a physician or poison control center right away (1-800-222-1222).

**Directions**

- chew or dissolve in mouth
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Adults and children: (12 years and over)**

- chew 2 tablets every 1/2 to 1 hour as needed
- do not exceed 16 tablets in 24 hours
- use until diarrhea stops, but not more than 2 days

**Children under 12 years:** Ask a doctor

**Children under 12 years:** Do not give to children under 12 years of age.

**Other information**

- **phenylketonurics:** contains phenylalanine 1.1mg per tablet
- **calcium** content per tablet: 73 mg
- store at room temperature 59°-86°F (15°-30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

**Inactive ingredients**

acacia gum, aspartame, calcium carbonate, D&C red #27 aluminum lake, dextrans, magnesium stearate, maltodextrin, microcrystalline cellulose, peppermint flavoring, silicon dioxide

**Questions or comments?**

1-800-634-7680

**Medi-First Pep-T-Med Label**

36 Chewable Tablets

Medi-First®

PEP-T-MED

## Chewable Tablets

Compare active ingredients to:

Pepto Bismol®

Registered Trademark of Proctor & Gamble

*Upset Stomach/Diarrhea Relief • Bismuth Subsalicylate 262mg*

Chew or crush tablets completely before swallowing.

Do not swallow tablets whole.

Tamper Evident

**MEDI-FIRST®**  
**Pep-T-Med Chewable Tablets**

41220BOX  
Rev. 6/8/21  
Mfg. 1

36 Chewable Tablets

**Drug Facts**  
**Active ingredient (in each tablet)** Bismuth Subsalicylate 262 mg .....Upset stomach reliever/antidiarrheal (each tablet contains 102 mg salicylate) **Purpose**

**Uses**  
relieves ■ travelers' diarrhea ■ diarrhea  
■ upset stomach due to overindulgence in food and drink, including:  
■ heartburn ■ indigestion ■ nausea  
■ gas ■ belching ■ fullness

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome a rare but serious illness.

**Allergy alert:** Contains salicylate. Do not take if you are: ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

**Do not use if you have**  
■ bloody or black stool ■ an ulcer ■ a bleeding problem

**Ask a doctor before use if you have** ■ fever ■ mucus in the stool

**Ask a doctor or pharmacist if you are taking any drug for**  
■ anticoagulation (thinning of the blood) ■ diabetes ■ gout ■ arthritis

**When using this product** a temporary and harmless darkening of the tongue and/or stool may occur.

**Stop use and ask a doctor if**  
■ symptoms get worse or lasts for more than 2 days  
■ ringing in the ears or loss of hearing occurs  
■ diarrhea lasts more than 2 days

**Drug Facts (continued)**  
**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children.** In case of accidental overdose, contact a physician or poison control center right away (1-800-222-1222).

**Directions**  
■ chew or dissolve in mouth  
■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Adults and children (12 years and over)**  
■ chew 2 tablets every 1/2 to 1 hour as needed  
■ do not exceed 16 tablets in 24 hours  
■ use until diarrhea stops, but not more than 2 days

**Children under 12 years:** Ask a doctor

**Other information**  
■ phenylketonurics: contains phenylalanine 1.1 mg per tablet  
■ calcium content per tablet: 73 mg  
■ store at room temperature 59°-86°F (15°-30°C)  
■ tamper-evident sealed packets  
■ do not use any opened or torn packets  
(continued on opposite panel)

**Drug Facts (continued)**  
**Inactive ingredients**  
acacia gum, aspartame, calcium carbonate, D&C red #27 aluminum lake, dextrates, magnesium stearate, maltodextrin, microcrystalline cellulose, peppermint flavoring, silicon dioxide

**Questions or comments?**  
800-634-7680

**Compare active ingredient to:**  
**Pepto-Bismol®**  
Registered Trademark of Proctor & Gamble

✓ Upset Stomach / Diarrhea Relief • Bismuth Subsalicylate 262 mg

**Tamper Evident**

Reorder #41220  
Manufactured for  
Medique Products  
Fort Myers, FL 33967  
1-800-634-7680  
www.mediqueproducts.com

3 47682 41220 3

## MEDI-FIRST PEP-T-MED

bismuth subsalicylate tablet, chewable

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:47682-406
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (SALICYLIC ACID -	BISMUTH	262 mg

UNII:O414PZ4LPZ)

SUBSALICYLATE

202 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>ASPARTAME</b> (UNII: Z0H242BBR1)	
<b>CALCIUM CARBONATE</b> (UNII: H0G9379FGK)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>D&amp;C RED NO. 27</b> (UNII: 2LRS185U6K)	
<b>DEXTRATES</b> (UNII: G263MI44RU)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>ACACIA</b> (UNII: 5C5403N26O)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	

**Product Characteristics**

<b>Color</b>	pink	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	16mm
<b>Flavor</b>	PEPPERMINT	<b>Imprint Code</b>	RH;046
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-406-20	6 in 1 BOX	12/01/2011	
1	NDC:47682-406-69	6 in 1 PACKET; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	12/01/2011	

**Labeler** - Unifirst First Aid Corporation (832947092)

Revised: 9/2025

Unifirst First Aid Corporation