

MEDI-FIRST PEP-T-MED- bismuth subsalicylate tablet, chewable
Unifirst First Aid Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

MEDI-FIRST PEP-T-MED

Drug Facts

Active ingredient (in each tablet)

Bismuth Subsalicylate 262 mg
(each tablet contains 102 mg salicylate)

Purpose

Upset stomach reliever/antidiarrheal

Uses

relieves

- traveler's diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
 - indigestion
 - nausea
 - gas
 - belching
 - fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are:

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- bloody or black stool
- an ulcer
- a bleeding problem

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist if you are taking any drug for

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

When using this product a temporary and harmless darkening of the tongue and/or stool may occur.

Stop use and ask a doctor if

- symptoms get worse or lasts for more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, contact a physician or poison control center right away (1-800-222-1222).

Directions

- chew or dissolve in mouth
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Adults and children: (12 years and over)

- chew 2 tablets every 1/2 to 1 hour as needed
- do not exceed 16 tablets in 24 hours
- use until diarrhea stops, but not more than 2 days

Children under 12 years: Ask a doctor

Children under 12 years: Do not give to children under 12 years of age.

Other information

- **phenylketonurics:** contains phenylalanine 1.1mg per tablet
- **calcium** content per tablet: 73 mg
- store at room temperature 59°-86°F (15°-30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

acacia gum, aspartame, calcium carbonate, D&C red #27 aluminum lake, dextrans, magnesium stearate, maltodextrin, microcrystalline cellulose, peppermint flavoring, silicon dioxide

Questions or comments?

1-800-634-7680

Medi-First Pep-T-Med Label

36 Chewable Tablets

Medi-First®

PEP-T-MED

Chewable Tablets

Compare active ingredients to:

Pepto Bismol®

Registered Trademark of Proctor & Gamble

Upset Stomach/Diarrhea Relief • Bismuth Subsalicylate 262mg

Chew or crush tablets completely before swallowing.

Do not swallow tablets whole.

Tamper Evident

41220BOX
Rev. 6/8/21
Mfg. 1

36
Chewable
Tablets

Drug Facts
Active ingredient (in each tablet) Purpose
Bismuth Subsalicylate 262 mgUpset stomach reliever/antidiarrheal
(each tablet contains 102 mg salicylate)

Uses
relieves ■ travelers' diarrhea ■ diarrhea
■ upset stomach due to overindulgence in food and drink, including:
■ heartburn ■ indigestion ■ nausea
■ gas ■ belching ■ fullness

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are: ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

Do not use if you have
■ bloody or black stool ■ an ulcer ■ a bleeding problem

Ask a doctor before use if you have ■ fever ■ mucus in the stool

Ask a doctor or pharmacist if you are taking any drug for
■ anticoagulation (thinning of the blood) ■ diabetes ■ gout ■ arthritis

When using this product a temporary and harmless darkening of the tongue and/or stool may occur.

Stop use and ask a doctor if
■ symptoms get worse or lasts for more than 2 days
■ ringing in the ears or loss of hearing occurs
■ diarrhea lasts more than 2 days

Drug Facts (continued)
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of accidental overdose, contact a physician or poison control center right away (1-800-222-1222).

Directions
■ chew or dissolve in mouth
■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Adults and children: (12 years and over)
■ chew 2 tablets every 1/2 to 1 hour as needed
■ do not exceed 16 tablets in 24 hours
■ use until diarrhea stops, but not more than 2 days

Children under 12 years: Ask a doctor

Other information
■ phenylketonurics: contains phenylalanine 1.1 mg per tablet
■ calcium content per tablet: 73 mg
■ store at room temperature 59°-86°F (15°-30°C)
■ tamper-evident sealed packets
■ do not use any opened or torn packets
(continued on opposite panel)

Drug Facts (continued)
Inactive ingredients
acacia gum, aspartame, calcium carbonate, D&C red #27 aluminum lake, dextrans, magnesium stearate, maltodextrin, microcrystalline cellulose, peppermint flavoring, silicon dioxide

Questions or comments?
800-634-7680

Compare active ingredient to:
Pepto-Bismol®
Registered Trademark of Proctor & Gamble

✓ Upset Stomach / Diarrhea Relief • Bismuth Subsalicylate 262mg

Tamper Evident

Reorder #41220
Manufactured for
Medique Products
Fort Myers, FL 33967
1-800-634-7680
www.mediqueproducts.com

3 47682 41220 3

MEDI-FIRST PEP-T-MED

bismuth subsalicylate tablet, chewable

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:47682-406 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|--------------------------|-----------------|
| BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ) | BISMUTH SUBSALICYLATE | 262 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|-----------------|
| ASPARTAME (UNII: Z0H242BBR1) | |
| CALCIUM CARBONATE (UNII: H0G9379FGK) | |
| CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) | |
| D&C RED NO. 27 (UNII: 2LRS185U6K) | |
| DEXTRATES (UNII: G263MI44RU) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| ACACIA (UNII: 5C5403N26O) | |
| MALTODEXTRIN (UNII: 7CVR7L4A2D) | |

Product Characteristics

| | | | |
|-----------------|------------|---------------------|----------|
| Color | pink | Score | no score |
| Shape | ROUND | Size | 16mm |
| Flavor | PEPPERMINT | Imprint Code | RH;046 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|----------|------------------|--|-----------------------------|---------------------------|
| 1 | NDC:47682-406-20 | 6 in 1 BOX | 12/01/2011 | |
| 1 | NDC:47682-406-69 | 6 in 1 PACKET; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------------|---|-----------------------------|---------------------------|
| OTC monograph final | part335 | 12/01/2011 | |

Labeler - Unifirst First Aid Corporation (832947092)

Revised: 3/2022

Unifirst First Aid Corporation