

DYE FREE IBUPROFEN- ibuprofen tablet, film coated
Walgreen Company

Walgreens 44-438-Motrin

Active ingredient (in each white tablet)

Ibuprofen USP, 200 mg (NSAID)*
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - backache
 - the common cold
 - minor pain of arthritis
 - toothache
 - menstrual cramps
 - muscular aches
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- rash
- facial swelling
- asthma (wheezing)
- hives
- skin reddening
- shock
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - slurred speech
 - leg swelling
 - trouble breathing
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
 - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- see end flap for expiration date and lot number

Inactive ingredients

colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, sodium starch glycolate, stearic acid, triacetin

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

Walgreens

WALGREENS PHARMACIST RECOMMENDED†

Compare to the active ingredient
in Motrin® IB Tablets††

NDC 0363-4381-12

Dye-Free

Ibuprofen 200

IBUPROFEN TABLETS USP, 200 MG / PAIN RELIEVER / FEVER REDUCER **(NSAID)**

100

TABLETS

ACTUAL SIZE

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

DISTRIBUTED BY: **WALGREEN CO.**
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2021 Walgreen Co.

†Our pharmacists recommend the Walgreens brand.
We invite you to compare to national brands.

††This product is not manufactured or distributed by
Johnson & Johnson Corporation, owner of the
registered trademark Motrin® IB Tablets.

50844 ORG122143812

Do not print/Varnish

B-2201-438-12-HR
ORG122143812



Compare to the active ingredient in Motrin® IB Tablets*

Dye-Free

Ibuprofen 200

IBUPROFEN TABLETS USP, 200 mg / PAIN RELIEVER / FEVER REDUCER (NSAID)

100
TABLETS



ACTUAL SIZE

NDC 0363-4381-12

W3ORG0122-F
FEV0422

ITEM 500731 W0000-0000-0
3 1 1917108932 4

TAMPER EVIDENT: DO NOT USE IF
PRINTED SAFETY SEAL UNDER
CAP IS BROKEN OR MISSING

how2recycle.info

PAPER BOX
PLASTIC BOTTLE

Drug Facts (continued)

■ redness or swelling is present in the painful area
■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ do not take more than directed
■ the smallest effective dose should be used
■ adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
■ if pain or fever does not respond to 1 tablet, 2 tablets may be used
■ do not exceed 6 tablets in 24 hours, unless directed by a doctor
■ children under 12 years: ask a doctor

Other information

■ store between 20-25°C (68-77°F)
■ avoid excessive heat 40°C (104°F)
■ see end flap for expiration date and lot number

Inactive ingredients colloidal silicon dioxide, corn starch, hydroxymethylcellulose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, sodium starch glycolate, stearic acid, triacetin

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ right before or after heart surgery

Ask a doctor before use if ■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn ■ you are taking a diuretic
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
■ you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint
■ have bloody or black stools ■ vomit blood
■ you have symptoms of heart problems or stroke: ■ chest pain ■ slurred speech
■ leg swelling ■ trouble breathing
■ weakness in one part or side of body
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days

Drug Facts (continued)

Questions or comments?
Call 1-800-428-9391 8:30 AM-4:00 PM ET,
Monday-Friday

DISTRIBUTED BY: WALGREEN CO.,
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2021 Walgreen Co.
Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Motrin® IB Tablets.
50844 ORG122143812

Drug Facts

Active ingredient
(in each white tablet)

Ibuprofen USP, 200 mg (NSAID) for pain relief, fever reduction, and inflammation relief. Pain reliever/fever reducer.

Uses

■ temporarily relieves minor aches and pains due to:
■ headache ■ toothache
■ backache ■ menstrual cramps
■ the common cold ■ muscular aches
■ minor pain of arthritis
■ temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:
■ rash ■ facial swelling ■ asthma (wheezing)
■ hives ■ skin reddening ■ shock ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
■ take more or for a longer time than directed
■ take a blood thinning (anticoagulant) or steroid drug
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ have 3 or more alcoholic drinks every day while using this product
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Purpose

NDC 0363-4381-12

DYE FREE IBUPROFEN			
ibuprofen tablet, film coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-4381
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;438
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-4381-12	1 in 1 CARTON	03/01/1999	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075139	03/01/1999	

Labeler - Walgreen Company (008965063)**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(0363-4381)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(0363-4381)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(0363-4381)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(0363-4381)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(0363-4381)

Revised: 5/2023

Walgreen Company