

**4342 FIRST AID KIT- 4342 first aid kit
Honeywell Safety Products USA, INC**

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

0498-4342: First Aid Kit (BZK wipe, neomycin, alcohol wipes, sting relief, ASA, aypanal- Z019845)

Alcohol Wipe
Active ingredient

Isopropyl alcohol 70%

Alcohol Wipe
Purpose

First aid antiseptic

Alcohol Wipe
Uses

- first aid to help prevent infection in minor cuts, scrapes, and burns

Alcohol Wipe
Warnings

For external use only

Do not use

- in the eyes
- over large areas of the body

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burn

When using this product

- do not use longer than one week unless directed by a doctor

Stop use and consult a doctor

- if condition persists or gets worse

Keep out of reach of children

- If swallowed, get medical help or contact a Poison Control Center right away.

Alcohol Wipe***Directions***

- clean the affected area
- apply wipe to affected area 1 to 3 times daily
- may be covered with a sterile bandage
- discard wipe after single use

Alcohol Wipe***Other information***

store at room temperature 15 ° to 25 ° C (59 ° to 77 °F)

Alcohol Wipe***Inactive ingredient***

water

Alcohol Wipe***Questions***

1-800-430-5490

Aspirin***Active ingredient (in each tablet)***

Aspirin 325 mg (NSAID)* *nonsteroidal anti-inflammatory drug

Aspirin***Purpose***

Pain reliever/fever reducer

Aspirin***Uses***

temporarily reduces fever and relieves minor aches and pains associated with:

- a cold
- headache
- toothache
- muscular aches
- backache
- minor pain of arthritis
- premenstrual and menstrual periods

Aspirin

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:are:

- age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- ringing in the ears or loss of hearing occurs
- any new symptoms appear

If pregnant or breast-feeding,

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last three months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

- In case of overdose, get medical help or contact Poison Control Center right away.

Aspirin

Directions

- drink a full glass of water with each dose
- adults and children 12 years of age and older: take 1 or 2 tablets every 4 hours while symptoms last, not more than 12 tablets in 24 hours
- children under 12 years of age: consult a doctor

Aspirin

Other information

- store at room temperature 15° - 30°C (59° - 86°F)
- TAMPER EVIDENT PACKETS
- DO NOT USE IF OPEN OR TORN

Aspirin

Inactive ingredients

corn starch, croscarmellose sodium*, hypromellose*, microcrystalline cellulose*, mineral oil*, polyethylene glycol*, povidone, propylene glycol, silicon dioxide, stearic acid*, titanium dioxide*

*may contain these ingredients

Aspirin

Questions or Comments

1-800-430-5490

Sting Relief

Active ingredient

Ethyl alcohol 50.0%

Lidocaine HCl 2.0%

Sting Relief

Purpose

Antiseptic

Topical pain relief

Sting Relief

Uses

- prevent infection in minor scrapes, and temporary relief of itching of insect bites

Sting Relief

Warnings

For external use only

Flammable, keep away from open fire or flame

Do not use

- over large areas of the body
- in eyes
- over raw or blistered areas

Stop use and ask a doctor

- if conditions worsen or persist for more than 7 days or clear up and occur again within a few days

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

sting Relief

Directions

- adults and children 2 years and older: Apply to cleaned affected area not more than 3 times daily.
- children under 2 years of age: consult a doctor.

Sting Relief

Inactive ingredients

benzalkonium chloride, menthol, and purified water

Sting Relief

Questions or Comments?

1-800-430-5490

BZK

Active ingredient

Benzalkonium chloride 0.13% w/v

BZK

Purpose

First aid antiseptic

BZK

Uses

Antiseptic cleansing of face, hands, and body without soap and water

BZK

Warnings

For external use only

Do not use

- in the eyes or over large areas of the body
- on mucous membranes
- on irritated skin
- in case of deep puncture wounds, animal bites or serious burns, consult a doctor
- longer than 1 week unless directed by a doctor

Stop use and ask a doctor if

- if irritation, redness or other symptoms develop
- the condition persists or gets worse

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

BZK

Directions

- tear open packet and use as a washcloth

BZK

Other information

- store at room temperature 15 ° to 30 ° C (59 ° - 86 ° F)
- do not reuse towelette

BZK

Inactive ingredient

water

BZK

Questions

1-800-430-5490

Neomycin

Active ingredient (each gram contains)

Neomycin sulfate (5 mg equivalent to 3.5 mg Neomycin base)

Neomycin

Purpose

First aid antibiotic

Neomycin

Uses

- first aid to help prevent infection in
- minor cuts
- scrapes
- burns

Do not use

- in the eyes
- over large areas of the body

Neomycin

Warnings

For external use only

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Stop use and ask a doctor if

- the condition persists or gets worse
- a rash or other allergic reaction develops
- you need to use longer than 1 week

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away

Neomycin

Directions

- clean the affected area
- apply a small amount of the product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage

Neomycin

Other information

- store at 15⁰ to 25⁰ C (59⁰ to 77⁰ F)

Neomycin

Inactive ingredient

petrolatum

Neomycin

Questions

1-800-430-5490

Aypanal EX

Active ingredient

Acetaminophen 500 mg

Aypanal EX

PURpose

Pain reliever/fever reducer

Aypanal EX

Uses

- temporarily relieves minor aches and pains due to the common cold and headache - temporarily reduces fever

Aypanal EX

Warnings

Liver Warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount.
- with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If skin reaction occurs, stop use and seek medical help right away

Do Not Use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

- liver disease

Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present
- If pregnant or breastfeeding,
- ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Aypanal EX

Directions

- **do not take more than directed (see overdose warning)**
- adults and children 12 years of age and over: Take 2 tablets with water every 6 hours while symptoms last.
- do not take any more than 8 tablets in 24 hours.
- children under 12: consult a doctor

Aypanal EX

Other information

- store at room temperature 15^o -30^o C (59^o -86^o F)
- TAMPER EVIDENT- DO NOT USE IF OPEN OR TORN

Aypanal EX

Inactive ingredients

microcrystalline cellulose, povidone, sodium starch glycolate, starch, stearic acid

Aypanal EX

Questions or Comments?

1-800-430-5490


4342

Z019845 KIT CONTENTS

- 1 XTRA LRG 2" X 4" AWC
- 1 FIRST AID GUIDE ASHI
- LBL STOCK 6-3/8"X4"
- LBL STOCK 4"X2-7/8"
- 1 LBL STOCK 4"X2-7/8"
- 1 LBL STOCK 3"x1-7/8"
- 3 BZK ANTISEPTIC WIPE, BULK
- 1 PR LRG NITRILE GLVES ZIP BAG
- 1 LIGHT STICKS 6" 12 HR
- 1 TAPE ADHESIVE 1 X 2.5 125135
- 2 POUCH NEOMYCIN ANTIBIOTIC .9 G
- 6 WIPE ALCOHOL PREP IPA 70% (DUKAL)
- 1 KIT, PP 10 UNIT FA
- 1 SAFETEC STING RELIEF WIPES BULK
- 10 PLASTIC BANDAGE 3/4" X 3"
- 2 WOVEN FINGERTIP BANDAGE 2"
- 1 WOVEN KNUCKLE BANDAGE
- 1 ASPIRIN BULK 2/PK
- 1 AYPANAL EXTRA BULK 2/PK
- 1 BANDAGE, PATCH, ADHESIVE, 2X3

Alcohol Wipe Label

825362T Rev. A
 white printed
 Pms 072 blue, Pms 186 red,
 Pms 7459 blue and black



Aspirin Principal Display Panel




Sting Relief Principal Display Panel

825366 Rev B

Honeywell

Honeywell 032043P




Sting Relief Wipes

Use for:
Minor Cuts • Scrapes • Insect Bites

Single Use Pouches
Saturated Wipes

100 wipes

Honeywell 032043P



Sting Relief Wipes

Use for:
Minor Cuts • Scrapes • Insect Bites

Single Use Pouches
Saturated Wipes

100 wipes

Drug Facts

Active Ingredients	Purpose
Ethyl alcohol 50.0%	First aid antiseptic
Lidocaine HCl 2.0%	Topical analgesic

Uses First aid to help prevent infection in minor scrapes and temporary relief of itching of insect bites.

Warnings
For external use only.
Flammable, keep away from fire or flame.

Do not use • over large areas of the body • in eyes • over raw or blistered areas

Stop use and ask a doctor • if conditions worsen or persist for more than 7 days or clear up and occur again within a few days.

Keep out of reach of children.
If swallowed get medical help or contact Poison Control center right away.

Directions
• adults and children 2 years and older, apply to cleaned affected area not more than 3 times daily.
• children under 2 years of age, consult a doctor.


Inactive Ingredients
benzalkonium chloride, menthol, purified water

Questions or comments? 1-800-430-5490

Distributed by
Honeywell Safety
Products, USA, Inc.
Smithfield, RI 02917

www.honeywellsafety.com

USA
1-800-430-5490
0-401-943-4400



8 21812 01284 1

Made in USA
Packaged in Mexico

BZK Principal Display Panel

47001083
Rev B

Antiseptic Towelettes

Honeywell

02-16-35MD

Antiseptic Towelettes

Benzalkonium chloride
First aid antiseptic

Six-Saturated Towelettes

Distributed by
Honeywell Safety
Products USA, Inc.
Smithfield, RI 02917

47001083
Rev B

Antiseptic Towelettes

Honeywell

Drug Facts

Active Ingredient	Purpose
Benzalkonium chloride 0.133% w/v	First aid antiseptic

Uses • antiseptic cleaning of face, hands and body without soap and water. • air dries in seconds

Warnings

For external use only

When using this product • do not use in the eyes or apply over large areas of the body

Ask a doctor before use • in case of deep or puncture wounds, animal bites, or serious burns

Stop use and consult a doctor if

• irritation, redness or other symptoms develop • condition persists or gets worse

Do not use • longer than 1 week unless directed by doctor

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions • tear open packet, unfold and use as washcloth

Other information



• store at room temperature 15° -30° C (59° -86° F) • do not reuse towelette

Inactive ingredient water

Questions or comments 1-800-430-5490

Neomycin
Principal Display Panel

796041-25 Rev A Unit Carton Printing Plate for "C" size carton.



FIRST AID

VEHICLE, BULK
5 PERSON



GRAINGER
FOR THE ONES WHO GET IT DONE

GRAINGER.COM®

470017126A

Distributed by Honeywell Safety Products USA, Inc. Smithfield, RI 02917

4342 FIRST AID KIT

4342 first aid kit kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0498-4342
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-4342-01	1 in 1 KIT; Type 0: Not a Combination Product	10/18/2018	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 PACKET	2
Part 2	6 POUCH	2.4 mL
Part 3	1 PACKET	2
Part 4	2 PACKET	1.8 g
Part 5	3 PACKET	4.2 mL
Part 6	1 POUCH	0.4 mL

Part 1 of 6

AYPANAL EX

acetaminophen tablet

Product Information

Item Code (Source) NDC:0498-2110

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE (UNII: FZ989GH94E)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	FR1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-2110-01	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		01/02/2017	

Part 2 of 6

ALCOHOL WIPE

isopropyl alcohol swab

Product Information

Item Code (Source) NDC:0498-0143

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	0.7 mL in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0143-04	0.4 mL in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	

Part 3 of 6

ASPIRIN

aspirin tablet

Product Information

Item Code (Source) NDC:0498-0114

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE 2208 (100 MPA.S) (UNII: B1QE5P712K)	
MINERAL OIL (UNII: T5L8T28FGP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	FR21
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0114-01	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	

Part 4 of 6**NEOMYCIN**

antibiotic ointment

Product Information

Item Code (Source)	NDC:0498-0730
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN SULFATE	3.5 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
PETROLATUM (UNII: 4T6H12BN9U)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0730-01	0.9 g in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		03/31/2010	

Part 5 of 6

ANTISEPTIC TOWELETTE

benzalkonium chloride liquid

Product Information

Item Code (Source)	NDC:0498-0501
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	1.3 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0501-00	1.4 mL in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/22/2017	

Part 6 of 6

STING RELIEF PAD

ethyl alcohol, lidocaine swab

Product Information

Item Code (Source)	NDC:0498-0733
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE ANHYDROUS	20 mg in 1 mL
ALCOHOL (UNII: 3K9958V90M) (ALCOHOL - UNII:3K9958V90M)	ALCOHOL	0.5 mL in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
MENTHOL (UNII: L7T10EIP3A)	
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0733-	0.4 mL in 1 POUCH; Type 0: Not a Combination		

00	Product		
----	---------	--	--

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/23/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		10/18/2018	

Labeler - Honeywell Safety Products USA, INC (118768815)

Revised: 1/2024

Honeywell Safety Products USA, INC