

REGULAR STRENGTH PINK BISMUTH- bismuth subsalicylate suspension
Raritan Pharmaceuticals Inc

DRx Choice Regular Strength Bismuth Subsalicylate 525 mg Drug Facts

Active ingredient (in each 30 mL dose)

Bismuth subsalicylate 525 mg

Purposes

Upset stomach reliever and antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea
- gas
- belching
- fullness

Warnings

Reye's syndrome:Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea or vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate.Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- shake well before use
- only use dose cup provided
- **adults and children 12 years and over:**(30 mL) 1 dose every 1/2 or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
- 30 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (240 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- **children under 12 years:**ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- **each 30 mL contains:**sodium 10 mg
- **salicylate 261 mg**
- low sodium
- sugar free
- store at room temperature
- protect from freezing
- avoid excessive heat (over 104°F 40°C)

Inactive ingredients

carboxymethylcellulose sodium, D&C Red No. 22, D&C Red No. 28, flavor, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum

Questions or comments?

1-866-467-2748

Do not use if imprinted shrink band is missing or broken.

Principal Display Panel

DRx CHOICE[®]

NDC# **68163-706-04**

***Compare to the active ingredient Pepto-Bismol[®] Regular Strength**

Regular Strength

Pink Bismuth

Bismuth Subsalicylate

Upset Stomach Reliever/Antidiarrheal

Relieves nausea, heartburn, indigestion, upset stomach, diarrhea

ORIGINAL FLAVOR

4 FL. OZ. (118 mL)

Manufactured by:

Raritan Pharmaceuticals

8 Joanna Court

East Brunswick, NJ 08816

DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN.

***This product is not manufactured or distributed by The Procter & Gamble, distributor of Pepto-Bismol[®] Regular Strength.**

Package Label for 118 mL

PEEL BACK FOR DRUG FACTS

DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN

DR CHOICE

*Compare to the active ingredient in Pepto-Bismol® Regular Strength

Regular Strength Pink Bismuth

Bismuth Subsalicylate Upset Stomach Reliever/Antidiarrheal

Relieves nausea, heartburn indigestion, upset stomach, diarrhea

ORIGINAL FLAVOR

4 FL OZ (118 mL)

NDC 68163-706-04

Drug Facts (continued)

Allergy alert: Contains salicylate. Do not take if you are allergic to salicylates (including aspirin) or taking other salicylate products

Do not use if you have an ulcer a bleeding problem bloody or black stool

Ask a doctor before use if you have fever mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for anticoagulation (thinning the blood) diabetes gout arthritis

Manufactured by:
Raritan Pharmaceuticals
8 Joanna Court,
East Brunswick, NJ 08816

70604DCLR



LOT:

UNVARNISHED AREA

EXP:

Drug Facts

Active ingredient (in each 30 mL dose) Bismuth subsalicylate 525 mg.....**Purposes** Upset stomach reliever and antidiarrheal

Uses relieves travelers' diarrhea diarrhea upset stomach due to overindulgence in food and drink, including: heartburn indigestion nausea gas belching fullness

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

*This product is not manufactured or distributed by Procter & Gamble, distributor of Pepto-Bismol® Regular Strength.

Drug Facts (continued)

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if symptoms get worse or last more than 2 days ringing in the ears or loss of hearing occurs diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222

Directions shake well before use only use dose cup provided adults and children 12 years and over: 30 mL (1 dose) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea 30mL (1dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) do not exceed 8 doses (240 mL) in 24 hours use until diarrhea stops but not more than 2 days children under 12 years: ask a doctor drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Drug Facts (continued)

Other information

- each 30 mL contains: sodium 10 mg
- salicylate 261 mg low sodium sugar free
- store at room temperature
- protect from freezing
- avoid excessive heat (more than 104°F or 40°C)

Inactive ingredients carboxymethylcellulose sodium, D&C Red No 22, D&C Red No. 28, flavor, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum

Questions or comments? 1-866-467-2748

STOP PEELING HERE

Package Label for 354 mL

Drug Facts (continued)

Directions ■ shake well before use ■ only use dose cup provided
 ■ **adults and children 12 years and over:**
 ■ 30 mL (1 dose) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea ■ 30 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) ■ do not exceed 8 doses (240 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ **children under 12 years:** ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information ■ each 30 mL contains: sodium 10 mg ■ salicylate 261 mg ■ low sodium ■ sugar free ■ store at room temperature ■ protect from freezing ■ avoid excessive heat (more than 104°F or 40°C)

Inactive ingredients carboxymethylcellulose sodium, D&C Red No. 22, D&C Red No. 28, flavor, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum

Questions or comments? 1-866-467-2748

Manufactured by: Raritan Pharmaceuticals
 8 Joanna Court, East Brunswick, NJ 08816

LOT:
 UNVARNISHED
 AREA
 EXP:



*Compare to the active ingredient in
 Pepto-Bismol® Regular Strength

Dr. Choice®

NDC 68163-706-12

regular strength
pink bismuth
Bismuth Subsalicylate /
Upset Stomach Reliever/Antidiarrheal
Relieves: Nausea, Heartburn, Indigestion,
Upset Stomach & Diarrhea
Original Flavor
12 FL. OZ. (354 mL)

Do not use if imprinted shrink band is missing or broken

Drug Facts

Active ingredient (in each 30 mL dose)	Purposes
Bismuth subsalicylate 525 mg	Upset stomach reliever and antidiarrheal

Uses relieves ■ traveler's diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including:
 ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are
 ■ allergic to salicylates (including aspirin)
 ■ taking other salicylate products

Do not use if you have ■ an ulcer ■ a bleeding problem ■ bloody or black stool

Ask a doctor before use if you have ■ fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for
 ■ anticoagulation (thinning the blood) ■ diabetes ■ gout ■ arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.

*This product is not manufactured or distributed by Procter & Gamble, distributor of Pepto-Bismol® Regular Strength.

REGULAR STRENGTH PINK BISMUTH

bismuth subsalicylate suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68163-706
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
D&C RED NO. 28 (UNII: 767IPOY5NH)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
POTASSIUM SORBATE (UNII: 1VPU26JZ4)	
WATER (UNII: 059QF0KO0R)	
SALICYLIC ACID (UNII: O414PZ4LPZ)	
DIMETHICONE, UNSPECIFIED (UNII: 92RU3N3Y1O)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

XANTHAN GUM (UNII: TTV12P4NEE)

Product Characteristics

Color	pink (viscous)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68163-706-04	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/13/2019	
2	NDC:68163-706-12	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/13/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	05/13/2019	

Labeler - Raritan Pharmaceuticals Inc (127602287)

Revised: 9/2025

Raritan Pharmaceuticals Inc