

PARASITEXX- pyrantel pamoate suspension
Dannso Corp./d.b.a. Essential Products

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

PARASITEXX

Active Ingredient

Active Ingredient (in each 1 ml) Purpose

Active Ingredient (in each 1 ml)	Purpose
PYRANTEL PAMOATE	Pinworm Treatment

Inactive Ingredient

acesulfame potassium, cherry flavor, citric acid, glycerin, polysorbate 80, potassium sorbate, propylene glycol, sodium benzoate, sodium carboxymethylcellulose, sodium citrate, sorbitol, water, xanthan gum.

Dosage

Directions Read package insert careful before taking this medication. Take only according to directions and do not exceed the recommended dosage unless directed by a doctor. Medication should only be taken one time as a single dose; do not repeat treatment unless directed by a doctor. When one individual in a household has pinworms, the entire household should be treated unless otherwise advised. See warnings. If any worms other than pinworms are present before or after treatment, or pinworms are still present after treatment, consult a doctor.

WEIGHTS SINGLE DOSE

Less than 26 pounds or under 2 years of age	Do not use unless directed by a doctor
25 to 37 pounds	½ teaspoonful (2.5 ml)
38 to 62 pounds	1 teaspoonful (5 ml)
63 to 87 pounds	1 ½ teaspoonfuls (7.5 ml)
88 to 112 pounds	2 teaspoonfuls (10 ml)
113 to 137 pounds	2 ½ teaspoonfuls (12.5 ml)
138 to 162 pounds	3 teaspoonfuls (15 ml)
163 to 187 pounds	3 ½ teaspoonfuls (17.5 ml)
188 pounds and over	4 teaspoonfuls (20 ml)

Shake well before use. This product can be taken any time of the day, with or without meals. It may be taken alone or with milk or juice. Use of a laxative is not necessary prior to, during, or after medication.

Uses

For the treatment of pinworms.

Warnings

Do not exceed recommended dosage. When using this product abdominal cramps, nausea, diarrhea, headache, or dizziness sometimes occurs after taking this drug. If this condition persists consult doctor.

Ask a doctor before use if you have liver disease.

If pregnant or breast-feeding, ask a health professional before use.

Uses for the treatment of pinworms

Keep out of reach of children.

In case of accidental overdose, seek medical attention or contact a Poison Control Center immediately.

Other information

Protect from excessive heat and freezing. Store at room temperature.

Tamper-Evident Feature: do not use if printed seal around bottle neck is torn, broken or missing.

Questions? 305-261-8762

Monday - Friday 9am to 5pm EST.

NDC 70242-134-05

Pyrantel Pamoate Suspension

Parasitexx

For the Treatment of Pinworms
(Pyrantel Base 50 mg/mL)

LIQUID/LIQUIDO

Shake Well / Agitar bien
Measuring Cup Included/Vaso Dosificador Incluido
Contents: 1 Oz FL (30mL)

Suspensión Pamoato de Pirantel
Para el Tratamiento de los Oxiuros
(Base de Pirantel 50 mg/mL)
Parasitexx

Contenido: 1 Oz FL (30mL)
MADE IN USA



Suspensión Pamoato de Pirantel
Para el Tratamiento de los Oxiuros
(Base de Pirantel 50 mg/mL)
Parasitexx

KEEP OUT OF REACH OF CHILDREN

Distributed by
essential Products
Miami, FL 33126
EssentialProductsUSA.com



MADE IN USA
Parasitexx

Parasitexx

LIQUID/LIQUIDO

Drug Facts	Factores Farmacológicos																																				
Active ingredient (in each 1 mL): Pyrantel Pamoate 144 mg (Pyrantel treatment equivalent to 50 mg pyrantel base)	Agente activo (en cada 1 mL): Pirantel Pamoato 144 mg (Tratamiento de los oxiuros equivalente a 50 mg de base de pirantel)																																				
Uses for the treatment of pinworms.	Usos para el tratamiento de los oxiuros.																																				
Warnings Do not exceed recommended dosage. When using this product additional cramps, nausea, diarrhea, headache, or dizziness sometimes occur after taking this drug. If any of these conditions persist consult a doctor.	Advertencias No exceder la dosis recomendada. Cuando se utiliza este producto a veces ocurren espasmos abdominales, náuseas, vómitos, diarrea, dolor de cabeza o mareos. Si alguno de estas condiciones persisten consulte a un médico.																																				
Ask a doctor before use if you have: liver disease.	Pregunte a su médico antes de usar si usted tiene: enfermedad del hígado.																																				
If pregnant or breast-feeding, ask a health professional before use.	Si está embarazada o lactando, pregunte a un profesional de la salud antes de usarlo.																																				
Keep out of reach of children. In case of accident or overdose, wash residual amount in contact with Poison Control Center immediately.	Mantener fuera del alcance de los niños. En caso de sobredosis accidental, busque ayuda médica o contacto a un Centro de Envenenamiento inmediatamente.																																				
Directions: Read package insert carefully before taking the medicine. Take only as directed and do not exceed the recommended dosage unless directed by a doctor. If a doctor of child only has taken one time as a single dose, do not repeat treatment unless directed by a doctor. When one individual in a household has pinworms, the rest of household should be treated unless otherwise advised. See Warnings. If any worms other than pinworms are present before or after treatment, or pinworms are still present after treatment, consult a doctor.	Instrucciones: Leer prospecto detalladamente antes de tomar este medicamento. Tome sólo de acuerdo a las instrucciones y no exceda la dosis recomendada a menos que lo indique un médico. El medicamento sólo debe tomarse una vez en una sola dosis; no repita el tratamiento a menos que lo indique un médico. Cuando una persona en la familia tiene oxiuros, toda la familia deberá ser tratada a menos que se le indique lo contrario. Véase advertencias. Si alguno de los parásitos que no son los oxiuros está presente antes o después del tratamiento, consulte a un médico.																																				
<table border="1"> <thead> <tr> <th>Weight</th> <th>Single Dose</th> </tr> </thead> <tbody> <tr> <td>Less than 20 pounds or under 2 years of age</td> <td>Do not use unless directed by a doctor</td> </tr> <tr> <td>20 to 24 pounds</td> <td>1 teaspoon (5 mL)</td> </tr> <tr> <td>25 to 34 pounds</td> <td>1 1/2 teaspoons (7.5 mL)</td> </tr> <tr> <td>35 to 44 pounds</td> <td>2 teaspoons (10 mL)</td> </tr> <tr> <td>45 to 54 pounds</td> <td>2 1/2 teaspoons (12.5 mL)</td> </tr> <tr> <td>55 to 64 pounds</td> <td>3 teaspoons (15 mL)</td> </tr> <tr> <td>65 to 74 pounds</td> <td>3 1/2 teaspoons (17.5 mL)</td> </tr> <tr> <td>75 to 84 pounds</td> <td>4 teaspoons (20 mL)</td> </tr> </tbody> </table>	Weight	Single Dose	Less than 20 pounds or under 2 years of age	Do not use unless directed by a doctor	20 to 24 pounds	1 teaspoon (5 mL)	25 to 34 pounds	1 1/2 teaspoons (7.5 mL)	35 to 44 pounds	2 teaspoons (10 mL)	45 to 54 pounds	2 1/2 teaspoons (12.5 mL)	55 to 64 pounds	3 teaspoons (15 mL)	65 to 74 pounds	3 1/2 teaspoons (17.5 mL)	75 to 84 pounds	4 teaspoons (20 mL)	<table border="1"> <thead> <tr> <th>Age</th> <th>Dose (Single)</th> </tr> </thead> <tbody> <tr> <td>Menos de 20 libras o menor de 2 años</td> <td>No utilizar. Consultar a un médico</td> </tr> <tr> <td>25 a 24 libras</td> <td>1 cucharadita (5 mL)</td> </tr> <tr> <td>25 a 34 libras</td> <td>1 1/2 cucharaditas (7.5 mL)</td> </tr> <tr> <td>35 a 44 libras</td> <td>2 cucharaditas (10 mL)</td> </tr> <tr> <td>45 a 54 libras</td> <td>2 1/2 cucharaditas (12.5 mL)</td> </tr> <tr> <td>55 a 64 libras</td> <td>3 cucharaditas (15 mL)</td> </tr> <tr> <td>65 a 74 libras</td> <td>3 1/2 cucharaditas (17.5 mL)</td> </tr> <tr> <td>75 libras o más</td> <td>4 cucharaditas (20 mL)</td> </tr> </tbody> </table>	Age	Dose (Single)	Menos de 20 libras o menor de 2 años	No utilizar. Consultar a un médico	25 a 24 libras	1 cucharadita (5 mL)	25 a 34 libras	1 1/2 cucharaditas (7.5 mL)	35 a 44 libras	2 cucharaditas (10 mL)	45 a 54 libras	2 1/2 cucharaditas (12.5 mL)	55 a 64 libras	3 cucharaditas (15 mL)	65 a 74 libras	3 1/2 cucharaditas (17.5 mL)	75 libras o más	4 cucharaditas (20 mL)
Weight	Single Dose																																				
Less than 20 pounds or under 2 years of age	Do not use unless directed by a doctor																																				
20 to 24 pounds	1 teaspoon (5 mL)																																				
25 to 34 pounds	1 1/2 teaspoons (7.5 mL)																																				
35 to 44 pounds	2 teaspoons (10 mL)																																				
45 to 54 pounds	2 1/2 teaspoons (12.5 mL)																																				
55 to 64 pounds	3 teaspoons (15 mL)																																				
65 to 74 pounds	3 1/2 teaspoons (17.5 mL)																																				
75 to 84 pounds	4 teaspoons (20 mL)																																				
Age	Dose (Single)																																				
Menos de 20 libras o menor de 2 años	No utilizar. Consultar a un médico																																				
25 a 24 libras	1 cucharadita (5 mL)																																				
25 a 34 libras	1 1/2 cucharaditas (7.5 mL)																																				
35 a 44 libras	2 cucharaditas (10 mL)																																				
45 a 54 libras	2 1/2 cucharaditas (12.5 mL)																																				
55 a 64 libras	3 cucharaditas (15 mL)																																				
65 a 74 libras	3 1/2 cucharaditas (17.5 mL)																																				
75 libras o más	4 cucharaditas (20 mL)																																				
Other information: Protect from excessive heat and freezing; store at room temperature. Tamper-Evident Feature: Do not use if printed seal around bottle neck is torn, broken, or missing.	Other information: Proteja de calor excesivo y de la congelación; Almacene a temperatura ambiente. Evite la exposición al frío. No use este producto si el sello impreso alrededor del cuello o la botella está roto, roto, o faltante.																																				
Active ingredients: acacia, calcium phosphate, cherry flavor, dicalcium phosphate, polyethylene glycol, polyethylene glycol, sodium benzoate, sodium carboxymethylcellulose, sodium citrate, sodium hydroxide, xanthan gum.	Agente(s) activo(s): sacarín, sabor de cereza, colorante blanco, ácido cítrico, glicerina, polietileno glicol, sorbitol de sodio, polietileno glicol, estearato de sodio, carboximetilcelulosa sódica, citrato de sodio, hidróxido de sodio, goma de xantano.																																				
Questions? 305-261-4762 Monday-Friday 9am to 5pm EST	Preguntas? 305-261-4762 Lunes-Viernes 9am a 5pm EST																																				

KEEP OUT OF REACH OF CHILDREN



PARASITEXX

pyrantel pamoate suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70242-134
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PYRANTEL PAMOATE (UNII: 81BK194Z5M) (PYRANTEL - UNII:4QIH0N49E7)	PYRANTEL	144 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
CHERRY (UNII: BUC5I9595W)	
CITRIC ACID ACETATE (UNII: DSO12WL7AU)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
WATER (UNII: 059QF0KO0R)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70242-134-05	1 in 1 CARTON	07/01/2017	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part357B	05/01/2017	

Labeler - Danso Corp./d.b.a. Essential Products (059741071)

Registrant - Danso Corp./d.b.a. Essential Products (059741071)

Establishment

Name	Address	ID/FEI	Business Operations
All Pharma LLC		117605075	MANUFACTURE(70242-134)

Revised: 1/2022

Danso Corp./d.b.a. Essential Products