

**NIGHTTIME COLD AND FLU RELIEF- acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled  
Harris Teeter, LLC**

-----  
**Safeway Nighttime Cold & Flu**

**Active Ingredients (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 15 mg

Doxylamine Succinate 6.25 mg

**Purpose**

**Pain reliever/fever reducer**

Cough suppressant

Antihistamine

**Purpose**

Pain reliever / fever reducer

Cough suppressant

Antihistamine

**Uses**

- temporarily relieves common cold and flu symptoms:
  - sore throat
  - headache
  - minor aches and pains
  - fever
  - runny nose and sneezing
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough that such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

### **Ask a doctor of pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1888-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use

### **Other Information**

- store between 15-30°C (59-86°F)
- avoid excessive heat

### **Inactive Ingredients**

D&C yellow #10, FD&C blue #1, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

### **Questions or Comments?**

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

### **Principal Display Panel**

†Compare to the active ingredients in Vick® NyQuil® Cold & Flu LiquiCaps™

Nighttime

Cold & Flu Relief

MULTI-SYMPTOM RELIEF

**ACETAMINOPHEN** 325 mg

Pain Reliever / Fever Reducer

DEXTROMETHORPHAN HBr 15 mg • Cough Suppressant

DOXYLAMINE SUCCINATE 6.25 mg • Antihistamine

COLD & FLU

NightTime Relief

- Aches, Fever, SoreThroat
- Cough
- Runny Nose, Sneezing

ALCOHOL-FREE

SOFTGELS\*\*

(\*\*Liquid Filled Capsules)

†This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, NyQuil™, and LiquiCaps™ are registered trademarks of The Procter & Gamble Company

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

PROUDLY DISTRIBUTED BY:

HARRIS TEETER, LLC

MATTHEWS, NC 28105

### Package Label

PROUDLY DISTRIBUTED BY:  
HARRIS TEETER, LLC  
MATTHEWS, NC 28105

This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, NyQuil™, and LiquiCaps™ are registered trademarks of The Procter & Gamble Company.

**Parents:**  
Keep this and all other medications out of the reach of children. See important information about this product on page 2 of this carton.

Full details: 1-800-432-5111 or harrisiteeter.com.

FD-03839C    F3089702

0 720361 73617 8



## NightTime Cold & Flu Relief

**MULTI-SYMPTOM RELIEF**

**ACETAMINOPHEN 325 mg**  
Pain Reliever/Fever Reducer,  
**DEXTROMETHORPHAN HBr 15 mg**  
Cough Suppressant,  
**DOXYLAMINE SUCCINATE 6.25 mg**  
Antihistamine

**COLD & FLU**

NightTime Relief

- Aches, Fever, Sore Throat
- Cough
- Sneezing, Runny Nose

**ALCOHOL-FREE**

**24 Softgels™**  
(\*Liquid Filled Capsules)



**ACTUAL SIZE**



**Drug Facts**

**Active ingredients (in each softgel)**

Acetaminophen 325 mg.....Pain reliever/fever reducer  
Dextromethorphan HBr 15 mg.....Cough suppressant  
Doxylamine succinate 6.25 mg.....Antihistamine

**Purposes**

**Uses** ■ temporarily relieves common cold and flu symptoms: ■ sore throat ■ headache ■ minor aches and pains ■ fever ■ runny nose and sneezing ■ cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema ■ cough that occurs with too much phlegm (mucus)

**Drug Facts (continued)**

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product** ■ do not exceed recommended dosage ■ excitability may occur, especially in children ■ avoid alcoholic drinks ■ marked drowsiness may occur ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if** ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ do not take more than directed (see Overdose warning) ■ do not take more than 4 doses in 24 hours ■ adults and children 12 years and over: take 2 softgels with water every 6 hours ■ swallow whole; do not crush, chew, or dissolve ■ children under 12 years: do not use

**Other information** ■ store between 15-30°C (59-86°F) ■ avoid excessive heat

**Inactive ingredients** ■ D&C yellow #10, FD&C blue #1, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

**Questions or comments?**  
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

Lot No.:  
Exp. Date:

**HARRIS TEETER Nighttime Cold & flu Relief**

# NIGHTTIME COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69256-899
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>LIGHT MINERAL OIL</b> (UNII: N6K5787QVP)	

## Product Characteristics

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	PC10
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69256-899-24	24 in 1 CARTON	05/28/2021	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/28/2021	

**Labeler** - Harris Teeter, LLC (048463103)

Revised: 12/2025

Harris Teeter, LLC