

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 540- octinoxate and titanium dioxide emulsion
SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 550- octinoxate and titanium dioxide emulsion
SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 560- octinoxate and titanium dioxide emulsion
SHISEIDO AMERICAS CORPORATION

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION

Drug Facts

<i>Active ingredients</i>	<i>Purpose</i>
OCTINOXATE 6.9%	Sunscreen
TITANIUM DIOXIDE 1.0%	Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see ***Directions***), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. – 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses

Inactive Ingredients

WATER■DIMETHICONE■DIPHENYLSILOXY PHENYL
TRIMETHICONE■GLYCERIN■TRIFLUOROPROPYLDIMETHYL/TRIMETHYLSILOXYSILICATE■BUTYLENE

GLYCOL■HYDROGENATED POLYDECENE■PEG-10 DIMETHICONE■BIS-BUTYLDIMETHICONE POLYGLYCERYL-3■DISTEARDIMONIUM
HECTORITE■ERYTHRITOL■XYLITOL■SORBITAN
SESQUIISOSTEARATE■POLYQUATERNIUM-51■CITRUS UNSHIU PEEL EXTRACT■THYMUS SERPYLLUM EXTRACT■FAGUS SYLVATICA BUD EXTRACT■HYDROLYZED CONCHIOLIN PROTEIN■DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER■DIMETHICONE/PEG-10/15 CROSSPOLYMER■BARIUM SULFATE■SYNTHETIC FLUORPHLOGOPITE■ALUMINUM HYDROXIDE■POLYSILICONE-2■DISODIUM EDTA■STEARIC ACID■CALCIUM ALUMINUM BOROSILICATE■TOCOPHEROL■DIPROPYLENE
GLYCOL■TRIETHOXYCAPRYLYLSILANE■TRIETHOXSILYLETHYL
POLYDIMETHYLSILOXYETHYL
DIMETHICONE■BHT■ALUMINA■POLYMETHYLSILSESQUIOXANE■PHENOXYETHANOL■TITANIUM DIOXIDE■IRON OXIDES■MICA■

Other information

- protect this product in this container from excessive heat and direct sun.

Questions or comments?

Call toll free 1-800-906-7503

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 110 Alabaster

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- **Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

Drug Facts
(continued)

Inactive ingredients

WATER • DIMETHICONE •
C12-PHENYLSILOXY PHENYL TRIMETHICONE • GLYCERIN •
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSEBACATE •
BUTYLENE GLYCOL • HYDROGENATED POLYDECENE •
PEG-10 DIMETHICONE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
DISTEAROINDOLIUM HECTORITE • ERYTHRITOL • XYLITOL •
SORBITAN SESQUISOSEBATE • POLYQUATERNIUM-15 •
CITRUS UNSHNU PEEL EXTRACT •
THYMUS SERPYLLUM EXTRACT •
FAGUS SYLVATICA BUD EXTRACT •
HYDROLYZED CONCHILIN PROTEIN •
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER •
DIMETHICONE PEG-10/15 CROSSPOLYMER •
BARIUM SULFATE • SYNTHETIC FLUOROPHLOGHITE •
ALUMINUM HYDROXIDE • POLYSILOXANE-2 •
SODIUM EDTA • STEARIC ACID •
CALCIUM ALUMINUM BORSILICATE • TOCOPHEROL •
DIPROPYLENE GLYCOL • TRIETHOXYCAPRYLYLSILANE •
TRIMETHOXYSILYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE •
BHT • ALUMINA • POLYMETHYLSILOXANOLANE •
PHENOXYETHANOL • TITANIUM DIOXIDE • IRON OXIDES •
MICA •

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com



<http://s1872.com/16732AA>
See our website for more information.

30mL NET WT. 1.2 OZ.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16732 2
12711602



110 / Alabaster

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 120 Ivory

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

Drug Facts
(continued)

Inactive ingredients

WATER • DIMETHICONE •
COPHENYLSILOXY PHENYL TRIMETHICONE • GLYCERIN •
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE •
BUTYLENE GLYCOL • HYDROGENATED POLYDECENE •
PEG-10 DIMETHICONE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
DISTEAROINDIUM HECTORITE • ERYTHRITOL • XYLITOL •
SORBITAN SESQUISOOSTEARATE • POLYQUATERNIUM-15 •
CITRUS UNSHNU PEEL EXTRACT •
THYMUS SERPYLLUM EXTRACT •
FAGUS SYLVATICA BUD EXTRACT •
HYDROLYZED CONCHILIN PROTEIN •
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER •
DIMETHICONE/PEG-10/15 CROSSPOLYMER •
BARIUM SULFATE • SYNTHETIC FLUOROPHLOGHITE •
ALUMINUM HYDROXIDE • POLYSILOXANE-2 •
SODIUM EDTA • STEARIC ACID •
CALCIUM ALUMINUM BORSILICATE • TOCOPHEROL •
PROPYLENE GLYCOL • TRIETHOXYCAPRYLYLSILANE •
TRIMETHOXYSYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE •
BHT • ALLYLVA • POLYMETHYLSILOXOXANE •
PHENOXYETHANOL • TITANIUM DIOXIDE • IRON OXIDES •
MICA •

Other Information

protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com




<http://s1872.com/16732AA>
See our website for more information.

30mL NET WT. 1.2 OZ.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16733 9
12714960



120 / Ivory

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 130 Opal

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0% ...
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- **Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

Drug Facts
(continued)

Inactive ingredients

WATER • DIMETHICONE •
COPHENYLSILOXY PHENYL TRIMETHICONE • GLYCERIN •
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE •
BUTYLENE GLYCOL • HYDROGENATED POLYDECENE •
PEG-10 DIMETHICONE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
DISTEARDINOLIUM HECTORITE • ERYTHRITOL • XYLITOL •
SORBITAN SESQUISOOSTEATE • POLYQUATERNIUM-15 •
CITRUS UNSHNU PEEL EXTRACT •
THYMUS SERPYLLUM EXTRACT •
FAGUS SYLVATICA BUD EXTRACT •
HYDROLYZED CONCHILIN PROTEIN •
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER •
DIMETHICONE PEG-10/15 CROSSPOLYMER •
BARIUM SULFATE • SYNTHETIC FLUORPHLOGHITE •
ALUMINUM HYDROXIDE • POLYSILOXANE-2 •
SODIUM EDTA • STEARIC ACID •
CALCIUM ALUMINUM BORSILICATE • TOCOPHEROL •
DIPROPYLENE GLYCOL • TRIETHOXYCAPRYLYLSILANE •
TRIMETHOXYSYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE •
BHT • ALLYLVA • POLYMETHYLSILOXANOLANE •
PHENOXYETHANOL • TITANIUM DIOXIDE • IRON OXIDES •
MICA •

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com




<http://s1872.com/16732AA>
See our website for more information.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16734 6

12714962



130 / Opal

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 140 Porcelain

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- **Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

Drug Facts
(continued)

Inactive ingredients

WATER • DIMETHICONE •
COPHENYLSILOXY PHENYL TRIMETHICONE • GLYCERIN •
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE •
BUTYLENE GLYCOL • HYDROGENATED POLYDECENE •
PEG-10 DIMETHICONE •
BIS-BUTYLDIMETHICONE POLYBICYCLERYL-3 •
DISTEAROINDIUM HECTORITE • ERYTHRITOL • XYLITOL •
SORBITAN SESQUISOOSTEARATE • POLYQUATERNIUM-15 •
CITRUS UNSHNU PEEL EXTRACT •
THYMUS SERPYLLUM EXTRACT •
FAGUS SYLVATICA BUD EXTRACT •
HYDROLYZED CONCHILIN PROTEIN •
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER •
DIMETHICONE PEG-10/15 CROSSPOLYMER •
BARIUM SULFATE • SYNTHETIC FLUOROPHLOGHITE •
ALUMINUM HYDROXIDE • POLYSILOXANE-2 •
SODIUM EDTA • STEARIC ACID •
CALCIUM ALUMINUM BORSILICATE • TOCOPHEROL •
DIPROPYLENE GLYCOL • TRIETHOXYCAPRYLYLSILANE •
TRIMETHOXYSYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE •
BHT • ALLYLVA • POLYMETHYLSILOXANOLANE •
PHENOXYETHANOL • TITANIUM DIOXIDE • IRON OXIDES •
MICA •

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com




<http://s1872.com/16732AA>
See our website for more information.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16735 3
12714964



140 / Porcelain

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 150 Lace

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

Drug Facts
(continued)

Inactive ingredients

WATER - DIMETHICONE -
CYPHENYLSILOXY PHENYL TRIMETHICONE - GLYCERIN -
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE -
BUTYLENE GLYCOL - HYDROGENATED POLYDECENE -
PEG-10 DIMETHICONE -
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 -
DISTEARDINOLIUM HECTORITE - ERYTHRITOL - XYLITOL -
SORBITAN SESQUISOOSTEARATE - POLYQUATERMIUM-15 -
CITRUS UNSHNU PEEL EXTRACT -
THYMUS SERPYLLUM EXTRACT -
FAGUS SYLVATICA BUD EXTRACT -
HYDROLYZED CONCHILIN PROTEIN -
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER -
DIMETHICONE PEG-10/15 CROSSPOLYMER -
BARIUM SULFATE - SYNTHETIC FLUOROPHLOGHITE -
ALUMINUM HYDROXIDE - POLYSILOXANE-2 -
SODIUM EDTA - STEARIC ACID -
CALCIUM ALUMINUM BORSILICATE - TOCOPHEROL -
DIPROPYLENE GLYCOL - TRIETHOXYCAPRYLYLSILANE -
TRIMETHOXYSILYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE -
BHT - ALUMINA - POLYMETHYLSILOXOXANE -
PHENOXYETHANOL - TITANIUM DIOXIDE - IRON OXIDES -
MICA -

Other Information

protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com




<http://s1872.com/16732AA>
See our website for more information.

30mL NET WT. 1.2 OZ.

12713286
AAA

縮小サイズ0.65倍 6mm×6mm
11mm×26mm

7 30852 16736 0
12714966



150 / Lace

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 160 Shell

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients

Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

■ helps prevent sunburn
■ if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- **Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

Drug Facts
(continued)

Inactive ingredients

WATER • DIMETHICONE •
COPHENYLSILOXY PHENYL TRIMETHICONE • GLYCERIN •
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE •
BUTYLENE GLYCOL • HYDROGENATED POLYDECENE •
PEG-10 DIMETHICONE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
DISTEARDINOLAM HEPTOATE • ERYTHRITOL • XYLITOL •
SORBITAN SESQUISOOSTEATE • POLYQUATERNIUM-15 •
CITRUS UNSHNU PEEL EXTRACT •
THYMUS SERPYLLUM EXTRACT •
FAGUS SYLVATICA BUD EXTRACT •
HYDROLYZED CONCHILIN PROTEIN •
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER •
DIMETHICONE PEG-10/15 CROSSPOLYMER •
BARIUM SULFATE • SYNTHETIC FLUOROPHLOGHITE •
ALUMINUM HYDROXIDE • POLYSILOXANE-2 •
SODIUM EDTA • STEARIC ACID •
CALCIUM ALUMINUM BORSILICATE • TOCOPHEROL •
DIPROPYLENE GLYCOL • TRIETHOXYCAPRYLYLSILANE •
TRIMETHOXYSILYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE •
BHT • ALLYLVA • POLYMETHYLSILOXOXANE •
PHENOXYETHANOL • TITANIUM DIOXIDE • IRON OXIDES •
MICA •

Other Information

■ protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com



<http://s1872.com/16732AA>
See our website for more information.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16737 7
12714968



160 / Shell

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 210 Birch

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

Drug Facts
(continued)

Inactive ingredients

WATER • DIMETHICONE • C12-PHENYLSILOXY PHENYL TRIMETHICONE • GLYCERIN • TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE • BUTYLENE GLYCOL • HYDROGENATED POLYDECENE • PEG-10 DIMETHICONE • BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 • DISTEARDINOLIUM HECTORITE • ERYTHRITOL • XYLITOL • SORBITAN SESQUISOOSTEATE • POLYQUATERNIUM-15 • CITRUS UNSHNU PEEL EXTRACT • THYMUS SERPYLLUM EXTRACT • FAGUS SYLVATICA BUD EXTRACT • HYDROLYZED CONCHILIN PROTEIN • DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER • DIMETHICONE PEG-10/15 CROSSPOLYMER • BARIUM SULFATE • SYNTHETIC FLUORPHLOGHITE • ALUMINUM HYDROXIDE • POLYSILOXANE-2 • SODIUM EDTA • STEARIC ACID • CALCIUM ALUMINUM BORSILICATE • TOCOPHEROL • DIPHENYLENE GLYCOL • TRIETHOXYCAPRYLYLSILANE • TRIMETHOXYSYLETHYL POLYDIMETHYLSILOXYETHYL DIMETHICONE • BHT • ALLYLVA • POLYMETHYLSILOXOXANE • PHENOXYETHANOL • TITANIUM DIOXIDE • IRON OXIDES • MICA

Other Information

protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com




<http://s1872.com/16732AA>
See our website for more information.

縮小サイズ0.65倍 6mmノブ
11mm×26mm

7 30852 16738 4

12714970

12713286
AAA



210 / Birch

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 220 Linen

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

Drug Facts
(continued)

Inactive ingredients

WATER • DIMETHICONE •
COPHENYLSILOXY PHENYL TRIMETHICONE • GLYCERIN •
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE •
BUTYLENE GLYCOL • HYDROGENATED POLYDECENE •
PEG-10 DIMETHICONE •
BIS-BUTYLDIMETHICONE POLYBICYCLERYL-3 •
DISTEARDINOLAM HEPTOATE • ERYTHRITOL • XYLITOL •
SORBITAN SESQUISOOSTEATE • POLYQUATERNIUM-15 •
CITRUS UNSHNU PEEL EXTRACT •
THYMUS SERPYLLUM EXTRACT •
FAGUS SYLVATICA BUD EXTRACT •
HYDROLYZED CONCHILIN PROTEIN •
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER •
DIMETHICONE PEG-10/15 CROSSPOLYMER •
BARIUM SULFATE • SYNTHETIC FLUOROPHLOGHITE •
ALUMINUM HYDROXIDE • POLYSILOXANE-2 •
SODIUM EDTA • STEARIC ACID •
CALCIUM ALUMINUM BORSILICATE • TOCOPHEROL •
DIPROPYLENE GLYCOL • TRIETHOXYCAPRYLYLSILANE •
TRIMETHOXYSYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE •
BHT • ALUMINA • POLYMETHYLSILOXANOLANE •
PHENOXYETHANOL • TITANIUM DIOXIDE • IRON OXIDES •
MICA •

Other Information
protect this product in this container from excessive heat and direct sun.

Questions or Comments?
Call toll free 1-800-906-7503

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com



<http://s1872.com/16732AA>
See our website for more information.

30mL NET WT. 1.2 OZ.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16739 1
12714972



220 / Linen

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 230 Alder

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0% ...
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

Drug Facts
(continued)

Inactive ingredients

WATER - DIMETHICONE -
COPHENYLSILOXY PHENYL TRIMETHICONE - GLYCERIN -
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE -
BUTYLENE GLYCOL - HYDROGENATED POLYDECENE -
PEG-10 DIMETHICONE -
BIS-BUTYLDIMETHICONE POLYBICYCERYL-3 -
DISTEAROINDIUM HECTORITE - ERYTHRITOL - XYLITOL -
SORBITAN SESQUISOOSTEARATE - POLYQUATERNIUM-15 -
CITRUS UNSHNU PEEL EXTRACT -
THYMUS SERPYLLUM EXTRACT -
FAGUS SYLVATICA BUD EXTRACT -
HYDROLYZED CONCHILIN PROTEIN -
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER -
DIMETHICONE PEG-10/15 CROSSPOLYMER -
BARIUM SULFATE - SYNTHETIC FLUOROPHLOGHITE -
ALUMINUM HYDROXIDE - POLYSILOXANE-2 -
SODIUM EDTA - STEARIC ACID -
CALCIUM ALUMINUM BODORSULFATE - TOCOPHEROL -
DIPROPYLENE GLYCOL - TRIETHOXYCAPRYLYLSILANE -
TRIMETHOXYSYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE -
BHT - ALUMINA - POLYMETHYLSILOXOXANE -
PHENOXYETHANOL - TITANIUM DIOXIDE - IRON OXIDES -
MICA

Other Information

protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com




<http://s1872.com/16732AA>
See our website for more information.

30mL NET WT. 1.2 OZ.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16740 7
12714974



230 / Alder

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 240 Quartz

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients

Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

Drug Facts
(continued)

Inactive ingredients

WATER • DIMETHICONE •
COPHENYLSILOXY PHENYL TRIMETHICONE • GLYCERIN •
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE •
BUTYLENE GLYCOL • HYDROGENATED POLYDECENE •
PEG-10 DIMETHICONE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
DISTEARDINOLAM HEPTOATE • ERYTHRITOL • XYLITOL •
SORBITAN SESQUISOOSTEATE • POLYQUATERNIUM-15 •
CITRUS UNSHNU PEEL EXTRACT •
THYMUS SERPYLLUM EXTRACT •
FAGUS SYLVATICA BUD EXTRACT •
HYDROLYZED CONCHILIN PROTEIN •
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER •
DIMETHICONE PEG-10/15 CROSSPOLYMER •
BARIUM SULFATE • SYNTHETIC FLUOROPHLOGHITE •
ALUMINUM HYDROXIDE • POLYSILOXANE-2 •
SODIUM EDTA • STEARIC ACID •
CALCIUM ALUMINUM BODOSULFATE • TOCOPHEROL •
DIPROPYLENE GLYCOL • TRIETHOXYCAPRYLYLSILANE •
TRIMETHOXYSYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE •
BHT • ALUMINA • POLYMETHYLSILOXOXANE •
PHENOXYETHANOL • TITANIUM DIOXIDE • IRON OXIDES •
MICA •

Other Information

protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com




<http://s1872.com/16732AA>
See our website for more information.

30mL NET WT. 1.2 OZ.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16741 4
12714976



240 / Quartz

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 250 Sand

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- **Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

Drug Facts
(continued)

Inactive ingredients

WATER • DIMETHICONE •
COPHENYLSILOXY PHENYL TRIMETHICONE • GLYCERIN •
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSEBACATE •
BUTYLENE GLYCOL • HYDROGENATED POLYDECENE •
PEG-10 DIMETHICONE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
DISTEARDINOLAM HEPTOATE • ERYTHRITOL • XYLITOL •
SORBITAN SESQUISOOSTEATE • POLYQUATERNIUM-15 •
CITRUS UNSHNU PEEL EXTRACT •
THYMUS SERPYLLUM EXTRACT •
FAGUS SYLVATICA BUD EXTRACT •
HYDROLYZED CONCHILIN PROTEIN •
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER •
DIMETHICONE PEG-10/15 CROSSPOLYMER •
BARIUM SULFATE • SYNTHETIC FLUOROPHLOGHITE •
ALUMINUM HYDROXIDE • POLYSILOXANE-2 •
SODIUM EDTA • STEARIC ACID •
CALCIUM ALUMINUM BORSILICATE • TOCOPHEROL •
DIPROPYLENE GLYCOL • TRIETHOXYCAPRYLYLSILANE •
TRIMETHOXYSYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE •
BHT • ALUMINA • POLYMETHYLSILOXANOLANE •
PHENOXYETHANOL • TITANIUM DIOXIDE • IRON OXIDES •
MICA •

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com




<http://s1872.com/16732AA>
See our website for more information.

30mL NET WT. 1.2 OZ.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16742 1
12714978



250 / Sand

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 260 Cashmere

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

Drug Facts
(continued)

Inactive ingredients

WATER • DIMETHICONE •
C12-PHENYLSILOXY PHENYL TRIMETHICONE • GLYCERIN •
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE •
BUTYLENE GLYCOL • HYDROGENATED POLYDECENE •
PEG-10 DIMETHICONE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
DISTEAROINDIUM HECTORITE • ERYTHRITOL • XYLITOL •
SORBITAN SESQUISOOSTEARATE • POLYQUATERNIUM-15 •
CITRUS UNSHNU PEEL EXTRACT •
THYMUS SERPYLLUM EXTRACT •
FAGUS SYLVATICA BUD EXTRACT •
HYDROLYZED CONCHILIN PROTEIN •
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER •
DIMETHICONE PEG-10/15 CROSSPOLYMER •
BARIUM SULFATE • SYNTHETIC FLUOROPHLOGHITE •
ALUMINUM HYDROXIDE • POLYSILOXANE-2 •
SODIUM EDTA • STEARIC ACID •
CALCIUM ALUMINUM BORSILICATE • TOCOPHEROL •
DIPROPYLENE GLYCOL • TRIETHOXYCAPRYLYLSILANE •
TRIMETHOXYSILYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE •
BHT • ALUMINA • POLYMETHYLSILOXOXANE •
PHENOXYETHANOL • TITANIUM DIOXIDE • IRON OXIDES •
MICA •

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com





<http://s1872.com/16732AA>
See our website for more information.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16743 8
12714980



260 / Cashmere

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 310 Silk

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

Drug Facts
(continued)

Inactive ingredients

WATER - DIMETHICONE -
COPHENYLSILOXY PHENYL TRIMETHICONE - GLYCERIN -
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE -
BUTYLENE GLYCOL - HYDROGENATED POLYDECENE -
PEG-10 DIMETHICONE -
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 -
DISTEARDINOLAM HEPTOATE - ERYTHRITOL - XYLITOL -
SORBITAN SESQUISOOSTEATE - POLYQUATERMUM-81 -
CITRUS UNSHNU PEEL EXTRACT -
THYMUS SERPYLLUM EXTRACT -
FAGUS SYLVATICA BUD EXTRACT -
HYDROLYZED CONCHILIN PROTEIN -
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER -
DIMETHICONE PEG-10/15 CROSSPOLYMER -
BARIUM SULFATE - SYNTHETIC FLUORPHLOGHITE -
ALUMINUM HYDROXIDE - POLYSILOXANE-2 -
SODIUM EDTA - STEARIC ACID -
CALCIUM ALUMINUM BODOSULFATE - TOCOPHEROL -
DIPROPYLENE GLYCOL - TRIETHOXYCAPRYLYLSILANE -
TRIMETHOXYSYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE -
BHT - ALUMINA - POLYMETHYLSILOXOXANE -
PHENOXYETHANOL - TITANIUM DIOXIDE - IRON OXIDES -
MICA -

Other Information

protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com





<http://s1872.com/16732AA>
See our website for more information.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16744 5
12714982



310 / Silk

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 320 Pine

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

Drug Facts
(continued)

Inactive ingredients

WATER - DIMETHICONE -
DIPHENYLSILOXY PHENYL TRIMETHICONE - GLYCERIN -
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE -
BUTYLENE GLYCOL - HYDROGENATED POLYOLEFENE -
PEG-10 DIMETHICONE -
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 -
DISTEARDINOLIUM HECTORITE - ERYTHRITOL - XYLITOL -
SORBITAN SESQUISOOSTEATE - POLYQUATERNIUM-15 -
CITRUS UNSHNU PEEL EXTRACT -
THYMUS SERPYLLUM EXTRACT -
FAGUS SYLVATICA BUD EXTRACT -
HYDROLYZED CONCHILIN PROTEIN -
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER -
DIMETHICONE PEG-10/15 CROSSPOLYMER -
BARIUM SULFATE - SYNTHETIC FLUOROPHLOGHITE -
ALUMINUM HYDROXIDE - POLYSILOXANE-2 -
SODIUM EDTA - STEARIC ACID -
CALCIUM ALUMINUM BORSILICATE - TOCOPHEROL -
PROPYLENE GLYCOL - TRIETHOXYCAPRYLYLSILANE -
TRIMETHOXYSYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE -
BHT - ALUMINA - POLYMETHYLSILOXOXANE -
PHENOXYETHANOL - TITANIUM DIOXIDE - IRON OXIDES -
MICA -

Other Information

protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com




<http://s1872.com/16732AA>
See our website for more information.

縮小サイズ0.65倍 6mmパー
11mm×26mm

7 30852 16745 2

12714984

12713286
AAA



320 / Pine

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 330 Bamboo

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

Drug Facts
(continued)

Inactive ingredients

WATER • DIMETHICONE •
COPHENYLSILOXY PHENYL TRIMETHICONE • GLYCERIN •
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE •
BUTYLENE GLYCOL • HYDROGENATED POLYDECENE •
PEG-10 DIMETHICONE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
DISTEAROINDIUM HECTORITE • ERYTHRITOL • XYLITOL •
SORBITAN SESQUISOOSTEARATE • POLYQUATERNIUM-15 •
CITRUS UNSHNU PEEL EXTRACT •
THYMUS SERPYLLUM EXTRACT •
FAGUS SYLVATICA BUD EXTRACT •
HYDROLYZED CONCHILIN PROTEIN •
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER •
DIMETHICONE/PEG-10/15 CROSSPOLYMER •
BARIUM SULFATE • SYNTHETIC FLUOROPHLOGHITE •
ALUMINUM HYDROXIDE • POLYSILOXANE-2 •
SODIUM EDTA • STEARIC ACID •
CALCIUM ALUMINUM BORSILICATE • TOCOPHEROL •
DIPROPYLENE GLYCOL • TRIETHOXYCAPRYLYLSILANE •
TRIMETHOXYSYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE •
BHT • ALUMINA • POLYMETHYLSILOXOXANE •
PHENOXYETHANOL • TITANIUM DIOXIDE • IRON OXIDES •
MICA •

Other Information

protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com




<http://s1872.com/16732AA>
See our website for more information.

30mL NET WT. 1.2 OZ.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16746 9
12714986



330 / Bamboo

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 340 Oak

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients

Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only
Do not use on damaged or broken skin
When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:
 ■ apply liberally 15 minutes before sun exposure
 ■ use a water resistant sunscreen if swimming or sweating
 ■ reapply at least every 2 hours
 ■ children under 6 months: Ask a doctor
Sun Protection Measures.
 Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 • limit time in the sun, especially from 10 a.m. - 2 p.m.
 • wear long-sleeve shirts, pants, hats, and sunglasses

Drug Facts
(continued)

Inactive ingredients

WATER • DIMETHICONE • C12-PHENYLSILOXY PHENYL TRIMETHICONE • GLYCERIN • TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE • BUTYLENE GLYCOL • HYDROGENATED POLYDECENE • PEG-10 DIMETHICONE • BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 • DISTEARDINOLAM HEPTOATE • ERYTHRITOL • XYLITOL • SORBITAN SESQUISOOSTEATE • POLYOLATERMUM-81 • CITRUS UNSHNU PEEL EXTRACT • THYMUS SERPYLLUM EXTRACT • FAGUS SYLVATICA BUD EXTRACT • HYDROLYZED CONCHILIN PROTEIN • DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER • DIMETHICONE PEG-10/15 CROSSPOLYMER • BARIUM SULFATE • SYNTHETIC FLUOROPHOSPHITE • ALUMINUM HYDROXIDE • POLYSILOXANE-2 • SODIUM EDTA • STEARIC ACID • CALCIUM ALUMINUM BORSILICATE • TOCOPHEROL • DIPROPYLENE GLYCOL • TRIETHOXYCAPRYLYLSILANE • TRIMETHOXYSILYLETHYL POLYDIMETHYLSILOXYETHYL DIMETHICONE • BHT • ALLYL DIMETHYLSESQUOXANE • PHEVOXYETHANOL • TITANIUM DIOXIDE • IRON OXIDES • MICA

Other Information

protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com




<http://s1872.com/16732AA>
See our website for more information.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16747 6
12714988



340 / Oak

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 350 Maple

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients

Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only
Do not use on damaged or broken skin
When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:
 ■ apply liberally 15 minutes before sun exposure
 ■ use a water resistant sunscreen if swimming or sweating
 ■ reapply at least every 2 hours
 ■ children under 6 months: Ask a doctor
 ■ **Sun Protection Measures.**
 Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 • limit time in the sun, especially from 10 a.m. - 2 p.m.
 • wear long-sleeve shirts, pants, hats, and sunglasses

Drug Facts
(continued)

Inactive ingredients

WATER • DIMETHICONE • C12-PHENYLSILOXY PHENYL TRIMETHICONE • GLYCERIN • TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE • BUTYLENE GLYCOL • HYDROGENATED POLYDECENE • PEG-10 DIMETHICONE • BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 • DISTEARDINOLAM HEPTOATE • ERYTHRITOL • XYLITOL • SORBITAN SESQUISOOSTEATE • POLYQUATERNIUM-15 • CITRUS UNSHNU PEEL EXTRACT • THYMUS SERPYLLUM EXTRACT • FAGUS SYLVATICA BUD EXTRACT • HYDROLYZED CONCHILIN PROTEIN • DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER • DIMETHICONE PEG-10/15 CROSSPOLYMER • BARIUM SULFATE • SYNTHETIC FLUOROPHLOGHITE • ALUMINUM HYDROXIDE • POLYSILOXANE-2 • SODIUM EDTA • STEARIC ACID • CALCIUM ALUMINUM BORSILICATE • TOCOPHEROL • PROPYLENE GLYCOL • TRIETHOXYCAPRYLYLSILANE • TRIMETHOXYETHYL POLYDIMETHYLSILOXYETHYL DIMETHICONE • BHT • ALUMINA • POLYMETHYLSILOXOXANE • PHENOXYETHANOL • TITANIUM DIOXIDE • IRON OXIDES • MICA

Other Information

protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com



<http://s1872.com/16732AA>
See our website for more information.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm
7 30852 16748 3
12714990



350 / Maple

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 360 Citrine

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

■ helps prevent sunburn
■ if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- **Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

Drug Facts
(continued)

Inactive ingredients

WATER • DIMETHICONE •
C12-PHENYLSILOXY PHENYL TRIMETHICONE • GLYCERIN •
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE •
BUTYLENE GLYCOL • HYDROGENATED POLYDECENE •
PEG-10 DIMETHICONE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
DISTEARDIMETHICONE HECTORITE • ERYTHRITOL • XYLITOL •
SORBITAN SESQUISOOSTEARATE • POLYQUATERNIUM-15 •
CITRUS UNSHNU PEEL EXTRACT •
THYMUS SERPYLLUM EXTRACT •
FAGUS SYLVATICA BUD EXTRACT •
HYDROLYZED CONCHILIN PROTEIN •
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER •
DIMETHICONE PEG-10/15 CROSSPOLYMER •
BARIUM SULFATE • SYNTHETIC FLUOROPHLOGHITE •
ALUMINUM HYDROXIDE • POLYSILOXANE-2 •
SODIUM EDTA • STEARIC ACID •
CALCIUM ALUMINUM BORSILICATE • TOCOPHEROL •
DIPROPYLENE GLYCOL • TRIETHOXYCAPRYLYLSILANE •
TRIMETHOXYSYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE •
BHT • ALLYLVA • POLYMETHYLSILOXOXANE •
PHENOXYETHANOL • TITANIUM DIOXIDE • IRON OXIDES •
MICA •

Other Information

■ protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com




<http://s1872.com/16732AA>
See our website for more information.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16749 0

12714002



12114992

360 / Citrine

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 410 Sunstone

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

Drug Facts
(continued)

Inactive ingredients

WATER • DIMETHICONE •
COPHENYLSILOXY PHENYL TRIMETHICONE • GLYCERIN •
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE •
BUTYLENE GLYCOL • HYDROGENATED POLYDECENE •
PEG-10 DIMETHICONE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
DISTEARDIMONIUM HECTORITE • ERYTHRITOL • XYLITOL •
SORBITAN SESQUISOOSTEARATE • POLYQUATERNIUM-15 •
CITRUS UNSHNU PEEL EXTRACT •
THYMUS SERPYLLUM EXTRACT •
FAGUS SYLVATICA BUD EXTRACT •
HYDROLYZED CONCHILIN PROTEIN •
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER •
DIMETHICONE/PEG-10/15 CROSSPOLYMER •
BARIUM SULFATE • SYNTHETIC FLUOROPHLOGHITE •
ALUMINUM HYDROXIDE • POLYSILOXANE-2 •
SODIUM EDTA • STEARIC ACID •
CALCIUM ALUMINUM BORSILICATE • TOCOPHEROL •
DIPROPYLENE GLYCOL • TRIETHOXYCAPRYLYLSILANE •
TRIMETHOXYETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE •
BHT • ALUMINA • POLYMETHYLSILOXOXANE •
PHENOXYETHANOL • TITANIUM DIOXIDE • IRON OXIDES •
MICA •

Other Information

protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com



<http://s1872.com/16732AA>
See our website for more information.



30mL NET WT. 1.2 OZ.

12713286
AAA

縮小サイズ0.65倍 6mmパー
11mm×26mm

7 30852 16750 6

12714994



410 / Sunstone

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 420 Bronze

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- **Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

Drug Facts
(continued)

Inactive ingredients

WATER - DIMETHICONE -
C12-PHENYLSILOXY PHENYL TRIMETHICONE - GLYCERIN -
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE -
BUTYLENE GLYCOL - HYDROGENATED POLYDECENE -
PEG-10 DIMETHICONE -
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 -
DISTEARDINOLAM HEPTOATE - ERYTHRITOL - XYLITOL -
SORBITAN SESQUISOOSTEATE - POLYQUATERNIUM-15 -
CITRUS UNSHNU PEEL EXTRACT -
THYMUS SERPYLLUM EXTRACT -
FAGUS SYLVATICA BUD EXTRACT -
HYDROLYZED CONCHILIN PROTEIN -
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER -
DIMETHICONE PEG-10/15 CROSSPOLYMER -
BARIUM SULFATE - SYNTHETIC FLUOROPHOSPHITE -
ALUMINUM HYDROXIDE - POLYSILOXANE-2 -
SODIUM EDTA - STEARIC ACID -
CALCIUM ALUMINUM BORSILICATE - TOCOPHEROL -
DIPROPYLENE GLYCOL - TRIETHOXYCAPRYLYLSILANE -
TRIMETHOXYSILYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE -
BHT - ALUMINA - POLYMETHYLSILOXOXANE -
PHENOXYETHANOL - TITANIUM DIOXIDE - IRON OXIDES -
MICA -

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com




<http://s1872.com/16732AA>
See our website for more information.

30mL NET WT. 1.2 OZ.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16751 3
12714996



420 / Bronze

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 430 Cedar

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- **Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

Drug Facts
(continued)

Inactive ingredients

WATER • DIMETHICONE •
COPHENYLSILOXY PHENYL TRIMETHICONE • GLYCERIN •
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSEBACATE •
BUTYLENE GLYCOL • HYDROGENATED POLYOLEFENE •
PEG-10 DIMETHICONE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
DISTEAROINDIUM HECTORITE • ERYTHRITOL • XYLITOL •
SORBITAN SESQUISOSEBATE • POLYQUATERNIUM-15 •
CITRUS UNSHNU PEEL EXTRACT •
THYMUS SERPYLLUM EXTRACT •
FAGUS SYLVATICA BUD EXTRACT •
HYDROLYZED CONCHILIN PROTEIN •
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER •
DIMETHICONE PEG-10/15 CROSSPOLYMER •
BARIUM SULFATE • SYNTHETIC FLUOROPHLOGHITE •
ALUMINUM HYDROXIDE • POLYSILOXANE-2 •
SODIUM EDTA • STEARIC ACID •
CALCIUM ALUMINUM BORSILICATE • TOCOPHEROL •
DIPROPYLENE GLYCOL • TRIETHOXYCAPRYLYLSILANE •
TRIMETHOXYETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE •
BHT • ALLYLVA • POLYMETHYLSILOXOXANE •
PHENOXYETHANOL • TITANIUM DIOXIDE • IRON OXIDES •
MICA •

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com




<http://s1872.com/16732AA>
See our website for more information.

30mL NET WT. 1.2 OZ.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16752 0
12715101



430 / Cedar

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 440 Amber

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients

Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only
Do not use on damaged or broken skin
When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:
 ■ apply liberally 15 minutes before sun exposure
 ■ use a water resistant sunscreen if swimming or sweating
 ■ reapply at least every 2 hours
 ■ children under 6 months: Ask a doctor
 ■ **Sun Protection Measures.**
 Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 • limit time in the sun, especially from 10 a.m. - 2 p.m.
 • wear long-sleeve shirts, pants, hats, and sunglasses

Drug Facts
(continued)

Inactive ingredients

WATER • DIMETHICONE • CYPHENYLSILOXY PHENYL TRIMETHICONE • GLYCERIN • TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSEBACATE • BUTYLENE GLYCOL • HYDROGENATED POLYDECENE • PEG-10 DIMETHICONE • BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 • DISTEARDINOLAM HEPTOATE • ERYTHRITOL • XYLITOL • SORBITAN SESQUISOSEBATE • POLYQUATERNIUM-15 • CITRUS UNSHNU PEEL EXTRACT • THYMUS SERPYLLUM EXTRACT • FAGUS SYLVATICA BUD EXTRACT • HYDROLYZED CONCHILIN PROTEIN • DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER • DIMETHICONE PEG-10/15 CROSSPOLYMER • BARIUM SULFATE • SYNTHETIC FLUOROPHLOGHITE • ALUMINUM HYDROXIDE • POLYSILOXANE-2 • SODIUM EDTA • STEARIC ACID • CALCIUM ALUMINUM BODOSILICATE • TOCOPHEROL • DIPROPYLENE GLYCOL • TRIETHOXYCAPRYLYLSILANE • TRIMETHOXYSYLETHYL POLYDIMETHYLSILOXYETHYL DIMETHICONE • BHT • ALLYLVA • POLYMETHYLSILOXANOLANE • PHEVOXYETHANOL • TITANIUM DIOXIDE • IRON OXIDES • MICA

Other Information

protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com




<http://s1872.com/16732AA>
See our website for more information.

縮小サイズ0.65倍 6mmノバー
11mm×26mm

7 30852 16753 7

12715103

12713286
AAA



440 / Amber

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 450 Copper

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

Drug Facts
(continued)

Inactive ingredients

WATER • DIMETHICONE •
CIPHENYLSILOXY PHENYL TRIMETHICONE • GLYCERIN •
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE •
BUTYLENE GLYCOL • HYDROGENATED POLYOLEFENE •
PEG-10 DIMETHICONE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
DISTEAROINDIUM HECTORITE • ERYTHRITOL • XYLITOL •
SORBITAN SESQUISOOSTEARATE • POLYQUATERNIUM-15 •
CITRUS UNSHNU PEEL EXTRACT •
THYMUS SERPYLLUM EXTRACT •
FAGUS SYLVATICA BUD EXTRACT •
HYDROLYZED CONCHILIN PROTEIN •
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER •
DIMETHICONE PEG-10/15 CROSSPOLYMER •
BARIUM SULFATE • SYNTHETIC FLUOROPHLOGHITE •
ALUMINUM HYDROXIDE • POLYSILOXANE-2 •
SODIUM EDTA • STEARIC ACID •
CALCIUM ALUMINUM BORSILICATE • TOCOPHEROL •
DIPROPYLENE GLYCOL • TRIETHOXYCAPRYLYLSILANE •
TRIMETHOXYSYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE •
BHT • ALLYLVA • POLYMETHYLSILOXOXANE •
PHENOXYETHANOL • TITANIUM DIOXIDE • IRON OXIDES •
MICA •

Other Information

protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com




<http://s1872.com/16732AA>
See our website for more information.

30mL NET WT. 1.2 OZ.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16754 4
12715105



450 / Copper

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 460 Topaz

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

Drug Facts
(continued)

Inactive ingredients

WATER - DIMETHICONE -
COPHENYLSILOXY PHENYL TRIMETHICONE - GLYCERIN -
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE -
BUTYLENE GLYCOL - HYDROGENATED POLYDECENE -
PEG-10 DIMETHICONE -
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 -
DISTEARDINOLAM HEPTOATE - ERYTHRITOL - XYLITOL -
SORBITAN SESQUISOESTEATE - POLYQUATERNIUM-15 -
CITRUS UNSHNU PEEL EXTRACT -
THYMUS SERPYLLUM EXTRACT -
FAGUS SYLVATICA BUD EXTRACT -
HYDROLYZED CONCHILIN PROTEIN -
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER -
DIMETHICONE PEG-10/15 CROSSPOLYMER -
BARIUM SULFATE - SYNTHETIC FLUOROPHOSPHITE -
ALUMINUM HYDROXIDE - POLYSILOXANE-2 -
SODIUM EDTA - STEARIC ACID -
CALCIUM ALUMINUM BORSILICATE - TOCOPHEROL -
PROPYLENE GLYCOL - TRIETHOXYCAPRYLYLSILANE -
TRIMETHOXYSYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE -
BHT - ALUMINA - POLYMETHYLSILOXOXANE -
PHENOXYETHANOL - TITANIUM DIOXIDE - IRON OXIDES -
MICA -

Other Information

protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com




<http://s1872.com/16732AA>
See our website for more information.

30mL NET WT. 1.2 OZ.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16755 1
12715107



460 / Topaz

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 510 Suede

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

Drug Facts
(continued)

Inactive ingredients

WATER • DIMETHICONE •
COPHENYLSILOXY PHENYL TRIMETHICONE • GLYCERIN •
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE •
BUTYLENE GLYCOL • HYDROGENATED POLYDECENE •
PEG-10 DIMETHICONE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
DISTEAROINDONIUM HECTORITE • ERYTHRITOL • XYLITOL •
SORBITAN SESQUISOOSTEARATE • POLYQUATERNIUM-15 •
CITRUS UNSHNU PEEL EXTRACT •
THYMUS SERPYLLUM EXTRACT •
FAGUS SYLVATICA BUD EXTRACT •
HYDROLYZED CONCHILIN PROTEIN •
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER •
DIMETHICONE PEG-10/15 CROSSPOLYMER •
BARIUM SULFATE • SYNTHETIC FLUOROPHLOGHITE •
ALUMINUM HYDROXIDE • POLYSILOXANE-2 •
SODIUM EDTA • STEARIC ACID •
CALCIUM ALUMINUM BORSILICATE • TOCOPHEROL •
DIPROPYLENE GLYCOL • TRIETHOXYCAPRYLYLSILANE •
TRIMETHOXYSYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE •
BHT • ALUMINA • POLYMETHYLSILOXOXANE •
PHENOXYETHANOL • TITANIUM DIOXIDE • IRON OXIDES •
MICA •

Other Information

protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com




<http://s1872.com/16732AA>
See our website for more information.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16756 8
12715109



510 / Suede

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 520 Rosewood

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

Drug Facts
(continued)

Inactive ingredients

WATER - DIMETHICONE -
C12-PHENYLSILOXY PHENYL TRIMETHICONE - GLYCERIN -
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE -
BUTYLENE GLYCOL - HYDROGENATED POLYDECENE -
PEG-10 DIMETHICONE -
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 -
DISTEAROINDIUM HECTORITE - ERYTHRITOL - XYLITOL -
SORBITAN SESQUISOOSTEARATE - POLYQUATERNIUM-15 -
CITRUS UNSHNU PEEL EXTRACT -
THYMUS SERPYLLUM EXTRACT -
FAGUS SYLVATICA BUD EXTRACT -
HYDROLYZED CONCHILIN PROTEIN -
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER -
DIMETHICONE PEG-10/15 CROSSPOLYMER -
BARIUM SULFATE - SYNTHETIC FLUOROPHLOGHITE -
ALUMINUM HYDROXIDE - POLYSILOXANE-2 -
SODIUM EDTA - STEARIC ACID -
CALCIUM ALUMINUM BORSILICATE - TOCOPHEROL -
DIPROPYLENE GLYCOL - TRIETHOXYCAPRYLYLSILANE -
TRIMETHOXYSYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE -
BHT - ALLYLVA - POLYMETHYLSILOXOXANE -
PHENOXYETHANOL - TITANIUM DIOXIDE - IRON OXIDES -
MICA -

Other Information

protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com




<http://s1872.com/16732AA>
See our website for more information.

30mL NET WT. 1.2 OZ.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16757 5
12715111



520 / Rosewood

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 530 Henna

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

Drug Facts
(continued)

Inactive ingredients

WATER - DIMETHICONE -
COPHENYLSILOXY PHENYL TRIMETHICONE - GLYCERIN -
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE -
BUTYLENE GLYCOL - HYDROGENATED POLYDECENE -
PEG-10 DIMETHICONE -
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 -
DISTEARDINOLAM HEPTOATE - ERYTHRITOL - XYLITOL -
SORBITAN SESQUISOOSTEATE - POLYOLATERMUM-851 -
CITRUS UNSHNU PEEL EXTRACT -
THYMUS SERPYLLUM EXTRACT -
FAGUS SYLVATICA BUD EXTRACT -
HYDROLYZED CONCHILIN PROTEIN -
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER -
DIMETHICONE PEG-10/15 CROSSPOLYMER -
BARIUM SULFATE - SYNTHETIC FLUOROPHLOGHITE -
ALUMINUM HYDROXIDE - POLYSILOXANE-2 -
SODIUM EDTA - STEARIC ACID -
CALCIUM ALUMINUM BORSILICATE - TOCOPHEROL -
DIPROPYLENE GLYCOL - TRIETHOXYCAPRYLYLSILANE -
TRIMETHOXYSYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE -
BHT - ALLYLVA - POLYMETHYLSILOXOXANE -
PHENOXYETHANOL - TITANIUM DIOXIDE - IRON OXIDES -
MICA -

Other Information

protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com



<http://s1872.com/16732AA>
See our website for more information.


30mL NET WT. 1.2 OZ.

12713286
AAA

縮小サイズ0.65倍 6mmノバー
11mm×26mm

7 30852 16758 2

12715113



530 / Henna

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 540 Mahogany

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

Drug Facts
(continued)

Inactive ingredients

WATER • DIMETHICONE •
C12-PHENYLSILOXY PHENYL TRIMETHICONE • GLYCERIN •
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE •
BUTYLENE GLYCOL • HYDROGENATED POLYDECENE •
PEG-10 DIMETHICONE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
DISTEARDINOLAM HEPTOATE • ERYTHRITOL • XYLITOL •
SORBITAN SESQUISOOSTEATE • POLYQUATERNIUM-15 •
CITRUS UNSHNU PEEL EXTRACT •
THYMUS SERPYLLUM EXTRACT •
FAGUS SYLVATICA BUD EXTRACT •
HYDROLYZED CONCHILIN PROTEIN •
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER •
DIMETHICONE PEG-10/15 CROSSPOLYMER •
BARIUM SULFATE • SYNTHETIC FLUOROPHOSPHITE •
ALUMINUM HYDROXIDE • POLYSILOXANE-2 •
SODIUM EDTA • STEARIC ACID •
CALCIUM ALUMINUM BORSILICATE • TOCOPHEROL •
DIPROPYLENE GLYCOL • TRIETHOXYCAPRYLYLSILANE •
TRIMETHOXYSYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE •
BHT • ALLYL DIMETHYLSESQUOXANE •
PHENOXYETHANOL • TITANIUM DIOXIDE • IRON OXIDES •
MICA •

Other Information

protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com



<http://s1872.com/16732AA>
See our website for more information.


30mL NET WT. 1.2 OZ.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16759 9

12715115



12719119

540 / Mahogany

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 550 Jasper

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

Drug Facts
(continued)

Inactive ingredients

WATER - DIMETHICONE -
COPHENYLSILOXY PHENYL TRIMETHICONE - GLYCERIN -
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE -
BUTYLENE GLYCOL - HYDROGENATED POLYDECENE -
PEG-10 DIMETHICONE -
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 -
DISTEARDIMONIUM HECTORITE - ERYTHRITOL - XYLITOL -
SORBITAN SESQUISULFATE - POLYQUATERNIUM-15 -
CITRUS UNSHNU PEEL EXTRACT -
THYMUS SERPYLLUM EXTRACT -
FAGUS SYLVATICA BUD EXTRACT -
HYDROLYZED CONCHILIN PROTEIN -
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER -
DIMETHICONE PEG-10/15 CROSSPOLYMER -
BARIUM SULFATE - SYNTHETIC FLUOROPHOSPHITE -
ALUMINUM HYDROXIDE - POLYSILOXANE-2 -
SODIUM EDTA - STEARIC ACID -
CALCIUM ALUMINUM BORSILICATE - TOCOPHEROL -
PROPYLENE GLYCOL - TRIETHOXYCAPRYLYLSILANE -
TRIMETHOXYSYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE -
BHT - ALLYLVA - POLYMETHYLSILOXOXANE -
PHENOXYETHANOL - TITANIUM DIOXIDE - IRON OXIDES -
MICA -

Other Information

protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com



<http://s1872.com/16732AA>
See our website for more information.


30mL NET WT. 1.2 OZ.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16760 5

12715117



550 / Jasper

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton - 560 Obsidian

SHISEIDO
GINZA TOKYO

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

30mL NET WT. 1.2 OZ.

RADIANT LIFTING
Foundation



SHISEIDO
GINZA TOKYO

東京銀座

Lift your skin with light-adjusting radiance. SHISEIDO Light-Adjusting technology instantly adapts and responds to every light source to smooth and diffuse fine lines and create a luminous, lifted effect on the skin. All day wear. Weightless comfort. Radiant finish. Medium to full, buildable coverage.

DERMATOLOGIST-TESTED.
NON-COMEDOGENIC.

Drug Facts

Active ingredients
Purpose
OCTINOXATE 6.9%
Sunscreen
TITANIUM DIOXIDE 1.0%
Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Drug Facts
(continued)

Warnings

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable: Do not use near heat, flame or while smoking.

Directions

For sunscreen use:

- apply liberally 15 minutes before sun exposure
- use a water resistant sunscreen if swimming or sweating
- reapply at least every 2 hours
- children under 6 months: Ask a doctor
- **Sun Protection Measures.**

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.
- wear long-sleeve shirts, pants, hats, and sunglasses

SYNCHRO SKIN
RADIANT LIFTING

Foundation
Oil-Free

BROAD SPECTRUM
SPF 30

Sunscreen

Drug Facts
(continued)

Inactive ingredients

WATER • DIMETHICONE •
C-PHENYLSILOXY PHENYL TRIMETHICONE • GLYCERIN •
TRIFLUOROPROPYLDIMETHYLTRIMETHYLSILOXYSULFATE •
BUTYLENE GLYCOL • HYDROGENATED POLYDECENE •
PEG-10 DIMETHICONE •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
DISTEARDIMONIUM HECTORITE • ERYTHRITOL • XYLITOL •
SORBITAN SESQUISOOSTEATE • POLYQUATERNIUM-15 •
CITRUS UNSHNU PEEL EXTRACT •
THYMUS SERPYLLUM EXTRACT •
FAGUS SYLVATICA BUD EXTRACT •
HYDROLYZED CONCHILIN PROTEIN •
DIMETHICONE VINYL DIMETHICONE CROSSPOLYMER •
DIMETHICONE PEG-10/15 CROSSPOLYMER •
BARIUM SULFATE • SYNTHETIC FLUORPHLOGHITE •
ALUMINUM HYDROXIDE • POLYSILOXANE-2 •
SODIUM EDTA • STEARIC ACID •
CALCIUM ALUMINUM BORSILICATE • TOCOPHEROL •
DIPROPYLENE GLYCOL • TRIETHOXYCAPRYLYLSILANE •
TRIMETHOXYSILYLETHYL POLYDIMETHYLSILOXYETHYL
DIMETHICONE •
BHT • ALUMINA • POLYMETHYLSILOXOXANE •
PHENOXYETHANOL • TITANIUM DIOXIDE • IRON OXIDES •
MICA •

Other Information

- protect this product in this container from excessive heat and direct sun.

Questions or Comments?

Call toll free 1-800-906-7503

SHISEIDO AMERICAS
CORPORATION DIST.
NEW YORK, NY 10017
MADE IN U.S.A.
www.shiseido.com



<http://s1872.com/16732AA>
See our website for more information.

30mL NET WT. 1.2 OZ.

12713286
AAA

縮小サイズ0.65倍 6mmバー
11mm×26mm

7 30852 16761 2

560 Obsidian

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 110

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-563
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3COOX)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
TANGERINE PEEL (UNII: JU3D414057)	
THYMUS SERPYLLUM WHOLE (UNII: 86H456K51N)	
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
BARIUM SULFATE (UNII: 25BB7EKE2E)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALUMINUM OXIDE (UNII: LMI26O6933)	

POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-563-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 120

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-564
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3C0OX)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL	

METHACRYLATE; 3:7 (UNII: EL9825H96J)
TANGERINE PEEL (UNII: JU3D414057)
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)
BARIUM SULFATE (UNII: 25BB7EKE2E)
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)
EDETATE DISODIUM (UNII: 7FLD91C86K)
STEARIC ACID (UNII: 4ELV7Z65AP)
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
DIPROPYLENE GLYCOL (UNII: E107L85C40)
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
ALUMINUM OXIDE (UNII: LM26O6933)
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-564-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 130

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-565
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3COOX)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
TANGERINE PEEL (UNII: JU3D414057)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
BARIUM SULFATE (UNII: 25BB7EKE2E)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-565-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 140

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-566
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3COOX)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
TANGERINE PEEL (UNII: JU3D414057)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
BARIUM SULFATE (UNII: 25BB7EKE2E)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALUMINUM OXIDE (UNII: LM26O6933)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-566-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 150

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-567
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3COOX)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
TANGERINE PEEL (UNII: JU3D414057)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
BARIUM SULFATE (UNII: 25BB7EKE2E)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)	

.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
DIPROPYLENE GLYCOL (UNII: E107L85C40)
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
ALUMINUM OXIDE (UNII: LMI2606933)
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-567-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 160

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-568
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3COOX)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	

DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)
ERYTHRITOL (UNII: RA96B954X6)
XYLITOL (UNII: VCQ006KQ1E)
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)
TANGERINE PEEL (UNII: JU3D414057)
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)
BARIUM SULFATE (UNII: 25BB7EKE2E)
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)
EDETATE DISODIUM (UNII: 7FLD91C86K)
STEARIC ACID (UNII: 4ELV7Z65AP)
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
DIPROPYLENE GLYCOL (UNII: E107L85C40)
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
ALUMINUM OXIDE (UNII: LMI26O6933)
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-568-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 210
 octinoxate and titanium dioxide emulsion

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-569
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3C00X)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
TANGERINE PEEL (UNII: JU3D414057)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
BARIUM SULFATE (UNII: 25BB7EKE2E)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-569-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

OTC MONOGRAPH DRUG	M020	09/20/2020	
-----------------------	------	------------	--

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 220

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-570
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3C0OX)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
TANGERINE PEEL (UNII: JU3D414057)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
BARIUM SULFATE (UNII: 25BB7EKE2E)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-570-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 230

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-571
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3C0OX)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
TANGERINE PEEL (UNII: JU3D414057)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
BARIUM SULFATE (UNII: 25BB7EKE2E)	

ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)
EDETATE DISODIUM (UNII: 7FLD91C86K)
STEARIC ACID (UNII: 4ELV7Z65AP)
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
DIPROPYLENE GLYCOL (UNII: E107L85C40)
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
ALUMINUM OXIDE (UNII: LMI26O6933)
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-571-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 240

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-572
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	

GLYCERIN (UNII: PDC6A3C0OX)
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)
ERYTHRITOL (UNII: RA96B954X6)
XYLITOL (UNII: VCQ006KQ1E)
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)
TANGERINE PEEL (UNII: JU3D414057)
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)
BARIUM SULFATE (UNII: 25BB7EKE2E)
ALUMINUM HYDROXIDE (UNII: 5QB0T2IU0)
EDETATE DISODIUM (UNII: 7FLD91C86K)
STEARIC ACID (UNII: 4ELV7Z65AP)
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
DIPROPYLENE GLYCOL (UNII: E107L85C40)
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
ALUMINUM OXIDE (UNII: LMI26O6933)
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-572-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 250

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-573
---------------------	----------------	---------------------------	---------------

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3C0OX)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
TANGERINE PEEL (UNII: JU3D414057)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4C00W6C5)	
BARIUM SULFATE (UNII: 25BB7EKE2E)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-573-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 260

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-574
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3C00X)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
TANGERINE PEEL (UNII: JU3D414057)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
BARIUM SULFATE (UNII: 25BB7EKE2E)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	

PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-574-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 310

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-575
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3C00X)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	

TANGERINE PEEL (UNII: JU3D414057)
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)
BARIUM SULFATE (UNII: 25BB7EKE2E)
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)
EDETATE DISODIUM (UNII: 7FLD91C86K)
STEARIC ACID (UNII: 4ELV7Z65AP)
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
DIPROPYLENE GLYCOL (UNII: E107L85C40)
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
ALUMINUM OXIDE (UNII: LMI26O6933)
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-575-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 320

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-576
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3C0OX)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
TANGERINE PEEL (UNII: JU3D414057)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
BARIUM SULFATE (UNII: 25BB7EKE2E)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-576-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-577
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3C0OX)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
TANGERINE PEEL (UNII: JU3D414057)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
BARIUM SULFATE (UNII: 25BB7EKE2E)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start	Marketing End
---	-----------	---------------------	-----------------	---------------

#	Item Code	Package Description	Date	Date
1	NDC:58411-577-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 340

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-578
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3C0OX)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
TANGERINE PEEL (UNII: JU3D414057)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
BARIUM SULFATE (UNII: 25BB7EKE2E)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	

DIPROPYLENE GLYCOL (UNII: E107L85C40)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-578-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 350

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-579
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3C0OX)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	

ERYTHRITOL (UNII: RA96B954X6)
XYLITOL (UNII: VCQ006KQ1E)
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)
TANGERINE PEEL (UNII: JU3D414057)
THYMUS SERPYLLUM WHOLE (UNII: 86H456K51N)
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)
BARIUM SULFATE (UNII: 25BB7EKE2E)
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)
EDETATE DISODIUM (UNII: 7FLD91C86K)
STEARIC ACID (UNII: 4ELV7Z65AP)
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
DIPROPYLENE GLYCOL (UNII: E107L85C40)
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
ALUMINUM OXIDE (UNII: LMI26O6933)
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSO FERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-579-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 360

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-580
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3C0OX)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
TANGERINE PEEL (UNII: JU3D414057)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
BARIUM SULFATE (UNII: 25BB7EKE2E)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-580-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH	M020	09/20/2020	

DRUG	MU20	09/20/2020	
------	------	------------	--

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 410

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-581
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y10)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3C00X)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
TANGERINE PEEL (UNII: JU3D414057)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
BARIUM SULFATE (UNII: 25BB7EKE2E)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-581-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 420

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-582
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3C0OX)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
TANGERINE PEEL (UNII: JU3D414057)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
BARIUM SULFATE (UNII: 25BB7EKE2E)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	

EDETATE DISODIUM (UNII: 7FLD91C86K)
STEARIC ACID (UNII: 4ELV7Z65AP)
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
DIPROPYLENE GLYCOL (UNII: E107L85C40)
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
ALUMINUM OXIDE (UNII: LMI26O6933)
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSO FERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-582-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 430

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-583
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3COOX)	

BUTYLENE GLYCOL (UNII: 3XUS85K0RA)
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)
ERYTHRITOL (UNII: RA96B954X6)
XYLITOL (UNII: VCQ006KQ1E)
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)
TANGERINE PEEL (UNII: JU3D414057)
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)
BARIUM SULFATE (UNII: 25BB7EKE2E)
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)
EDETATE DISODIUM (UNII: 7FLD91C86K)
STEARIC ACID (UNII: 4ELV7Z65AP)
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
DIPROPYLENE GLYCOL (UNII: E107L85C40)
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
ALUMINUM OXIDE (UNII: LMI26O6933)
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-583-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 440

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-584
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3C0OX)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
TANGERINE PEEL (UNII: JU3D414057)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
BARIUM SULFATE (UNII: 25BB7EKE2E)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-584-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 450

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-585
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3C0OX)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
TANGERINE PEEL (UNII: JU3D414057)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
BARIUM SULFATE (UNII: 25BB7EKE2E)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-585-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 460

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-586
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3C0OX)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
TANGERINE PEEL (UNII: JU3D414057)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	

FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)
BARIUM SULFATE (UNII: 25BB7EKE2E)
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)
EDETATE DISODIUM (UNII: 7FLD91C86K)
STEARIC ACID (UNII: 4ELV7Z65AP)
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
DIPROPYLENE GLYCOL (UNII: E107L85C40)
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
ALUMINUM OXIDE (UNII: LMI26O6933)
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-586-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 510

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-587
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

WATER (UNII: 059QF0KO0R)
DIMETHICONE (UNII: 92RU3N3Y1O)
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)
GLYCERIN (UNII: PDC6A3C0OX)
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)
ERYTHRITOL (UNII: RA96B954X6)
XYLITOL (UNII: VCQ006KQ1E)
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)
TANGERINE PEEL (UNII: JU3D414057)
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)
BARIUM SULFATE (UNII: 25BB7EKE2E)
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)
EDETATE DISODIUM (UNII: 7FLD91C86K)
STEARIC ACID (UNII: 4ELV7Z65AP)
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
DIPROPYLENE GLYCOL (UNII: E107L85C40)
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
ALUMINUM OXIDE (UNII: LMI26O6933)
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-587-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 520

octinoxate and titanium dioxide emulsion

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-588	
Route of Administration	TOPICAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)		OCTINOXATE	2.484 g in 30 mL	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)		TITANIUM DIOXIDE	0.36 g in 30 mL	
Inactive Ingredients				
Ingredient Name				Strength
WATER (UNII: 059QF0KO0R)				
DIMETHICONE (UNII: 92RU3N3Y1O)				
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)				
GLYCERIN (UNII: PDC6A3C0OX)				
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)				
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)				
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)				
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)				
ERYTHRITOL (UNII: RA96B954X6)				
XYLITOL (UNII: VCQ006KQ1E)				
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)				
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)				
TANGERINE PEEL (UNII: JU3D414057)				
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)				
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)				
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)				
BARIUM SULFATE (UNII: 25BB7EKE2E)				
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)				
EDETATE DISODIUM (UNII: 7FLD91C86K)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)				
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)				
DIPROPYLENE GLYCOL (UNII: E107L85C40)				
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)				
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)				
ALUMINUM OXIDE (UNII: LMI26O6933)				
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)				
PHENOXYETHANOL (UNII: HIE492ZZ3T)				
FERRIC OXIDE RED (UNII: 1K09F3G675)				
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)				
FERROSFERRIC OXIDE (UNII: XM0M87F357)				
MICA (UNII: V8A1AW0880)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-588-10	1 in 1 CARTON	09/20/2020	

1	30 mL in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 530

octinoxate and titanium dioxide emulsion

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-589
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL	

Inactive Ingredients		
Ingredient Name	Strength	
WATER (UNII: 059QF0K00R)		
DIMETHICONE (UNII: 92RU3N3Y1O)		
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)		
GLYCERIN (UNII: PDC6A3C0OX)		
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)		
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)		
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)		
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)		
ERYTHRITOL (UNII: RA96B954X6)		
XYLITOL (UNII: VCQ006KQ1E)		
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)		
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)		
TANGERINE PEEL (UNII: JU3D414057)		
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)		
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)		
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)		
BARIUM SULFATE (UNII: 25BB7EKE2E)		
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)		
EDETATE DISODIUM (UNII: 7FLD91C86K)		
STEARIC ACID (UNII: 4ELV7Z65AP)		
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)		
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)		
DIPROPYLENE GLYCOL (UNII: E107L85C40)		
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)		
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)		

ALUMINUM OXIDE (UNII: LMI26O6933)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-589-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 540

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-590
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3C0OX)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	

POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)
TANGERINE PEEL (UNII: JU3D414057)
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)
BARIUM SULFATE (UNII: 25BB7EKE2E)
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)
EDETATE DISODIUM (UNII: 7FLD91C86K)
STEARIC ACID (UNII: 4ELV7Z65AP)
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
DIPROPYLENE GLYCOL (UNII: E107L85C40)
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
ALUMINUM OXIDE (UNII: LMI26O6933)
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)
PHENOXYETHANOL (UNII: HIE492ZZ3T)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-590-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 550

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-591
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3C0OX)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
TANGERINE PEEL (UNII: JU3D414057)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
BARIUM SULFATE (UNII: 25BB7EKE2E)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-591-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

SHISEIDO SYNCHRO SKIN RADIANT LIFTING FOUNDATION 560

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-592
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	2.484 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	0.36 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPHENYLSILOXY PHENYL TRIMETHICONE (UNII: I445L28B12)	
GLYCERIN (UNII: PDC6A3C0OX)	
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	
HYDROGENATED POLYDECENE TYPE I (UNII: U333RI6EB7)	
PEG-10 DIMETHICONE (600 CST) (UNII: 8PR7V1SVM0)	
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)	
ERYTHRITOL (UNII: RA96B954X6)	
XYLITOL (UNII: VCQ006KQ1E)	
SORBITAN SESQUIISOSTEARATE (UNII: VU97D01BF9)	
POLYQUATERNIUM-51 (2-METHACRYLOYLOXYETHYL PHOSPHORYLCHOLINE/N-BUTYL METHACRYLATE; 3:7) (UNII: EL9825H96J)	
TANGERINE PEEL (UNII: JU3D414057)	
THYMUS SERPYLLUM WHOLE (UNII: 86H4S6K51N)	
FAGUS SYLVATICA FLOWER BUD (UNII: 6D5V13045W)	
DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE) (UNII: 9E4CO0W6C5)	
BARIUM SULFATE (UNII: 25BB7EKE2E)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CALCIUM ALUMINUM BOROSILICATE (UNII: 3JRB8A35M0)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MICA (UNII: V8A1AW0880)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-592-10	1 in 1 CARTON	09/20/2020	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH DRUG	M020	09/20/2020	

Labeler - SHISEIDO AMERICAS CORPORATION (193691821)

Establishment

Name	Address	ID/FEI	Business Operations
Shiseido America Inc.		782677132	ANALYSIS(58411-563, 58411-564, 58411-565, 58411-566, 58411-567, 58411-568, 58411-569, 58411-570, 58411-571, 58411-572, 58411-573, 58411-574, 58411-575, 58411-576, 58411-577, 58411-578, 58411-579, 58411-580, 58411-581, 58411-582, 58411-583, 58411-584, 58411-585, 58411-586, 58411-587, 58411-588, 58411-589, 58411-590, 58411-591, 58411-592) , MANUFACTURE(58411-563, 58411-564, 58411-565, 58411-566, 58411-567, 58411-568, 58411-569, 58411-570, 58411-571, 58411-572, 58411-573, 58411-574, 58411-575, 58411-576, 58411-577, 58411-578, 58411-579, 58411-580, 58411-581, 58411-582, 58411-583, 58411-584, 58411-585, 58411-586, 58411-587, 58411-588, 58411-589, 58411-590, 58411-591, 58411-592)

Revised: 4/2020

SHISEIDO AMERICAS CORPORATION