

**EQUATE PAIN AND FEVER INFANTS- acetaminophen suspension**  
**Wal-Mart Stores Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Wal-Mart Infants' Pain & Fever Drug Facts**

**Active ingredient (in each 5 mL)**

Acetaminophen 160 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

**Warnings**

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if your child has**

liver disease

**Ask a doctor or pharmacist before use if your child is**

taking the blood thinning drug warfarin

**When using this product**

do not exceed recommended dose (see overdose warning)

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present. These could be signs of a serious condition.

**Keep out of reach of children.**

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise, use age.
- push air out of syringe. Insert syringe tip into bottle opening.
- flip bottle upside down. Pull yellow part of syringe to the first dose line and then push product back into bottle.
- pull yellow part of syringe until it reaches and stays at the correct dose
- dispense liquid slowly into child's mouth, toward inner cheek
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- replace cap tightly to maintain child resistance

Dosing Chart		
Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL

\*or as directed by a doctor

Attention: use only enclosed syringe specifically designed for use with this product. Do not use any other dosing device.

**Other Information**

- store at 20-25°C (68-77°F)
- do not use if printed bottle wrap is broken or missing

### **Inactive ingredients**

anhydrous citric acid, butylparaben, calcium sulfate, carrageenan, D&C red #33, FD&C blue #1, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

### **Questions or comments?**

1-888-287-1915

### **Package/Label Principal Display Panel**

Compare to Infants' Tylenol® Oral Suspension active ingredient

infant's PAIN & FEVER

Acetaminophen 160 mg per 5 mL

Suspension Liquid

Pain Reliever/Fever Reducer

AGES 2 to 3 YEARS

Pediatrician preferred dosing system

Alcohol Free

Aspirin Free

Ibuprofen Free

Grape Flavored

1 FL OZ (30 mL)

Use only with enclosed syringe. See side panel for more information.



## EQUATE PAIN AND FEVER INFANTS

acetaminophen suspension

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49035-946
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	

BUTYL PARABEN (UNII: 3QPIIU3FV8)
CALCIUM SULFATE (UNII: WAT0DDB505)
CARRAGEENAN (UNII: 5C69YCD2YJ)
D&C RED NO. 33 (UNII: 9DBA0SBB0L)
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)
GLYCERIN (UNII: PDC6A3C0OX)
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
WATER (UNII: 059QF0KO0R)
SODIUM BENZOATE (UNII: OJ245FE5EU)
SORBITOL (UNII: 506T60A25R)
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)

### Product Characteristics

Color	PURPLE	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49035-946-10	1 in 1 CARTON	09/14/2011	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:49035-946-16	1 in 1 CARTON	08/08/2011	
2		59 mL in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	08/08/2011	

**Labeler** - Wal-Mart Stores Inc (051957769)