EQUATE PAIN AND FEVER INFANTS- acetaminophen suspension Wal-Mart Stores Inc

Wal-Mart Infants' Pain & Fever Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- new symptoms occur
- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise, use age.
- push air out of syringe. Insert syringe tip into bottle opening.
- flip bottle upside down. Pull yellow part of syringe to the first dose line and then push product back into bottle.
- · pull yellow part of syringe until it reaches and stays at the correct dose
- dispense liquid slowly into child's mouth, toward inner cheek
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- replace cap tightly to maintain child resistance

| Dosing | Chart |
|--------|-------|
| | |

| Weight (lb) | Age (yr) | Dose (mL)* | | | |
|-------------|---------------|--------------|--|--|--|
| under 24 | under 2 years | ask a doctor | | | |

| 24-35 | 2-3 years | 5 mL |
|-------|-----------|------|

*or as directed by a doctor

Attention: use only enclosed syringe specifically designed for use with this product. Do not use any other dosing device.

Other Information

- store at 20-25°C (68-77°F)
- do not use if printed neckband is broken or missing

Inactive ingredients

anhydrous citric acid, calcium sulfate, carrageenan, D&C red #33, FD&C blue #1, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions or comments?

1-888-287-1915

Package/Label Principal Display Panel

equate™

Compare to Infants' Tylenol® active ingredient

infants PAIN & FEVER

Acetaminophen 160 mg per 5 mL

Suspension Liquid

Pain Reliever/Fever Reducer

AGES 2 to 3 YEARS

Pediatrician preferred dosing system

- Alcohol-Free
- Aspirin-Free
- Ibuprofen-Free

PARABEN FREE

Grape Flavored

2 FL OZ (59 mL)

Use only with enclosed syringe. See side panel for more information.



EQUATE PAIN AND FEVER INFANTS

acetaminophen suspension

| Product Information | | | | |
|-------------------------|----------------|--------------------|---------------|--|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:49035-946 | |
| Route of Administration | ORAL | | | |

| Active Ingredient/Active Moiety | | | | |
|--|--------------------------|----------------|--|--|
| Ingredient Name | Basis of Strength | Strength | | |
| ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D) | ACETAMINOPHEN | 160 mg in 5 mL | | |

| Inactive Ingredients | | | | |
|---|----------|--|--|--|
| Ingredient Name | Strength | | | |
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | | | | |
| CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WATODDB505) | | | | |
| CARRAGEENAN (UNII: 5C69YCD2YJ) | | | | |
| D&C RED NO. 33 (UNII: 9DBA0SBB0L) | | | | |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | | | | |
| GLYCERIN (UNII: PDC6A3C0OX) | | | | |
| HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S) | | | | |
| HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D) | | | | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | | | | |
| CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K6790BS311) | | | | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | | | | |
| WATER (UNII: 059QF0KO0R) | | | | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | | | | |
| SORBITOL SOLUTION (UNII: 8KW3E207O2) | | | | |
| SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X) | | | | |

| Product Characteristics | | | | |
|-------------------------|--------|--------------|--|--|
| Color | PURPLE | Score | | |
| Shape | | Size | | |
| Flavor | GRAPE | Imprint Code | | |
| Contains | | | | |

| P | Packaging | | | | | |
|---|----------------------|--|-------------------------|-----------------------|--|--|
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date | | |
| 1 | NDC:49035-946- 10 | 1 in 1 CARTON | 09/14/2011 | 07/31/2021 | | |
| 1 | | 30 mL in 1 BOTTLE; Type 0: Not a Combination Product | | | | |
| 2 | NDC:49035-946- 16 | 1 in 1 CARTON | 08/08/2011 | | | |
| 2 | | 59 mL in 1 BOTTLE; Type 0: Not a Combination Product | | | | |

| Marketing Information | | | | |
|-----------------------|---|-------------------------|-----------------------|--|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | |
| OTC Monograph Drug | M013 | 08/08/2011 | | |
| | | | | |

Labeler - Wal-Mart Stores Inc (051957769)

Revised: 7/2024 Wal-Mart Stores Inc