# COLD- acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride tablet, coated Marc Glassman, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

#### 1138 - MAR - 2018-1211

#### **Drug Facts**

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

#### Uses

- temporarily relieves these common cold/flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - nasal congestion
  - cough
  - sinus congestion and pressure
- helps clear nasal passages
- promotes nasal and sinus drainage
- temporarily reduces fever

#### **Warnings**

# Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

# Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

## Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

### When using this product do not exceed recommended dosage

## Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

# Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### **Directions**

• do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul> <li>swallow whole - do not crush, chew, or dissolve</li> <li>do not take more than 10 caplets in 24 hours</li> </ul>
children under 12 years	ask a doctor

#### Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information

# **Inactive ingredients**

acesulfame potassium, colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

#### PRINCIPAL DISPLAY PANEL

See New Warnings Information & Directions

†Compare to the active ingredients in Tylenol® Cold Multi-Symptom Daytime

Marc's®

Daytime, Non-drowsy

Cold Multi-Symptom

For Adults

Pain Reliever/Fever Reducer, Nasal Decongestant, Cough Suppressant

Acetaminophen, Phenylephrine HCl, Dextromethorphan HBr

For Relief Of:

- Headache, Fever & Sore Throat
- Nasal Congestion
- Cough

24 Cool Taste Caplets

F113805MAR

- persistent or chronic cough such as occurs with smoking, asthma, m trouble uninating due to an enlarged prostate glass
  - liver disease Ask a doctor before use if you have
- y hon pane ener pad an allergic reaction to this product or any of its ingredients an MAOI, ask a doctor or pharmacist before taking this product. the MAOI drug. If you do not know if your prescription drug contains conditions, or Parkinson's disease), or for 2 weeks after stopping (MAOI) (certain drugs for depression, psychiatric, or emotional
- if you are now taking a prescription monoamine oxidase inhibitor acetaminophen, ask a doctor or pharmacist. uoubusecubnou): Il you are not sure whether a drug contains
  - with any other drug containing acetaminophen (prescription or esn jou og

vomiting, consult a doctor promptly.

- days, is accompanied or followed by fever, headache, rash, nausea, or Sore throat warning: If sore throat is severe, persists for more than 2
  - 3 or more alcoholic drinks every day while using this product миди одиек qunda confaining acetaminophen
    - more than 4,000 mg of acetaminophen in 24 hours 24 hours. Severe liver damage may occur if you take
- dose of this product is 10 caplets (3,250 mg acetaminophen) in Liver warning: This product contains acetaminophen. The maximum

nund racts (continued)

12 years children under ■ do not take more than 10 caplets in 24 hours years and over swellow whole – do not crush, chew, or dissolve children 12 ■ take 2 caplets every 4 hours bns stlubs do not take more than directed (see overdose warning)

Directions

attention is critical for adults as well as for children even if you do not Poison Control Center right away (1-800-222-1222). Quick medical Overdose warning: In case of overdose, get medical help or contact a keep out of reach of children.

If pregnant or breast-feeding, ask a health professional before use.

- These could be signs of a serious condition. conditionnes back or occurs with rash or headache that lasts ■ redness or swelling is present
  ■ new symptoms occur ■ Jever gets worse or lasts more than 3 days
- pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ nervousness, dizziness, or sleeplessness occur Stop use and ask a doctor if

When using this product do not exceed recommended dosage ninehew gunb gninnirð

Ask a doctor or pharmacist before use if you are taking the blood сопду для осспиг миду доо шпсу бунедци (шпспя)

nund racts (continued)

 pembocsuj
 , tequces perent pelbe cjest usesj basesôes

 bromotes nasal and sinus drainage a siuns coudestion and pressure ■ condy шодзабиоо језеи в

в рездесув mujuor aches and pains sone throat NS63 # femborarily relieves these common cold/flu symptoms:

nund Lacts (continued)

Masal decongestant Phenylephrine HCl 5 mg. guessauddns u6nog Dextromethorphan HBr 10 mg.

Active ingredients (in each caplet) Purpose

†Compare to the active

redients in Tylenol® Cold Multi-Symptom Daytime

See New Warnings Information & Directions



Marc's.



PAIN RELIEVER/FEVER REDUCER, NASAL DECONGESTANT, COUGH SUPPRESANT

Acetaminophen, Phenylephrine HCl, Dextromethorphan HBr

For Relief of:

- Headache, Fever & Sore Throat
- Nasal Congestion
- Cough

24 Cool Taste Caplets

silicon dioxide, com starch, croscarmellose sodium, crospovidone, >

 retain carton for complete product information store between 20-25°C (68-77°F) in a dry place

Other information

Drug Facts (continued)

0

IF BLISTER UNITS IN OR BROKEN †This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenof® Cold Mutit-Symptom Daysime. USE IF TORN ( NOT ARE

Distributed by: Marc Glassman, Inc. West 130th Street Cleveland, 0H 44130

acesultame potassium, colloidal inactive ingredients

stearic acid, talc, titanium dioxide glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol flavor, magnesium stearate, microcrystalline cellulose, polyethylene

Drug Facts (continued)

# **COLD**

acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68998-138
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	<b>Basis of Strength</b>	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg	
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

Inactive Ingredients	
Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)	
CROSPOVIDONE (UNII: 2S7830E561)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
POVIDONE (UNII: FZ 989GH94E)	
STARCH, CORN (UNII: 08232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics				
Color	white	Score	no score	
Shape	OVAL	Size	17mm	
Flavor	MINT	Imprint Code	AAA;1138	
Contains				

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:68998- 138-05	2 in 1 CARTON	02/01/2010	06/30/2024	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part341	02/01/2010	06/30/2024	

Labeler - Marc Glassman, Inc. (094487477)

Revised: 12/2022 Marc Glassman, Inc.