

**TUSSIN COUGH EXPECTORANT COUGH SUPPRESSANT DM- dextromethorphan hydrobromide liquid**

**AptaPharma Inc.**

-----  
**Drug Facts**

**Active ingredients (in each 5mL tsp)**

Dextromethorphan HBr, USP 10mg

Guaifenesin, USP 100mg

**Purpose**

Dextromethorphan HBr, USP .... Cough Suppressant

Guaifenesin, USP ..... Expectorant

**Uses**

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

**Warnings**

**Do not use** ■ in a child under 12 years of age  
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before using this product.

**Ask a doctor before use if you have**

- cough that occurs with too much phlegm (mucus)
- cough that is chronic, such as occurs with smoking, asthma, chronic bronchitis or emphysema

**Stop use and ask a doctor if** cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than 6 doses in any 24-hour period.

-----

**Age**

**Dose**

-----  
Adults and children                      2 teaspoonfuls  
12 years and over                        every 4 hours  
-----

children under 12 years                do not use  
-----

**Other information**

- store at 20-25°C (68-77°F) ■ do not refrigerate
- dosage cup provided ■ sodium 3 mg per teaspoonful
- **Keep carton for full directions for use.**

**Inactive ingredients** Anhydrous citric acid, dextrose, FD&C red #40, flavor, glycerin, high fructose corn syrup, menthol, purified water. saccharin sodium, sodium benzoate.

**Package Label**

**AP SAFE** NDC 76281-503-24

**COMPARE TO THE ACTIVE  
INGREDIENTS IN ROBITUSSIN®  
PEAK COLD COUGH & CHEST  
CONGESTION DM\***

**Adult  
Tussin  
Cough**

**Dextromethorphan HBr / Guaifenesin**

**Cough Suppressant / Expectorant**

**Non-Drowsy  
Cough Formula**

- **Controls Cough**
- **Helps Loosen chest congestion**
- **For Ages 12 and over**

**DM**

4 FL OZ (118 mL)

**CAUTION: DO NOT USE IF IMPRINTED SEAL UNDER  
CAP IS BROKEN OR MISSING.**

3  
1013430  
3/20

This product is not manufactured or distributed

by Pfizer, owner of the registered trademark Robitussin® Peak Cold.

**Manufactured by:**  
**AptaPharma Inc.,**  
**1533 Union Ave.**  
**Pennsauken, NJ 08110**

**AP-BX-02**

**Carton**



NDC 76281-503-24

COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD COUGH & CHEST CONGESTION DM\*

# Adult Tussin Cough

Dextromethorphan HBr / Guaifenesin Cough Suppressant / Expectorant

**Non-Drowsy Cough Formula**

- Controls Cough
- Helps Loosen chest congestion
- For Ages 12 and over

**DM**

4 FL OZ (118 mL)

3  
1013430  
3/20

CAUTION: DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING.

### Drug Facts

**Active ingredients (in each 5mL tsp)**  
 Dextromethorphan HBr, USP 10mg..... Cough Suppressant  
 Guaifenesin, USP 100mg..... Expectorant

**Purpose**  
**Uses**  
 ■ temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold  
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

**Warnings**  
**Do not use** ■ in a child under 12 years of age  
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before using this product.

**Ask a doctor before use if you have**  
 ■ cough that occurs with too much phlegm (mucus)  
 ■ cough that lasts or chronic, such as occurs with smoking, asthma, chronic bronchitis or emphysema

**Stop use and ask a doctor** if cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 ■ do not take more than 6 doses in any 24-hour period.

Age	Dose
Adults and children 12 years and over	2 teaspoonsfuls every 4 hours
children under 12 years	do not use ▶

### Drug Facts (continued)

#### Other information

■ store at 20-25°C (68-77°F) ■ do not refrigerate  
 ■ dosage cup provided ■ sodium 3 mg per teaspoonful  
 ■ **Keep carton for full directions for use.**

**Inactive ingredients** Anhydrous citric acid, dextrose, FD&C red #40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate.

**Questions?** Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-796-5944

\*This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin® Peak Cold.

**Manufactured by:**  
**AptaPharma Inc.,**  
**1533 Union Ave.**  
**Pennsauken, NJ 08110**



3 76281 50324 1 AP-BX-02

NDC 76281-503-24

COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD COUGH & CHEST CONGESTION DM\*

# Adult Tussin Cough

Dextromethorphan HBr / Guaifenesin Cough Suppressant / Expectorant

**Non-Drowsy Cough Formula**

- Controls Cough
- Helps Loosen chest congestion
- For Ages 12 and over

**DM**

4 FL OZ (118 mL)

NDC 76281-503-24

**CAUTION: DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING.**

**AP SAFE**

COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD COUGH & CHEST CONGESTION DM\*

# Adult Tussin Cough

Dextromethorphan HBr / Guaifenesin Cough Suppressant / Expectorant

**Non-Drowsy Cough Formula**

- Controls Cough
- Helps Loosen chest congestion
- For Ages 12 and over

**DM**

4 FL OZ (118 mL)

3  
1013430  
3/20

**Drug Facts**

Active ingredients (in each 5mL tsp)	Purpose
Dextromethorphan HBr, USP 10mg	Cough Suppressant
Guaifenesin, USP 100mg	Expectorant

**Uses**

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

**Warnings**

**Do not use** if in a child under 12 years of age

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before using this product.

**Ask a doctor before use if you have**

- cough that occurs with too much phlegm (mucus)
- cough that lasts or chronic, such as occurs with smoking, asthma, chronic bronchitis or emphysema

**Stop use and ask a doctor** if cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than 6 doses in any 24-hour period.

Age	Dose
Adults and children 12 years and over	2 teaspoonsful every 4 hours
children under 12 years	do not use

NDC 76281-503-24

**AP SAFE**

Adult Tussin Cough

Expectorant Cough Suppressant DM

COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD COUGH & CHEST CONGESTION DM\*

# Adult Tussin Cough

Dextromethorphan HBr / Guaifenesin Cough Suppressant / Expectorant

**Non-Drowsy Cough Formula**

- Controls Cough
- Helps Loosen chest congestion
- For Ages 12 and over

**DM**

4 FL OZ (118 mL)

**Drug Facts (continued)**

**Other information**

- store at 20-25°C (68-77°F)
- do not refrigerate
- dosage cup provided
- sodium 3 mg per teaspoonful
- Keep carton for full directions for use.

**Inactive ingredients** Anhydrous citric acid, dextrose, FD&C red #40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate.

**Questions?** Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-796-5944

\*This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin® Peak Cold.

**Manufactured by:**  
AptaPharma Inc.,  
1533 Union Ave.  
Pennsauken, NJ 08110

3 7 6 2 8 1 5 0 3 2 4 1 AP-BX-02

**Bottle**



NDC 76281-503-24

# Adult Tussin Cough

Dextromethorphan HBr / Guaifenesin  
Cough Suppressant / Expectorant

**Non-Drowsy  
Cough Formula**

- Controls Cough
- Helps Loosen chest congestion
- For Ages 12 and over

4 FL OZ (118 mL)



DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING.

## Drug Facts

Active ingredients (in each 5mL tsp)	Purpose
Dextromethorphan HBr, USP 10mg	Cough Suppressant
Guaifenesin, USP 100mg	Expectorant

**Uses** ■ temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold ■ helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

### Warnings

**Do not use** ■ in a child under 12 years of age ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before using this product.

**Ask a doctor before use if you have** ■ cough that occurs with too much phlegm (mucus) ■ cough that lasts or chronic, such as occurs with smoking, asthma, chronic bronchitis or emphysema

**Stop use and ask a doctor if** cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ do not take more than 6 doses in any 24-hour period.

Age	Dose
Adults and children 12 years and over	2 teaspoonfuls every 4 hours
children under 12 years	do not use

**Other information** ■ store at 20-25°C (68-77°F) ■ do not refrigerate  
■ dosage cup provided ■ sodium 3 mg per teaspoonful  
■ **Keep carton for full directions for use.**

**Inactive ingredients** Anhydrous citric acid, dextrose, FD&C red #40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate.

**Questions?** Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

Manufactured by: Aptapharma Inc.,  
1533 Union Ave. Pennsauken, NJ 08110

AP-LR-02

LOT:

EXP:



NDC 76281-503-24

Adult  
**Tussin  
Cough**  
Dextromethorphan HBr / Guaifenesin  
Cough Suppressant / Expectorant

**Non-Drowsy  
Cough Formula**

- Controls Cough
- Helps Loosen chest congestion
- For Ages 12 and over

4 FL OZ (118 mL)



DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING.

**Drug Facts**

Active ingredients (in each 5mL tsp)	Purpose
Dextromethorphan HBr, USP 10mg	Cough Suppressant
Guaifenesin, USP 100mg	Expectorant

**Uses** ■ temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold ■ helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

**Warnings**

**Do not use** ■ in a child under 12 years of age ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before using this product.

**Ask a doctor before use if you have** ■ cough that occurs with too much phlegm (mucus) ■ cough that lasts or chronic, such as occurs with smoking, asthma, chronic bronchitis or emphysema

**Stop use and ask a doctor if** cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ do not take more than 6 doses in any 24-hour period.

Age	Dose
Adults and children 12 years and over	2 teaspoonfuls every 4 hours
children under 12 years	do not use

**Other information** ■ store at 20-25°C (68-77°F) ■ do not refrigerate ■ dosage cup provided ■ sodium 3 mg per teaspoonful ■ **Keep carton for full directions for use.**

**Inactive ingredients** Anhydrous citric acid, dextrose, FD&C red #40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate.

**Questions?** Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

Manufactured by: Aptapharma Inc.,  
1533 Union Ave. Pennsauken, NJ 08110

AP-LR-02

LOT:

EXP:

res

**TUSSIN COUGH EXPECTORANT COUGH SUPPRESSANT DM**  
dextromethorphan hydrobromide liquid

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:76281-503
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 5 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>DEXTROSE</b> (UNII: IY9XDZ35W2)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>HIGH FRUCTOSE CORN SYRUP</b> (UNII: XY6UN3QB6S)	
<b>MENTHOL</b> (UNII: L7T10EIP3A)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	

## Product Characteristics

<b>Color</b>	red	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	FRUIT PUNCH	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:76281-503-24	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/30/2020	

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M012	09/30/2020	

**Labeler** - AptaPharma Inc. (790523323)

**Registrant** - AptaPharma Inc. (790523323)

**Establishment**

Name	Address	ID/FEI	Business Operations
AptaPharma Inc.		790523323	manufacture(76281-503)

Revised: 12/2023

AptaPharma Inc.