

**NASAL DECONGESTANT MAXIMUM STRENGTH NON DROWSY-  
pseudoephedrine hcl tablet, film coated  
Meijer Distribution Inc**

-----  
**Meijer Distribution, Inc. Nasal Decongestant Drug Facts**

**Active ingredient (in each tablet)**

Pseudoephedrine HCl 30 mg

**Purpose**

Nasal decongestant

**Uses**

- temporarily relieves sinus congestion and pressure
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies

**Warnings**

**Do not use**

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

**When using this product**

**do not exceed recommended dosage**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever

**If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

adults and children 12 years and over	<ul style="list-style-type: none"><li>• take 2 tablets every 4 to 6 hours</li><li>• do not take more than 8 tablets in 24 hours</li></ul>
children ages 6 to 11 years	<ul style="list-style-type: none"><li>• take 1 tablet every 4 to 6 hours</li><li>• do not take more than 4 tablets in 24 hours</li></ul>
children under 6 years	do not use this product in children under 6 years of age

### **Other information**

- **each tablet contains:** calcium 20 mg
- store at 20°-25° C (68°-77° F)
- **do not use if blister unit is broken or torn**

### **Inactive ingredients**

carnauba wax, dibasic calcium phosphate dihydrate, FD&C red no. 40 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, silicon dioxide, titanium dioxide

### **Questions or comments?**

**1-800-719-9260**

### **Principal Display Panel**

meijer®

MAXIMUM STRENGTH

Nasal Decongestant

COMPARE TO SUDAFED® CONGESTION ACTIVE INGREDIENT

PSEUDOEPHEDRINE HCl TABLETS, 30 mg

(NASAL DECONGESTANT)

Non-Drowsy

Sinus Pressure + Congestion

24 TABLETS

Actual Size



43262 0E C12

Important: Read all product information before using. Keep this box for important information.

**Drug Facts**

**Active ingredient (in each tablet) Purpose**  
Pseudoephedrine HCl 30 mg..... Nasal decongestant

**Uses**

- temporarily relieves sinus congestion and pressure
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies

**Warnings**

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

When using this product do not exceed recommended dosage

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Drug Facts (continued)****Directions**

- |                                       |  |
|---------------------------------------|--|
| adults and children 12 years and over | <ul style="list-style-type: none"> <li>■ take 2 tablets every 4 to 6 hours</li> <li>■ do not take more than 8 tablets in 24 hours</li> </ul> |
| children ages 6 to 11 years           | <ul style="list-style-type: none"> <li>■ take 1 tablet every 4 to 6 hours</li> <li>■ do not take more than 4 tablets in 24 hours</li> </ul>  |
| children under 6 years                | do not use this product in children under 6 years of age   |

**Other information**

- each tablet contains: calcium 20 mg
- store at 20°-25°C (68°-77°F)
- do not use if blister unit is broken or torn

**Inactive ingredients** carnauba wax, dibasic calcium phosphate dihydrate, FD&C red no. 40 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, silicon dioxide, titanium dioxide

**Questions or comments? 1-800-719-9260**

\*This product is not manufactured or distributed by Johnson & Johnson, owner of the registered trademark Sudafed®.

DIST. BY MEIJER DISTRIBUTION, INC.  
GRAND RAPIDS, MI 49544  
www.meijer.com

**NASAL DECONGESTANT MAXIMUM STRENGTH NON DROWSY**

pseudoephedrine hcl tablet, film coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41250-432
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>PSEUDOEPHEDRINE HYDROCHLORIDE</b> (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>DIBASIC CALCIUM PHOSPHATE DIHYDRATE</b> (UNII: O7TSZ97GEP)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6130)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	

<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	

### Product Characteristics

<b>Color</b>	RED	<b>Score</b>	no score
<b>Shape</b>	ROUND (convex)	<b>Size</b>	7mm
<b>Flavor</b>		<b>Imprint Code</b>	L432
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-432-62	24 in 1 CARTON	02/15/1991	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:41250-432-67	48 in 1 CARTON	11/09/1995	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:41250-432-80	96 in 1 CARTON	08/14/1997	
3		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	02/15/1991	

**Labeler** - Meijer Distribution Inc (006959555)

Revised: 9/2024

Meijer Distribution Inc