

SOUND BODY COLD AND FLU RELIEF- acetaminophen, dextromethorphan hbr, doxylamine succinate solution

Big Lots Stores, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Big Lots, Inc Cold & Flu Relief Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening

- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be

signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 30 mL contains:** sodium 38 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

alcohol, anhydrous citric acid, D&C yellow no. 10, FD&C green no. 3, FD&C yellow no. 6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions?

1-800-719-9260

Package/Label Principal Display Panel

SOUNDBODY™

Compare to the Active Ingredients in Vicks® NyQuil® Cold & Flu

NIGHTTIME

Cold & Flu Relief

Pain Reliever / Fever Reducer – Acetaminophen

Cough Suppressant – Dextromethorphan HBr

Antihistamine - Doxylamine Succinate

10% ALCOHOL

ORIGINAL FLAVOR

12 FL OZ (355 mL)



Packaged by
Perrigo Company for:
Big Lots Stores, Inc.
P.O. Box 28523
Columbus, OH 43228-0523
V#5003185 ITEM#33540

PARENTS: **DO NOT USE IF PRINTED**
Learn about teen medicine abuse **HECKBAND IS BROKEN**
OR **MISSING**
www.StopMedicineAbuse.org

Drug Facts
Active ingredients Purpose
(in each 30 mL)
Acetaminophen Pain reliever/
650 mg fever reducer
Dextromethorphan hbr
30 mg Cough suppressant
Doxylamine succinate
12.5 mg Antihistamine

Uses temporarily relieves common cold/flu symptoms: cough due to minor throat and bronchial irritation sore throat headache minor aches and pains fever
nausea and vomiting

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include skin redness blisters rash
If a skin reaction occurs, stop use and seek medical help right away.
See throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, noise, or swelling, consult a doctor promptly.

7003001982

PEEL BACK AT CORNER FOR MORE INFORMATION : 33540 TH 81

Drug Facts
(continued)

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. If you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have:
liver disease glaucoma
cough that occurs with too much phlegm (mucus) a breathing problem such as emphysema or chronic bronchitis trouble urinating due to an enlarged prostate gland persistent or chronic cough that occurs with smoking, asthma, or emphysema
a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers taking the blood thinning drug warfarin

When using this product:
excitability may occur, especially in children marked drowsiness may occur
avoid alcoholic drinks

ADHESIVE AREA • NO COATING • NO VARNISH • NO TYPE

Drug Facts
(continued)

Be careful when driving a motor vehicle or operating machinery alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if:
pain or cough gets worse or lasts more than 7 days
fever gets worse or lasts more than 3 days
headache or swelling is present
new symptoms occur
cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults, as well as for children even if you do not notice any signs or symptoms.

Directions take only as directed - see Overdose warning only use the dose cup provided do not exceed 4 doses per 24 hrs
adults & children 12 yrs & over 30 mL every 6 hrs
children 4 to under 12 yrs ask a doctor
children under 4 yrs do not use

Other information each 30 mL contains sodium 38 mg store at 20-25°C (68-77°F)

Inactive ingredients alcohol, anhydrous citric acid, D&C yellow no. 10, FD&C green no. 3, FD&C yellow no. 6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions? 1-800-719-6280

GLUTEN FREE
*This product is not manufactured or distributed by Prechter & Gamble, distributor of Nicotrol Niquan Cold & Flu.

ADHESIVE AREA • NO COATING • NO VARNISH • NO TYPE

SOUND BODY COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50594-335
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C GREEN NO. 3 (UNII: 3P3ONR601S)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	

Product Characteristics

Color	GREEN (clear, bright green)	Score	
Shape		Size	
Flavor	FRUIT (anise / cooling menthol aroma)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50594-335-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/04/2015	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	03/04/2015	

Labeler - Big Lots Stores, Inc. (017885351)

Revised: 5/2023

Big Lots Stores, Inc.