

GOOD SENSE PAIN RELIEF- acetaminophen tablet

L. Perrigo Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Perrigo Pain Relief Tablets Drug Facts

Active ingredient (in each tablet)

Acetaminophen 325 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- the common cold
- headache
- minor pain of arthritis
- backache
- muscular aches
- toothache
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if the user has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if the user

- has liver disease
- is a child with pain of arthritis

Ask a doctor or pharmacist before use if the user is

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- new symptoms occur
- pain gets worse or lasts for more than 10 days (for adults) or 5 days (for children)
- fever gets worse or lasts for more than 3 days
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none"> • take 2 tablets every 4 to 6 hours while symptoms last • do not take more than 10 tablets in 24 hours • do not use for more than 10 days unless directed by a doctor
children 6-11 years	<ul style="list-style-type: none"> • take 1 tablet every 4 to 6 hours while symptoms last • do not take more than 5 tablets in 24 hours • do not use for more than 5 days unless directed by a doctor
children under 6 years	ask a doctor

Inactive ingredients

croscarmellose sodium*, povidone, pregelatinized starch, stearic acid *may contain this ingredient

Questions or comments?

1-800-719-9260

Principal Display Panel

Regular Strength

Pain Relief TABLETS

Pain Reliever/Fever Reducer

Acetaminophen

Compare to active ingredient of Tylenol® Tablets

100% SATISFACTION GUARANTEED

100 Tablets - 325 mg Each

GOODSENSE.

NDC 0113-0403-78

Regular Strength

Pain Relief

Pain Reliever/Fever Reducer
Acetaminophen

TABLETS



100 Tablets - 325 mg Each

Compare to active ingredient of
Tylenol® Tablets



GOODSENSE.

NDC 0113-0403-78

Regular Strength

Pain Relief

Pain Reliever/Fever Reducer
Acetaminophen

TABLETS



100 Tablets - 325 mg Each

Compare to active ingredient of
Tylenol® Tablets



EXP.

LOT NO.

70030 13032 0



Drug Facts

Active ingredient *Purpose*
(in each tablet)

Acetaminophen 325 mg...Pain reliever/fever reducer

Uses ■ temporarily relieves minor aches and pains due to:
■ the common cold ■ headache
■ minor pain of arthritis ■ backache
■ muscular aches ■ toothache
■ premenstrual and menstrual cramps
■ temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if
■ adult takes more than 4,000 mg of acetaminophen in 24 hours
■ child takes more than 5 doses in 24 hours
■ taken with other drugs containing acetaminophen
■ adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if the user has ever had an allergic reaction to this product or any of its ingredients

Drug Facts (continued)

Ask a doctor before use if the user
■ has liver disease ■ is a child with pain of arthritis
Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if ■ new symptoms occur
■ pain gets worse or lasts more than 10 days (for adults) or 5 days (for children)
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. **Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions


■ do not take more than directed (see overdose warning)

adults and children 12 years and over
■ take 2 tablets every 4 to 6 hours while symptoms last
■ do not take more than 10 tablets in 24 hours
■ do not use for more than 10 days unless directed by a doctor

children 6-11 years
■ take 1 tablet every 4 to 6 hours while symptoms last
■ do not take more than 5 tablets in 24 hours
■ do not use for more than 5 days unless directed by a doctor

children under 6 years
ask a doctor

Drug Facts
Inactive
sodium
acid
free
Quest
1-800-71
Store at 20
DO NOT USE
UNDER CA
Distribute
Per
Allergan, Inc



GOOD SENSE PAIN RELIEF

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0113-0403
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	WHITE	Score	no score
Shape	ROUND (beveled edge)	Size	10mm
Flavor		Imprint Code	325MG;L403
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0113-0403-78	1 in 1 CARTON	08/15/1989	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	08/15/1989	

