LOW DOSE ASPIRIN- aspirin tablet, delayed release CVS WOONSOCKET PRESCRIPTION CENTER, INCORPORATED

CVS 44-645

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID*)
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- have 3 or more alcoholic drinks every day while using this product
- have had stomach ulcers or bleeding problems
- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Do not use

if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if

stomach bleeding warning applies to you

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

colloidal anhydrous silica, corn starch, FD&C red #40, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-426-9391

Principal Display Panel

CVSTM

VALUE

SIZE

LOW DOSE

ASPIRIN 81 mg Pain Reliever (**NSAID**)

NDC 51316-456-51

Compare to St. Joseph® Safety Coated 81mg Low Dose Aspirin active ingredient†

[‡]Talk to your doctor or other healthcare provider before using this product for your heart.

365 ENTERIC COATED TABLETS

Safety Coated

Aspirin

Regimen[‡]

actual size

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

[†]This product is not manufactured or distributed by Foundation Consumer Healthcare, LLC, owner of the registered trademark St. Joseph® Safety Coated 81 mg Low Dose Aspirin. 50844 ORG102464551

Distributed by: CVS Pharmacy, Inc.

One CVS Drive Woonsocket, RI 02895 © 2025 CVS/pharmacy CVS.com® 1-800-SHOP CVS V-11112



CVS 44-645

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away

by a doctor because it may cause problems in the unborn

child or complications during delivery.

adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless

drink a full glass of water with each dose

do not take more than directed

Directions

children under 12 years: do not use unless directed by a

directed by a doctor

Other information

doctor

colloidal anhydrous silica, corn starch, FD&C red #40, FD&C

Inactive ingredients

yellow #6, hypromellose, methacrylic acid, microcrystalline

cellulose, polydextrose, polyethylene glycol, shellac wax,

Questions or comments? 1-800-426-9391

STOP PEELING

itanium dioxide, triacetin, triethyl citrate

LOW DOSE ASPIRIN

Do not use if you have ever had an allergic reaction to any

Drug Facts (continued)

other pain reliever/fever reduce

Ask a doctor before use if

Ask a doctor or pharmacist before use if you are taking a

■ you are taking a diuretic
■ you have asthma

or kidney disease

prescription drug for diabetes gout arthritis
Stop use and ask a doctor if an allergic reaction occurs. Seek medical help right away

you experience any of the following signs of stomach bleeding: ■ feel faint ■ womit blood

have bloody or black stools

■ ringing in the ears or a loss of hearing occurs ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days have stomach pain that does not get better

redness or swelling is present

any new symptoms appear

aspirin tablet, delayed release

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51316-456
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	AS PIRIN	81 mg

	Inactive Ingredients	
ı	Ingredient Name	Strength

SILICON DIOXIDE (UNII: ETJ7Z6XBU4) STARCH, CORN (UNII: O8232NY3SJ) FD&C RED NO. 40 (UNII: WZB9127XOA) FD&C YELLOW NO. 6 (UNII: H77VEI93A8) HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO) METHACRYLIC ACID (UNII: 1CS02G8656) MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) POLYDEXTROSE (UNII: VH2XOU12IE) POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) **SHELLAC** (UNII: 46N107B710) **DIMETHICONE** (UNII: 92RU3N3Y10) WATER (UNII: 059QF0KO0R) **SODIUM BICARBONATE** (UNII: 8MDF5V39QO) **SODIUM LAURYL SULFATE** (UNII: 368GB5141J) TALC (UNII: 7SEV7J4R1U) **TITANIUM DIOXIDE** (UNII: 15FIX9V2JP) TRIACETIN (UNII: XHX3C3X673) TRIETHYL CITRATE (UNII: 8Z96QXD6UM)

Product Characteristics					
Color	pink	Score	no score		
Shape	ROUND	Size	6mm		
Flavor		Imprint Code	L		
Contains					

l	Packaging					
	#	t Item Code Package Description		Marketing Start Date	Marketing End Date	
		NDC:51316- 456-51	365 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/08/2027		

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC Monograph Drug	M013	10/02/2025			

Labeler - CVS WOONSOCKET PRESCRIPTION CENTER, INCORPORATED (062312574)

Establishment					
Name	Address	ID/FEI	Business Operations		
LNK International, Inc.		038154464	pack(51316-456)		

Establishment			
Name	Address	ID/FEI	Business Operations

LNK International, Inc.	832867837	manufacture(51316-456)	
-------------------------	-----------	------------------------	--

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(51316-456)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(51316-456)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(51316-456)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(51316-456)

Revised: 11/2025 CVS WOONSOCKET PRESCRIPTION CENTER, INCORPORATED